


Without the contributions of Mr. Hart Brown and D. H. Shurbet and his students at Texas Tech University this map would not have been possible.




Prepared by the U.S. Army Topographic Command (BEPM), Washington, D.C. Compiled in 1956 by photogrammetric methods from aerial photographs taken 1954. Photographs field annotated 1954. Revised in 1975 by the U.S. Geological Survey from aerial photographs taken 1974.

REFERENCES

- Briggs, I. C., 1974, Machine contouring using minimum curvature: *Geophysics*, v. 38, p. 39-48.
Hammer, S., 1939, Terrain corrections for gravimeter stations: *Geophysics*, v. 4, p. 184-194.
Morelli, G., 1967, Geodetic standards for gravity surveys: *Geophysics*, v. 41, p. 1051.
Plouff, D., 1977, Preliminary documentation for a FORTRAN program to compute gravity terrain corrections based on topography digitized on a geographic grid: U.S. Geological Survey, Open-File Report 7-353, 45 p.
Sampson, R. J., 1978, Surface II graphics system: Kansas Geological Survey, 240 p.

10,000-meter Universal Transverse Mercator grid ticks, zone 13
+ = approximate location of station points


Major Sources of Data for the Perryton Sheet:
Defense Meteorological Project for the Environment, Dallas, The University of Texas at El Paso, Texas Tech University, Mr. Hart Brown, Shurbet, D. H., 1968, Gravity field and isostatic equilibrium of the Llano Estacado of Texas and New Mexico: *Geol. Soc. Am. Bull.*, v. 79, p. 213-222.



BOUGUER GRAVITY ATLAS OF TEXAS, PERRYTON SHEET

Compiled by: G. R. Keller and C. L. V. Aiken
Assisted by: M. R. Voight and Wayne Basden

1986



CARTOGRAPHY BY RICHARD M. PLATT UNDER THE SUPERVISION OF R. L. DILLON.

1986 MAGNETIC DECLINATION FOR CENTER OF THIS SHEET IS 8°58' EAST. MEAN ANNUAL CHANGE IS 45° WESTWARD.

CONTOUR INTERVAL = 2 MILLIGALS

TEXAS GRAVITY DATA BASE AND REDUCTION PARAMETERS

The data base of gravity measurements made in Texas gravity mapping project is the result of over 100,000 individual stations and many will contain more than 50,000 readings. Major sources of data for each sheet are indicated in the explanation. However, a particular note of thanks is due to Mr. and Mrs. Hart Brown because the regional data they so generously provided is the foundation of this project. Without these data, the mapping and compilation required would have been impossible.

Data for the Texas gravity data base are catalogued and stored in a standard format and are available to the public (Hammer, 1967; Morelli, 1976). Several forms were used as an elevation datum, and a density of 2.67 g/cm³ was used in the Bouguer correction. For stations west of 103° W longitude, outer-zone terrain corrections were calculated for zones extending from 0.89 km to 167 km from each station (Hammer, 1969). These calculations employed a terrain correction program written by Plouff (1977) and a set of average elevations on a grid interval of 30 km of latitude and longitude.

The data base is information contained in the maps, maps all boundaries at common boundaries. These maps were constructed by gridding the data using the minimum curvature technique (Briggs, 1974) and were contoured by using a modified version of the Surface II graphics system (Sampson, 1978).

All gravity data were reduced to Bouguer anomaly values using the following formulas:

$$BA = G_{TH} - G_{TM} + C_{TA} + (C_{GEO} + C_{TERR}) + C_{CURV}$$

where

$$BA = \text{Bouguer anomaly}$$

ρ' = Ratio of reduction density to the standard value of 2.67 g/cm³

G_{TH} = Observed gravity, in milligals, relative to the IGSN-71 gravity datum (Morelli, 1976)

G_{TM} = Theoretical gravity on the surface (sea level) of the 1967 reference spheroid:
$$g_{TM} = 97903.843 + \sigma (15727.66 + \phi (-15762.337 + \sigma (6083.534 + \sigma (-1089.748 + \sigma (69.43))))), \sigma = 0.0001 \phi^2, \phi = \text{latitude in degrees}$$
 (International Association of Geodesy, 1967; see Morelli, 1976)

C_{TA} = Free-air correction for the elevation of the station relative to sea level:
$$C_{TA} = -0.0303 \phi^2 + \sigma (-0.0013596 + \sigma (0.0013553 + \sigma (-0.000529 + \sigma (0.000911))))$$

 $\phi = \text{elevation of station in meters}$

C_{GEO} = Bouguer correction for rock mass positioned between the station and sea level. Correction is calculated from the formula for the attraction of an infinite horizontal slab of thickness h and density ρ to obtain the equation $C_{GEO} = -0.119h$

C_{CURV} = Curvature correction, a modification of the Bouguer slab approximation that corrects for the curvature of the Earth's surface:
$$C_{CURV} = -h (1.4639108 \times 10^{-7} + h (-3.532715 \times 10^{-7} + h (4.449648 \times 10^{-8})))$$

C_{TERR} = Correction for local topography