





Photography used in the study was 1-m-pixel digital imagery, photographed in 2012, that was obtained from Texas Natural Resources Information System (TNRIS). A 1-m-resolution digital-elevation model (DEM) constructed from a 2014 airborne-lidar survey was also used. Previous regional maps that cover this area include the 1:125,000-scale Environmental Geologic Atlas, Corpus Christi Sheet (Brown and others, 1976), the 1:125,000-scale map of Distribution of Wetlands and Benthic Macroinvertebrates (White and others, 1983), and the 1:250,000-scale Geologic Atlas of Texas, Beeville–Bay City Sheet (Aronow, 1975; revised 1987). Sand-rich facies, Qbsd, of the Beaumont Formation are adapted from review of 1:250,000-scale interpretations of the Beeville–Bay City Sheet (Aronow, 1975; revised 1987), study of soil maps (Guckian and Gacia, 1979; Guckian and others, 1980), field observations, and ground-based electromagnetic conductivity measurements measured for this study. The 1:250,000-scale interpretations of the Beeville–Bay City Sheet (Aronow, 1975; revised 1987) are displayed in digital data files developed by the Texas Water Development Board and U.S. Geological Survey as part of the 1:250,000-scale Statewide Digital Geologic Atlas of Texas created from hard-copy maps. Topographic scarps were mapped from the 2014 lidar-derived DEM. Review of Paine (1991) provided useful subsurface information for the Quaternary deposits. Digital files for topography adapted to this map came from TNRIS for U.S. Geological Survey topographic quadrangle maps, Bayside and Mission Bay. Digital files of roads and railroads were also obtained through TNRIS. Water bodies were adapted from aerial photographs, lidar-derived DEMs, and the U.S. Geological Survey topographic maps. BEG staff Kutalnis Saylam, Tiffany Caudle, Aaron Averett, Ruth Costley, John Andrews, and John Hupp acquired and processed airborne lidar data. Work for this map was supported partly by the U.S. Geological Survey, National Geologic Mapping Program, with the assistance of STATEMAP award G13AC00178, and partly by the BEG STARR funds for geologic mapping and study of geologic hazards. Views and conclusions contained in this map should not be interpreted as necessarily representing the official policies, either expressed or implied, of the U.S. Government. The authors disclaim any responsibility or liability for interpretations from this map or digital data or decisions based thereon.


Edward W. Collins, Project Coordinator
Graphics by Jana S. Robinson
Editing by Chris Parker

Selected References

- Aronow, S., Brown, T. E., Brewton, J. L., Eargle, D. H., and Barnes, V. E., 1975, revised 1987, Beeville – Bay City Sheet: The University of Texas at Austin, Bureau of Economic Geology, Geologic Atlas of Texas, scale 1:250,000.
- Brown, L. F., Jr., Brewton, J. L., McGowan, J. H., Evans, T. H., Fisher, W. L., and Groat, C. G., 1976, Environmental geologic atlas of the Texas Coastal Zone: Corpus Christi area. The University of Texas at Austin, Bureau of Economic Geology, Geologic Atlas of Texas, scales 1:250,000 and 1:125,000, 123 p.
- Guckian, W. J. and Garcia, R. N., 1979, Soil survey of San Patricio and Aransas Counties, Texas: U.S. Department of Agriculture, Soil Conservation Service, 122 p.
- Guckian, W. J., Molina, Ramiro, and Divin, J. A., 1988, Soil Survey of Refugio County, Texas: U.S. Department of Agriculture, Soil Conservation Service, 123 p.
- McNeill, J. D., 1980a, Electrical Conductivity of Soil and Rocks: Geonics Limited, Mississauga, Ontario, Canada, Technical Note TN-5, 22 p.
- McNeill, J. D., 1980b, Electromagnetic Terrain Conductivity Measurement at Low Induction Numbers: Geonics Limited, Mississauga, Ontario, Canada, Technical Note TN-6, 15 p.
- Paine, J. G., 1991, Late Quaternary depositional units, sea level, and vertical movement along the central Texas coast: The University of Texas at Austin, Ph.D. dissertation, 256 p.
- White, W. A., Calnan, T. R., Morton, R. A., Kimble, R. S., Littleton, T. G., McGowan, J. H., Nance, H. S., Schmedes, K. E., and others, 1983, Submerged lands of Texas, Port Lavaca area: sediments, geochemistry, benthic macroinvertebrates, and associated wetlands: The University of Texas at Austin, Bureau of Economic Geology, 154 p.

GEOLOGIC MAP OF THE MISSION BAY QUADRANGLE: MISSION DELTA AND COPANO BAY AREA TEXAS GULF OF MEXICO COAST

Jeffrey G. Paine and Edward W. Collins
2014



Subsurface depiction of units diagrammatic following local detailed borehole data from Paine (1991).
Subsurface Qb contains local, undivided, sand-rich intervals.