





References

Bishop, B. A., 1967, Stratigraphic study of the Kiamichi Formation of the lower Cretaceous of Texas, in Hendricks, L., ed., Comanchean (Lower Cretaceous) stratigraphy and paleontology of Texas: The Permian Basin Section, Society of Economic Paleontologists and Mineralogists, Publication No. 67-8, p. 159-180.

Bybee, H. P., and Bullard, F. M., 1927, The geology of Cooke County, Texas: University of Texas, Austin, Bulletin No. 2710, plate scale 1:250,000, text 170 p.

Fisher, W. L., and Rodda, P. U., 1966, Nomenclature revision of basal Cretaceous rocks between the Colorado and Red Rivers, Texas: The University of Texas at Austin, Bureau of Economic Geology Report of Investigations No. 58, 20 p.

1967, Lower Cretaceous sands of Texas: stratigraphy and resources: The University of Texas at Austin, Bureau of Economic Geology Report of Investigations No. 59, 116 p.

Hendricks, L., 1967, Comanchean stratigraphy of the Cretaceous of north Texas, in Hendricks, L., ed., Comanchean (Lower Cretaceous) stratigraphy and paleontology of Texas: The Permian Basin Section, Society of Economic Paleontologists and Mineralogists, Publication No. 67-8, p. 51-63.

Hill, R. T., 1901, Geography and geology of the Black and Grand prairies, Texas, with detailed descriptions of the Cretaceous formations and special reference to artesian waters: U.S. Geological Survey, 21st Annual Report of the United States Geological Survey, to the Secretary of the Interior, 666 p.

McFarlan, E., Jr., and Menes, L. S., 1991, Lower Cretaceous, in Salvador, A., ed., The Gulf of Mexico Basin: Geological Society of America, The Geology of North America, v. J, p. 181-204.

McGowan, J. H., Hentz, T. F., Owen, D. E., Pieper, M. K., Shelby, C. A., and Barnes, V. E., 1967, Sherman sheet: The University of Texas at Austin, Bureau of Economic Geology, Geologic Atlas of Texas, scale 1:250,000, revised 1991, reprinted 1997.

Scott, R. W., Fee, D., Magee, R., and Laali, H., 1978, Episodic depositional models for the lower Cretaceous Washita Group, North-Central Texas: The University of Texas at Austin, Bureau of Economic Geology Report of Investigations No. 84, 23 p.

Stock, S. F., 1967, Physical stratigraphy of the Georgetown limestone equivalents in Tarrant, Denton, and Cooke Counties, Texas, in Hendricks, L., ed., Comanchean (lower Cretaceous) stratigraphy and paleontology of Texas: The Permian Basin Section, Society of Economic Paleontologists and Mineralogists, Publication No. 67-8, p. 183-216.

Geology illustrated on this map is based on field and aerial-photograph interpretations following a review of previous work on the area's geology. Previous regional maps include the 1:250,000-scale Sherman sheet (McGowan and others, 1967) and the 1:250,000-scale geologic map of Cooke County (Bybee and Bullard, 1927). The topographic base was created from digital files of the Texas Natural Resources Information Services (TNRIS) for U.S. Geological Survey, Muenster West, Texas, 7.5 minute topographic quadrangle map. Digital files of roads, railroads, county boundaries, drainage, and lakes were also obtained through TNRIS.

Work on this map was supported by the Bureau of Economic Geology STARR program. The mapping also benefited from ongoing STATEMAP mapping in North-Central Texas and from subsurface data from the Texas Railroad Commission Groundwater Advisory Unit. Editing was by Joan Maltese. Computer graphics work done to construct this map was by Cathy Brown. The author disclaims any responsibility or liability for interpretations from this map.

GEOLOGIC MAP OF THE MUENSTER WEST QUADRANGLE, TEXAS

Edward W. Collins

2015

