THE UNIVERSITY OF TEXAS BULLETIN

No. 3042: November 8, 1930

# THE GEOLOGY OF THE GLASS MOUNTAINS, TEXAS

## PART II

### FAUNAL SUMMARY AND CORRELATION OF THE PERMIAN FORMATIONS WITH DESCRIPTION OF BRACHIOPODA

By

ROBERT EVANS KING

Bureau of Economic Geology

J. A. Udden, Director E. H. Sellards, Associate Director

PUBLISHED BY

THE UNIVERSITY OF TEXAS

AUSTIN

QAe3042

THE UNIVERSITY OF TEXAS BULLETIN
No. 3042: November 8, 1930

# THE GEOLOGY OF THE GLASS MOUNTAINS, TEXAS

## PART II

FAUNAL SUMMARY AND CORRELATION OF THE PERMIAN FORMATIONS WITH DESCRIPTION OF BRACHIOPODA

By

ROBERT EVANS KING

Bureau of Economic Geology

J. A. Udden, Director E. H. Sellards, Associate Director

PUBLISHED BY THE UNIVERSITY FOUR TIMES A MONTH, AND ENTERED AS SECOND-CLASS MATTER AT THE POSTOFFICE AT AUSTIN, TEXAS, UNDER THE ACT OF AUGUST 24, 1912

The benefits of education and of useful knowledge, generally diffused through a community, are essential to the preservation of a free government.

Sam Houston

Cultivated mind is the guardian genius of Democracy, and while guided and controlled by virtue, the noblest attribute of man. It is the only dictator that freemen acknowledge, and the only security which freemen desire.

Mirabeau B. Lamar

# CONTENTS


| | Page |
|---|---|
| Acknowledgments | 5 |
| Historical resumé | 5 |
| Stratigraphy and correlation | 6 |
| Summary of the Permian stratigraphy of Trans-Pecos Texas | 6 |
| The Glass Mountains section | 6 |
| The Delaware and Guadalupe Mountains section | 11 |
| The Sierra Diablo section | 14 |
| Permian of the Diablo Plateau | 15 |
| Permian of the Hueco and Franklin Mountains | 16 |
| Permian of the Finlay Mountains | 17 |
| Permian of the Carrizo and Wylie Mountains | 17 |
| Permian of the Shafter district | 17 |
| Permian near Las Delicias, Coahuila, Mexico | 18 |
| Correlation of the Permian sections of Trans-Pecos Texas with other American Permian sections | 18 |
| Permian of the mid-continent region | 18 |
| Permian of Arizona and New Mexico | 24 |
| Permian of the Great Basin and northern Rockies | 30 |
| Permian of northern California | 33 |
| Permian of Arctic America | 33 |
| "Upper Carboniferous" of the Andes | 35 |
| Paleontology | 36 |
| Summary of the Permian faunas | 36 |
| Descriptions of brachiopod genera and species | 42 |
| *Orbiculoidea* | 42 |
| *Crania* | 42 |
| *Rhipidomella* | 43 |
| *Orthotichia* | 44 |
| *Enteletes* | 45 |
| *Parenteletes*, nov. | 48 |
| *Streptorhynchus* | 49 |
| *Orthotetella*, nov. | 51 |
| *Orthothetina* | 52 |
| *Meekella* | 52 |
| *Geyerella* | 57 |
| *Derbya* | 58 |
| *Chonetes* | 60 |
| *Isogramma* | 64 |
| *Productus* | 65 |
| *Linoproductus* | 75 |
| *Striatifera* | 78 |
| *Pustula* | 79 |
| *Buxtonia* | 79 |
| *Waagenoconcha* | 80 |
| *Overtonia* | 82 |
| *Avonia* | 82 |
| *Horridonia* | 85 |
| *Marginifera* | 86 |
| *Scacchinella* | 91 |
| *Aulosteges* | 92 |
| *Strophalosia* | 96 |
| *Teguliferina* | 96 |
| *Prorichthofenia*, nov. | 97 |
| *Parakeyserlingina* | 100 |
| *Lyttonia* | 102 |

PAGE
Leiorhynchus ..... 104
Pugnoides ..... 106
Rhynchopora ..... 109
Camerophoria ..... 110
Uncinuloides, nov. ..... 112
Spirifer ..... 113
Squamularia ..... 118
Ambocoelia ..... 119
Martinia ..... 120
Spiriferina ..... 121
Punctospirifer ..... 124
Hustedia ..... 125
Composita ..... 128
Dielasma ..... 131
Dielasmina ..... 133
Index to localities ..... 133
Localities in The University of Texas collections ..... 139
Localities in Delo collection ..... 140
Localities in Schuchert collection ..... 140
Localities in R. F. Baker collection ..... 140
Bibliography ..... 140
Index ..... 241

# ILLUSTRATIONS

## FIGURES

Fig. 1. Correlation chart ..... Following 146
Fig. 2. Table of fossils ..... 147
Fig. 3. Table of fossils ..... 148
Fig. 4. Table of fossils ..... 149
Fig. 5. Table of fossils ..... 150

## PLATES

Plates I–XLIV. Brachiopoda of the Glass Mountains ..... Following 150

# THE GEOLOGY OF THE GLASS MOUNTAINS, TEXAS

## PART II

### Faunal Summary and Correlation of the Permian Formations with Description of Brachiopoda

ROBERT EVANS KING


## ACKNOWLEDGMENTS

This memoir is the result of field work in Trans-Pecos Texas by the writer and his brother, Philip B. King, during the summers of 1925 to 1928, and of the laboratory study of their own and various other collections of Permian fossils from this region. The work was presented as a doctor's dissertation at Yale University in 1929, and final work on the manuscript by the author was completed in August of that year.

The investigation was largely made possible by the financial support of the writer's parents, though a material aid to field work was given through generous purchase of our fossil collections by Prof. Charles Schuchert, and by allotments for field expenses by the Texas Bureau of Economic Geology. I am also indebted to Professor Schuchert for his contagious enthusiasm for problems of the Permian, and the inspiration and assistance which he has given in that branch of the major problem on which I worked. Prof. C. O. Dunbar, of Yale University, has given invaluable criticism and counsel as the work progressed. After its completion, when I was called abroad in commercial work, he freely gave his time to editorial work on the manuscript, and by his intimate acquaintance with the fauna herein described was able to keep the work up to date by following more recent studies of paleontologists on this and related topics.

In addition to the collections made especially for this study, which are now the property of Peabody Museum, the collections which Udden, Böse, Beede, and Baker made for the Texas Bureau during the earlier investigations on the Permian were available to me for study through the courtesy of Dr. E. H. Sellards, the associate director of the Bureau. A small collection of Leonard and Word fossils, which furnished valuable supplementary material, was generously submitted for identification by Mr. D. M. Delo, now of Northwestern University, while he was with the Southwestern Crude Oil Purchasing Company in Texas. A small collection from Trans-Pecos Texas made by Mr. Ray Baker of the Texas Company was also examined. Finally, the collections made by Professor Schuchert in western Texas and Arizona were freely placed at my disposal.

The collateral study of Permian faunas outside Trans-Pecos Texas was based very largely on an extensive review of the American and foreign literature. The writer also had access to the reference sets of fossil collections at Peabody Museum, which included not only many suites from important localities and formations in the west, but also from other parts of the continent and abroad.


## HISTORICAL RESUMÉ

The first fossil described from the Permian of Trans-Pecos Texas was *Composita mexicana,* which was collected by Emory's United States and Mexican Boundary Survey and described by Hall in 1857. One species from the Permian of the Delaware Mountains was described by Marcou in 1858. The first important contribution was made by the Shumards. As geologist of the expedition under Captain Pope, sent to discover artesian waters in the arid southwest, George C. Shumard collected some fossils from the Delaware, Guadalupe, and Hueco mountains, which he sent to his brother, B. F. Shumard, for study. Shumard described these collections in 1858 and 1859, and recognized the fossils from the Delaware and Guadalupe mountains as being of Permian age. The fauna of the Hueco Mountains was assigned to the Pennsylvanian. During the remainder of the nineteenth century no further paleontologic work was done in the region, and the fauna described by Shumard came generally to be regarded as Upper Carboniferous rather than Permian in age. In 1901 G. H. Girty accompanied B. F. Hill to the Delaware and Guadalupe mountains, and spent about two weeks collecting from the Permian rocks. Collections made by G. B. Richardson and R. T. Hill in other ranges in Trans-Pecos Texas were studied along

Printed August, 1931.

with Girty's own collections, and in 1908 the classic "Guadalupian Fauna" was published as United States Geological Survey Professional Paper 58. In this work, Shumard was shown to have been correct in regarding his faunas as of Permian age. The richness of the Permian faunas of this region was first made known as a result of this work. Unfortunately, circumstances prevented Dr. Girty from devoting enough time to collecting to obtain very complete material; many of his species are therefore established on only a few specimens and prove rather difficult to use. Incomplete field work also prevented any attempt at definite zonation of the faunas, and they were treated very much as a unit, to which the name "Guadalupian" was applied. The first zonation of the Trans-Pecos Permian section was made possible by the work of the late Emil Böse. Böse studied the collections he gathered and those made by Udden and his associates in the first work on the stratigraphy of the Glass Mountains, and in 1917 published detailed descriptions of the ammonoids from this area and discussed in detail their stratigraphic significance. As a result, four main ammonoid zones were made known, and were correlated with the ammonoid bearing formations in other parts of the world. In 1927 Schuchert reviewed the Glass Mountains sequence and published lists of some of the fossils. In 1929 J. P. Smith restudied the ammonoid fauna of the basal formation of the Permian, the Wolfcamp, bringing forth new facts in proof of its age.

## STRATIGRAPHY AND CORRELATION

### SUMMARY OF THE PERMIAN STRATIGRAPHY OF TRANS-PECOS TEXAS

#### THE GLASS MOUNTAINS SECTION

By accidents of structure and erosion, the most complete Permian section in Trans-Pecos Texas is in the Glass Mountains, north of the town of Marathon, where about 7000 feet of strata of this age dip northwestward away from the older rocks of the Marathon uplift. Most of the collections described in the present paper were made in this district.

The basal Permian beds rest with a considerable structural unconformity on the underlying rocks. Evidence indicates, however, that the youngest beds in the subjacent series, the upper part of the Gaptank formation, are not greatly older than the basal strata above the unconformity. West of Marathon they contain at one place the *Proudden-ites* fauna, in which A. K. Miller* has identified nine genera and 10 species of ammonoids; F. B. Plummer regards this as comparable to one of his Canyon ammonoid zones in Central Texas. Both to the east and west of Marathon the occurrence of such characteristic fusulinids as *Triticites secalicus* and *T. cullomensis* indicates the existence of strata as young as the Cisco group as well.*

*The Wolfcamp formation.*—The lowest Permian formation of the Glass Mountains section is the Wolfcamp, which varies from 700 feet in thickness at the type locality to only 50 feet toward the east of that locality and several hundred feet toward the west. At the type locality no angular break is apparent between it and the underlying Gaptank formation, of Pennsylvanian age, but to the west where the Gaptank beds are strongly folded there is a marked angular discordance. The section at Wolfcamp starts with the *Uddenites* zone, a shale with limestone layers, varying in thickness up to 100 feet. This unit contains a great number of ammonoids, which have been shown by Smith (**156**, pp. 63–80) to prove its lowest Permian age. The brachiopod fauna of the *Uddenites* zone comprises mostly species which occur also in the Pennsylvanian, but has in addition some notable innovations. The following is a numerical analysis of the fauna:

Number of species—34
Number also in middle and upper Wolfcamp—28
Number also in Hess—5
Number also in Leonard—3
Number also in Word—1
Number also in Pennsylvanian—22
Present in Bolivia—16
Present in Bolivia, not in Pennsylvanian—6
Present in higher Permian, not Pennsylvanian—12
Present in Pennsylvanian and higher Permian—17
Present in Pennsylvanian, not higher Permian—5
Restricted to this horizon—0

Among the lower Wolfcamp species occurring in higher beds of the Texas Permian section but not occurring in the Pennsylvanian are *Meekella irregularis texanus, Waagenoconcha montpelierensis, Avonia boulei, Scacchinella gigantea, Aulosteges wolfcampensis, Spirifer condor,* and *Martinia wolfcampensis.* All these species are definitely Permian types. An interesting element in the fauna is

*Miller, A. K., A New Ammonoid Fauna of Late Paleozoic Age from Western Texas, Jour. Pal., Vol. 4, pp. 383–412, 1930.

*For more detailed description of the Gaptank fauna see Part One of this memoir.

*Paraënteletes cooperi,* which occurs elsewhere in America only in the underlying Gaptank formation. Similarities may be recognized with two foreign faunas, that of the Andean "Upper Carboniferous" and that of the Trogkofel limestone of the Karnic Alps. Ten of the species in common with the Andean fauna occur also in the Pennsylvanian, but *Waagenoconcha montpelierensis, Avonia boulei, Marginifera capaci, Orthotichia kozlowskii, Spirifer condor,* and possibly *Linoproductus villiersi* do not occur below the *Uddenites* zone. This indicates that the brachiopod assemblage of the *Uddenites* zone is more closely allied to the Andean one than are those of the Pennsylvanian, which contain only 3 species in common with the Andean fauna and not occurring in the Permian of this country. The species *Meekella irregularis texanus, Scacchinella gigantea, Paraënteletes cooperi,* and *Teguliferina bösei* are very close to species in the Trogkofel limestone. In summary, it may be noted that the brachiopod fauna of the *Uddenites* zone has many identities with the Pennsylvanian, yet contains other species that are definite heralders of the Permian, and some of which are unlike American Pennsylvanian forms, but have foreign affinities.

The middle and upper parts of the Wolfcamp formation.—At the type locality, the part of the Wolfcamp formation above the *Uddenites* zone consists largely of shale, containing occasional limestone layers. Immediately above this zone *Schwagerina* makes its first appearance. This distinctive fusulinid genus is abundant there and in the succeeding beds of the Wolfcamp. Near the middle of the formation is the most prolific fossil locality (loc. 93). In the western part of the Glass Mountains the upper part of the Wolfcamp alone is fossiliferous, the lower part there consisting of conglomerate. On Dugout Mountain the upper Wolfcamp contains a rather large and well preserved fauna, including many ammonoids, among which are *Perrinites cumminsi* and *Prothalassoceras welleri.* Fossiliferous Wolfcamp is present also in the uplift northeast of the Hess Ranch, where there are a good many *Schwagerina,* associated with molluscs and echinoid spines and plates. The following is a numerical analysis of the brachiopods in the middle and upper parts of the Wolfcamp formation:

Number of species—46
Number also in Pennsylvanian—18
Number also in *Uddenites* zone—28.
Number also in Hess—12
Number also in Leonard—10
Number also in Word—5
Number also in Capitan—3
Present in Bolivia—13
Present in Bolivia, not in Pennsylvanian—4
Present in higher Permian, not lower—11
Present in lower and higher Permian—4
Present in lower and not higher Permian—24
Restricted to this horizon—6

Many interesting brachiopod types make their first appearance in this fauna, heralding the developments of succeeding faunas. Ancestors of *Prorichthofenia* and *Lyttonia* are present, and the productids show a great expansion. The following brachiopods are particularly characteristic of the fauna of the middle and upper Wolfcamp:

Orthotetella wolfcampensis
Derbya buchi
Productus gratiosus occidentalis
P. semistriatus
P. wolfcampensis
Avonia boulei
Marginifera capaci
Aulosteges wolfcampensis
Teguliferina bösei
Parakeyserlingina fredericksi
Camarophoria thevenini
Spirifer condor
Martinia wolfcampensis

*The Hess formation.*—Unconformably above the Wolfcamp is the Hess formation, which varies in thickness from 2100 feet in the east to 50 feet in the west. It consists of limestone and dolomite, with a basal conglomerate. In the east the dolomite predominates, while in the west there is mostly limestone. Fossils are abundant only at certain horizons, mainly in the upper part of the formation. One fossiliferous layer on the escarpment above Wolf Camp has yielded a large proportion of the fossils which have been collected from the formation. Localities 107, 108, 212, 215, and 222 are at this horizon. Fossils are more common in the limestone facies of the west, but as the limestone is very hard, they are difficult to collect.

The great decrease in thickness from east to west in the Glass Mountains is especially remarkable because of the corresponding thinning of the younger and conformable Leonard in the reverse direction. This fact might be interpreted as due to the intergradation of lower Leonard and upper Hess. However, the evidence that could be gathered in the field and in a study of the faunas indicates that the two formations are distinct time units. The upper Hess contains the ammonoid *Perrinites compressus,* while in the lower Leonard the more specialized species *Perrinites vidriensis*

is common; the two have not been found in association. *Enteletes dumblei,* a common upper Hess fossil, does not occur in the Leonard of the Glass Mountains. *Scacchinella gigantea* is abundant in the Hess at certain localities in both the western and eastern facies of the formation, in one place but a few feet below the top of the formation, yet it has not been collected from the Leonard. The primitive species of *Prorichthofenia, P. tegulifer-oides,* occurs in the upper Hess of both the Glass Mountains and the Sierra Diablo, but has not been found in the Leonard. While it is true that a large proportion of the characteristic Leonard brachiopods make their first appearance in the upper Hess, there are 13 species in the Hess that do not occur at a higher horizon, and several of these are present in the Hess of the Sierra Diablo and the Gym of the Hueco Mountains—a fact indicating that the Hess fauna is a separate biota. It might be argued that the reason for the differences between the Hess and Leonard faunas is that the two are of distinct faunal facies. This seems to be disproved by two facts. In the first place, different faunal facies are ordinarily to be correlated with differences in lithology, yet the lower Leonard contains limestone members which are lithologically similar to the underlying upper part of the Hess but contain Leonard types of fossils, including *Perrinites vidriensis.* In the second place, many of the species in the Hess do occur in the limestones of the Leonard, so that if there is any difference in faunal facies it is not great enough to prevent a large number of identical forms from occurring in both. A final argument is the evidence for a distinct faunal zone in the highest beds of the Hess and the lowest beds of the Leonard, characterized by the species *Enteletes plummeri.* This faunule occurs at the top of Leonard Mountain, and occupies the same stratigraphic position 5 miles to the southwest of there, though the Hess limestone underlying it has thinned from 750 feet to 250 feet.

As noted above, the ammonoid *Perrinites compressus* occurs at the upper Hess fossiliferous horizon. This form is intermediate in complexity between *P. cumminsi* of the Wolfcamp and *P. vidriensis* of the Leonard. Böse believed that the presence of the genus *Perrinites* at this horizon proved that the fauna was of Leonard age, but it has been shown that *Perrinites* is not restricted to the Leonard, making its first appearance in the Wolfcamp. The ammonoid *Prothalassoceras welleri* was collected by Baker and Böse from the conglomerate at the base of the Permian in the western Glass Mountains, at a horizon regarded by Böse as Hess. Hence the Hess was called the "zone of *Prothalassoceras*" in the "Ammonoids of the Glass Mountains." We have shown that the conglomerate from which this fossil was collected is actually part of the Wolfcamp formation, and furthermore we have collected *Prothalassoceras welleri* from the upper Wolfcamp on Dugout Mountain. (See section on Wolfcamp formation above. The genus *Prothalassoceras* has also been identified, by Miller, in the Gaptank formation.)

The following table indicates numerically the relations of the Hess brachiopod fauna:

Number of species—53
Number also in Pennsylvanian—2
Number also in *Uddenites* zone—4
Number also in Wolfcamp—9
Number also in Leonard—38
Number also in Word—20
Number also in Capitan—10
Present in Bolivia—1
Present in Bolivia, not in Pennsylvanian—1
Present in higher Permian, not lower—32
Present in lower and higher Permian—4
Present in lower and not higher Permian—8
Restricted to this horizon—10

One of the most notable features of the Hess fauna of the limestone facies is the abundance of brachiopods having high cardinal areas—especially *Scacchinella gigantea, Geyerella americana,* and *Streptorhynchus undulatus.* The former occurs in great abundance in some places. These same genera are present also in the Trogkofel formation of the Karnic Alps, and the first two in the Sosio limestone of Sicily. It is probable, however, that their presence is not of stratigraphic significance, but proves the presence of a facies, perhaps of reef nature, like the rudistid reef limestones of the Mesozoic. It is very likely that the Capitan is of the same nature, for it is, similarly, a dense hard limestone containing brachiopods having high, deformed cardinal areas. Notable innovations occur in the Hess, since here is the first appearance of *Prorichthofenia,* in a primitive species transitional from *Teguliferina.* The first specimens of *Lyttonia* occur here. Many of the species which are common in the Leonard and Word formations first appear in the Hess. It may therefore be said that the Wolfcamp fauna is transitional from the Pennsylvanian to the Permian, while that of the Hess is

characteristically Permian. The following are the more important brachiopods which occur in the Hess:

Rhipidomella hessensis
Enteletes dumblei
Streptorhynchus lamellatus
S.? undulatus
Meekella attenuata
Geyerella americana
Chonetes hessensis
Productus ivesi
Striatifera pinniformis
Marginifera whitei
Scacchinella gigantea
Aulosteges medlicottianus
Prorichthofenia likharewi
Prorichthofenia tegulifer-oides
Lyttonia nobilis americanus
Pugnoides elegans
P. texanus
P. transversus
Camarophoria venusta
Spirifer huecoensis
Squamularia guadalupensis
Spiriferina angulata
Hustedia hessensis
Composita mexicana
C. subtilita

At locality 223, 1.1 mile southwest of the Word Ranch, there is a very peculiar faunule in the highest beds of the Hess. *Composita mexicana* and *Pugnoides texanus* occur in great abundance, almost to the exclusion of other species of brachiopods. Several genera of ammonoids, including *Perrinites*, also are present. The abundance of the two species of brachiopods reminds one of the upper Gym, where the same forms are very common.

*The Leonard formation.*—This unit consists of limestone, shale, siliceous shale (radiolarite) and conglomeratic limestone, varying from 1800 feet in thickness in the western Glass Mountains to 300 feet in the east. Fossils are abundant in all the limestone and some of the shale layers. Some of the best localities are in the middle part of the formation north and northwest of Iron Mountain—localities 120 and 123. Ammonoids are abundant in places; *Perrinites vidriensis* is the most common species. The best ammonoid locality is in the middle part of the upper Leonard below the Clay Slide, northwest of Leonard Mountain.

The following table summarizes the brachiopods of the Leonard fauna:

Number of species—69
Number also in Pennsylvanian—1
Number also in *Uddenites* zone—1
Number also in middle and upper Wolfcamp—8
Number also in Hess—35
Number also in Word—39
Number also in Capitan—16
Present in higher Permian, not lower—21
Present in lower and higher Permian—21
Present in lower and not higher Permian—18
Restricted to this horizon—9

*Prorichthofenia* and *Lyttonia* occur abundantly in the Leonard, and there is a great diversity of productids. One of the most important species in the Leonard fauna is *Productus ivesi,* which is abundant in many places, but especially so 4.3 miles northeast of the Word Ranch, at locality 174, where the rock is a sandy limestone. *Productus leonardensis* is another of the most abundant and typical species. Affinity to the Kaibab and San Andres faunas of Arizona and New Mexico is seen in the abundance of *Productus ivesi,* as well as of two other forms which are common in the Leonard, *Marginifera manzanica* and *M. cristobalensis.*

The following are the more important species of brachiopods occurring in the Leonard:

Enteletes plummeri
Meekella difficilis
M. globosa
Chonetes subliratus
C. permianus
Productus ivesi
P. leonardensis
P. occidentalis
P. schucherti
Linoproductus waagenianus
Marginifera cristobalensis
M. manzanica
M. reticulata
M. reticulata angusta
M. sublaevis
Aulosteges magnicostatus
A. medlicottianus
A. subcostatus
Aulosteges trigonalis
Prorichthofenia likharewi
P. uddeni
Lyttonia nobilis americanus
Pugnoides bidentata
Camarophoria venusta
Uncinuloides guadalupensis
Spirifer marcoui infraplica
S. costella
S. mexicanus latus
S. pseudocameratus
Squamularia guadalupensis
Martinia rhomboidalis
Spiriferina hilli
Hustedia meekana
H. mormoni papillata
Composita mira
C. subtilita

*The Word formation.*—This division consists in the western Glass Mountains of three main limestone members separated by shale, sandstone, and siliceous shale (radiolarite), about 1500 feet thick. In the eastern Glass Mountains the formation consists only of about 300 feet of dolomite. Fossils are abundant in the limestones, occurring generally silicified. The following is a statistical summary of the brachiopods of the entire Word formation:

Number of species—77
Number also in Pennsylvanian—0
Number also in Wolfcamp—6
Number also in Hess—22
Number also in Leonard—40
Number also in Capitan—25
Present in higher Permian, not lower—11
Present in lower and higher Permian—14
Present in lower and not higher Permian—27
Restricted to this horizon—25

Each member of the Word formation has its characteristic fauna. The lower Word limestone, which is best developed in the western part of the

Glass Mountains, contains an abundance of silicified fossils. The best collecting localities are in the southern foothills of Cathedral Mountain. Ammonoids are rare, however, at this horizon. The more important species of brachiopods are:

| | |
|---|---|
| Chonetes subliratus | Camarophoria venusta |
| Productus indicus | Hustedia meekana |
| Linoproductus waagenianus | Composita mira |
| Avonia subhorrida rugatula | |

The middle limestone member, which occurs in its best development on the south side of Road and Hess canyons, and capping the mountain north of Leonard Mountain, is the most fossiliferous of the three divisions. There is a great abundance in places of ammonoids, of which the most important are *Waagenoceras dieneri, Medlicottia burckhardti, Gastrioceras roadense,* and *Stacheoceras bowmani. Prorichthofenia* and *Lyttonia* are especially abundant, as well as spinose productids. The following species of brachiopods are especially characteristic:

| | |
|---|---|
| Enteletes dumblei | Pugnoides swallowiana |
| Meekella attenuata | Rhynchopora taylori |
| Meekella skenoides | Camarophoria venusta |
| Productus multistriatus | Spirifer pseudocameratus |
| Avonia signata | S. sulcifer |
| A. walcottiana | Squamularia guadalupensis |
| Marginifera opimus | Spiriferina laxa |
| M. popei | Punctospirifer billingsii |
| M. wordensis | Hustedia meekena |
| Aulosteges guadalupensis | Composita emarginata affinis |
| A. tuberculatus | Composita mira |
| Prorichthofenia permiana | Dielasma spatulatum |
| Lyttonia nobilis americanus | Dielasmina schucherti minor |

The upper limestone member, which does not occur in its normal development to the west of Gilliland Canyon, is very profusely fossiliferous near Hess Canyon. Some species of productids are especially abundant. The fauna of this member has many identities with species of the upper dark limestone, which occupies a similar position at the top of the Delaware Mountain formation. Many of the species have Arctic affinities, and a large number occur also in the Phosphoria. The following brachiopods are especially characteristic:

| | |
|---|---|
| Meekella attenuata | Avonia signata |
| Derbya crenulata | Marginifera opimus |
| Chonetes quadratus | Marginifera popei |
| Productus arcticus | M. texana |
| P. guadalupensis | Spirifer sulcifer |
| Linoproductus nasutus | Spiriferina laxa |
| L. phosphaticus | Punctospirifer billingsii |
| Waagenoconcha montpelierensis | Hustedia meekana |
| | Composita emarginata affinis |

Permian brachiopod evolution culminates in the Word faunas, for the species of the Capitan show little further development. The Word is the highest formation in the Trans-Pecos Permian section which has a truly cosmopolitan fauna, and it is probable that thereafter the seas in this region became so restricted that there was no longer free interchanging of forms with the seas of other continents.

*The Capitan formation.*—This is the highest marine unit of the Glass Mountains Permian section. In the region of Hess Canyon, near the center of the Glass Mountains, it is about 3000 feet thick, but it thins greatly to the west and probably to a lesser extent to the east. It is divided into several members. In the Hess Canyon region the lower member is the Vidrio dolomite, the middle the Gilliam sandstone, red shale, dolomite, and gypsum, and the upper the Tessey dolomite. In the western part of the Glass Mountains the Altuda member of limestone and sandstone wedges into the dolomite comprising the main part of the formation. Except for *Fusulina elongata,* which is abundant throughout the Capitan, fossils are rare. No ammonoids have been found in the Capitan either here or in the Guadalupe Mountains. Some brachiopods were collected from the Altuda member, and some from the Gilliam. Where they occur, individuals are extremely abundant, but the number of species is small, the most common types being *Squamularia guadalupensis, Martinia subquadrata,* and *Composita emarginata affinis.* At one place, a great number of specimens of *Pleurophorus* were found in the Tessey. A total of 11 species of brachiopods was found in the Capitan in the Glass Mountains, all of which occur in lower formations. Two species occur also in the Wolfcamp, 4 in the Hess, 6 in the Leonard, and 10 in the Word. No species were found which occur also in the Pennsylvanian. A large part of the Capitan limestone, particularly in the more massive portions, is composed of the remains of lime-secreting algae. It is not improbable that these plants built up large reefs during Capitan time. This suggestion is more fully treated in Part One of this memoir.

*The Bissett formation.*—At the top of the Glass Mountains Permian section is the Bissett conglomerate and red shale, which attains a thickness of 700 feet. It contains a few layers of fossiliferous dolomite, containing ostracods, pelecypods, and

gastropods, all poorly preserved, as well as a flora regarded by David White as of Upper Permian age. The conglomerate contains cobbles from the underlying Permian. This formation apparently was deposited during the late Permian period of uplift, as the final phase of Permian sedimentation.

### THE DELAWARE AND GUADALUPE MOUNTAINS SECTION

The Delaware and Guadalupe mountains together form a more or less connected chain extending across northern Trans-Pecos Texas into New Mexico. They are composed entirely of Permian rocks of a similar facies to that in the Glass Mountains, with their strata tilted toward the east as a result of block faulting. The lowest beds are brought to view along their western base, but these rocks are no older than those near the middle of the section in the Glass Mountains. To the strata outcropping on the face of the mountains the name Guadalupe group has been applied, but the extended use of this title has proved unfortunate, since its indefinite lower limit depends on the accidents of structure, rather than on a stratigraphic or paleontologic definition.

*The Bone Canyon member.*—In general, the rocks of the Delaware and Guadalupe mountains resemble those of the Glass Mountains, but there are differences in detail. This is particularly true of the Leonard formation at the base of the section, which is here divisible into two members of limestone, of which the lower, or Bone Canyon member, consists of as much as 1000 feet of black limestone (the "basal black limestone" of Girty). It is exposed along the lower part of the Delaware and Guadalupe mountains escarpment as far south as Seven Heart Gap. Fossils are abundant in certain layers. From this horizon Girty described the ammonoids *Peritrochia erebus, Paraceltites elegans,* and *Agathiceras texanum. Perrinites* has also been identified in this horizon by other geologists working in the area. The following is a numerical analysis of the brachiopods in the Bone Canyon member:

Number of species—15
Number also in Pennsylvanian—0
Number also in Wolfcamp—1
Number also in Hess—10
Number also in Leonard—11
Number also in Word—10
Number also in Capitan—5
Present in Bone Canyon member, Sierra Diablo—5
Restricted to this member—0
Present in Leonard, not Hess—5
Present in Hess, not Leonard—2
Present in Word, not Leonard—1
Present in Leonard, not Word—4

Among the species which point clearly to a Leonard age for this unit are *Productus leonardensis, Marginifera cristobalensis, Pugnoides texanus, P. bidentatus,* and *Composita mexicana. Pugnoides texanus* and *Composita mexicana* are common also in the Hess, but the other forms are practically diagnostic of the Leonard.

*The Victorio Peak member.*—In the Delaware Mountains and at Guadalupe Point the Bone Canyon member is overlain without apparent break by the Delaware Mountain sandstone. On the west side of the Guadalupe Mountains, however, at Bone Canyon, there is a thick conglomerate at the base of the Delaware Mountain formation, most of its boulders consisting of white limestone. A few miles to the north the white limestone from which these boulders were derived appears beneath the Delaware Mountain sandstone and above the Bone Canyon black limestone. This limestone is known as the Victorio Peak member of the Leonard formation, the name being taken from Victorio Peak in the Sierra Diablo, to the southwest. In the Guadalupe Mountains this member is several hundred feet thick.[1] The fossils determined from the Victorio Peak member by Girty[2] do not indicate conclusively its age. *Enteletes* sp. c. Girty is *E. liumbonus* King, a species occurring in the Hess formation. It is possible that the species listed by Girty here is not conspecific with the form originally described and figured as *E.* sp. c., but *E. leonardensis,* which is somewhat similar. The original of *Enteletes* sp. d is supposed by me to be *E. wordensis,* but in any case such an identification is uncertain. *Meekella attenuata* is a long ranging species. *Productus guadalupensis* var. *comancheanus* occurs in the Word in the Glass Mountains. *Pustula subhorrida* var. *rugatula?* may be *Marginifera cristobalensis. Aulosteges guadalupensis* is restricted to the Word in the Glass Mountains, and *A. magnicostatus* is most common in the Leonard of the same area. The other forms listed are of very little importance or are uncertainly identified. Because of the very incomplete nature of the collections on which the descriptions in "The Guadalupian Fauna" are based, it is difficult to use that book in

[1]Darton, N. H., and Reeside, J. B.: Guadalupe Group, Bull. Geol. Soc. Amer., vol. 37, pp. 421–423, fig. 4, 1926.
[2]Idem, p. 421.

making some of the identifications, and it is probable that some of the species above mentioned have been misidentified. Probably the most reliable of the species listed is *Aulosteges magnicostatus,* for it is easily identified, and could hardly be confused with anything else. Its presence suggests a Leonard age for this horizon. However, there is little besides the lithologic similarity of this member with that on the other side of the Salt Basin (in the Sierra Diablo) to prove its age as Leonard.

*The Delaware Mountain formation* (in restricted sense).—This unit is about 2100 feet thick at Guadalupe Point. The lower 2000 feet consist largely of sandstone, while the upper 100 feet is the "upper dark limestone" member of Girty. To the north, along the west side of the Guadalupe Mountains, the Delaware Mountain formation thins gradually against the unconformity which is seen in Bone Canyon and northward thereof, so that it is little more than 100 feet thick at the Texas-New Mexico state line. South of Guadalupe Point it comprises the upper, main part of the escarpment of the Delaware Mountains as far south as Seven Heart Gap.

The fauna of the Delaware Mountain sandstone has been made known by Girty in "The Guadalupian Fauna." The ammonoid *Waagenoceras* is present, as well as *Paraceltites elegans, Gastrioceras? serratum,* and *Gastrioceras* sp. The brachiopod fauna gives still further proof of the Word age of this formation, as shown by the following statistics:

Number of species—52
Number also in Wolfcamp—0
Number also in Hess—8
Number also in Leonard—15
Number also in Word—33
Number also in Capitan—29
Restricted to this formation—7
Present in Word, not Leonard—28
Present in Leonard, not Word—3
Present in Word, not Capitan—20
Present in Capitan, not Word—10

The upper dark limestone member of the Delaware Mountain formation is a widespread unit which underlies the Capitan limestone at Guadalupe Point and the Castile gypsum farther to the east and south. In the Glass Mountains two species from this horizon, *Derbya elevata* and *Chonetes hillanus,* occur only in the upper limestone member of the Word formation and in the Capitan. *Marginifera popei, M. opimus, Spirifer sulcifer,* and *Spiriferina laxa,* which occur in the dark limestone of the Delaware Mountain, occur in both the middle and upper Word limestones but are especially characteristic of the upper Word limestone. Other species which are common in the upper Word, such as *Composita emarginata affinis,* show definite affinity to Capitan species. Another point of resemblance between the dark limestone and the upper Word limestone is that *Lyttonia* was not reported by Girty from the former, and it is very rare in the latter, though common in the middle Word limestone and the Delaware Mountain sandstone. The close similarity in faunas of the upper dark limestone of the Delaware Mountain formation with the upper limestone of the Word, along with their similar stratigraphic position, makes their equivalence fairly certain.

Until recently the upper dark limestone member at Guadalupe Point was thought to extend northeastward along the base of the Guadalupe Mountains to McKittrick Canyon, and from there to trace southward along the east slope of the Delaware Mountains between the main mass of the Delaware Mountain formation and the Castile gypsum. This eastward extension was named the Frijole limestone. In 1929, however, Lloyd* expressed the opinion that the upper dark limestone at McKittrick Canyon interfingers with steeply inclined foreset beds dipping off the upper part of the Capitan reef-mass, and that though the member was only 50 feet thick it was "the time equivalent of the upper part of the (Capitan) reef, which is 1000 feet or more thick." These seemingly inexplicable relationships are clarified by Blanchard and Davis* who have traced the Frijole limestone "to Guadalupe Canyon, of which Guadalupe Point forms the west side, and at the place where it lost its identity through transition into the massive dolomite it was above approximately 1000 feet of the dolomite of Capitan facies." The dark limestones from which Girty collected his fauna are therefore considerably lower than the Frijole, and must correlate with some horizon well down in the Delaware Mountain formation in the area to the south.

*Lloyd, E. Russel, Capitan Limestone and Associated Formations, Bull. Amer. Assoc. Petrol. Geol., Vol. 13, p. 650, 1929.

*Blanchard, W. G., and Davis, M. J., Permian Stratigraphy and Structure of Southeastern New Mexico and Southwestern Texas, Bull. Amer. Assoc. Petrol. Geol., Vol. 13, pp. 979-980, 1929.

By this interpretation the Delaware Mountain formation at its type section in the Delaware Mountains includes two stratigraphic components of identical facies, but different relationships. Below the level of the true "upper dark limestone" are strata which correlate with the Delaware Mountain sandstones at Guadalupe Point and to the north of it, as well as with the Word formation of the Glass Mountains. Between the level of the dark limestone and the Frijole limestone are beds equivalent in age to the Capitan limestone of the Guadalupe and Glass mountains, and of open sea rather than reef facies, and thus exactly comparable to the Altuda member of the Glass Mountains Capitan.

*The Capitan limestone.*—At the top of the Guadalupe Mountains section is the Capitan limestone, about 2000 feet thick, which forms the upper part of the mountains, terminating to the south at the bold bluff of Guadalupe Point. From it Girty has described a large fauna. A large proportion of the Capitan forms occur in the Word-Delaware Mountain, and the fauna as a whole can hardly be regarded as more than a development from the latter fauna. There is no evidence of migrants from other Permian seaways, for at this time the Permian sea in Trans-Pecos Texas probably had the form of an inclosed embayment from the south, so restricted in area that free intermigration of species was not possible. Ammonoids are not present in the formation. The following brachiopods are especially characteristic:

| | |
|---|---|
| Streptorhynchus gregarium | Prorichthofenia permiana |
| Geyerella americana | Pugnoides elegans |
| Chonetes hillanus | Camarophoria? indentata |
| Productus capitanensis | C.? longaeve |
| P.? pileolus | Uncinuloides guadalupensis |
| P.? limbatus | Spirifer mexicanus |
| Avonia latidorsata | S. sulcifer |
| Linoproductus waagenianus | Squamularia guadalupensis |
| Marginifera popei | Martinia rhomboidalis |
| Striatifera pinniformis | Spiriferina laxa |
| Leiorhynchus bisculatus | Punctospirifer billingsii |
| Lyttonia nobilis americanus | Composita emarginata |

Recently, various writers* have interpreted the Capitan limestone as a great reef mass built by corals and algae, a conclusion in harmony with that of the writers of this memoir for the Capitan in the Glass Mountains. As noted above, these writers show that the portion of the Delaware Mountain formation of the type area which lies above the horizon of Girty's upper dark limestone grades laterally into the Capitan reef dolomite, and is presumably its open sea facies like the Altuda member of the Glass Mountains. The Capitan reef, in its typical massive facies, extends northeast along the Guadalupe Mountains escarpment from Guadalupe Point toward Carlsbad. A few miles northwest of the reef front it merges into a complex of thinner-bedded deposits, presumably of lagoonal origin, divided into three main members. The lower, a bedded limestone, has been called the San Andres by Blanchard and Davis and the Chupadera by Crandall. Neither name is very appropriate since both of them imply equivalence to New Mexico formations whose relationship is decidedly dubious. This is succeeded by the sandstones, red beds, and gypsums of the Queen and Seven Rivers members, which thicken northward into New Mexico. The highest member of the lagoonal succession, known as the Carlsbad member, is again a well bedded limestone.

Southeast of the reef escarpment, the Frijole limestone is overlain by the Castile gypsum and Rustler limestone. At one time these were believed to be laterally equivalent to the Capitan, a relationship thought to result from rapid shoreward transition from limestone and dolomite into gypsum and red beds. The revised interpretation of the Frijole now makes it clear that these two formations are younger than the Capitan, as was originally announced by Darton and Reeside* and later by Crandall** and others. Unfortunately, the fauna of the Rustler limestone is too poor in variety and preservation to furnish much information on the life of these highest Permian deposits.

In the Apache Mountains, which lie south of Seven Heart Gap at the south end of the Delaware Mountains, massive dolomites† appear again and are considered by many geologists (as Crandall, p. 939) to be another segment of the Capitan reef.

*Ruedemann, R., quoted by King and King, **90**, p. 139, 1928.
Lloyd, *op. cit.*, pp. 645–658, 1929.
Crandall, K. H., Permian Stratigraphy of Southeastern New Mexico and Adjacent Parts of Western Texas, Bull. Amer. Assoc. Petrol. Geol., Vol. 13, pp. 927–944, 1929.
Blanchard and Davis, *op. cit.*, pp. 957, 996, 1929.

*Darton, N. H., and Reeside, J. B., Guadalupe Group, Bull. Geol. Soc. Amer., Vol. 37, p. 420, 1926.
**Crandall, *op. cit.*, pp. 941–943, 1929.
†Sometimes called the "Apache limestone," though this name is preoccupied by the Cambrian or Algonkian Apache group of southern Arizona.

THE SIERRA DIABLO SECTION[4]

In the Sierra Diablo, which lies to the west of the Delaware Mountains on the opposite side of the Salt Basin, the lower part of the Permian section is more complete than that on the east side of the Basin. Richardson placed the entire Permian section of the Sierra Diablo, as well as the small amount of Pennsylvanian underlying it, in the Hueco formation, which he supposed to be of Pennsylvanian age. It is actually a very heterogeneous group.

*The upper Wolfcamp formation.*—At the base of the Permian here there is a member about 175 feet thick made up of conglomerate, red shale, sandstone, and marly limestone, which in places contains abundant upper Wolfcamp fossils. In the northern part of its exposure it rests with angular discordance on Pennsylvanian limestone, and in the south it lies upon the pre-Cambrian. In many parts of the range, especially to the south, the formation is not present, and here the Hess limestone forms the base of the Permian. The following is a numerical analysis of its brachiopod fauna:

Number of species present—7
Number also in Pennsylvanian—2
Number also in *Uddenites* zone—5
Number also in Glass Mountains Wolfcamp—5
Number also in Hess—2
Number also in Leonard—1
Number also in Gym—4
Restricted to this unit—0
Present in Wolfcamp formation, not *Uddenites* zone—1
Present in *Uddenites* zone, not higher in Wolfcamp—1

The species of brachiopods present are:

Orthotichia kozlowskii
Isogramma millepunctata
Productus dartoni
Productus semistriatus
P. wolfcampensis
Linoproductus cora
Composita subtilita angusta

*The Hess limestone.*—Above the upper Wolfcamp in the northern and central parts of the Sierra Diablo, or resting on the pre-Cambrian, Cambrian, or Ordovician in the southern part of the range and in the Baylor Mountains, are 500 feet of limestone which contain an abundance of *Fusulina* and in the upper part a prolific Hess fauna. Some of the species occurring abundantly in the Hess of the Sierra Diablo are rare in the Hess of the Glass Mountains, and vice versa, but, as in the Hess of the Glass Mountains, *Lyttonia* and *Prorichthofenia teguliferoides* are common. The following is a numerical analysis:

Number of species—23
Number also in Pennsylvanian—2
Number also in *Uddenites* zone—2
Number also in middle and upper Wolfcamp—5
Number also in Hess of Glass Mountains—14
Number also in Leonard—13
Number also in Word—7
Number also in Capitan—2
Number also in Gym—7
Restricted to this unit—2
Present in Wolfcamp, not Hess—4
Present in Hess, not Wolfcamp—13
Present in Hess, not Leonard—4
Present in Leonard, not Hess—2

Among the more important brachiopods are:

Rhipidomella hessensis
Enteletes dumblei
Meekella attenuata
Derbya nasuta
Chonetes biplicatus
C. spinoliratus diabloensis
Productus ivesi
Buxtonia peruviana
Marginifera manzanica
Marginifera? whitei
Prorichthofenia likharewi
P. teguliferoides
Lyttonia nobilis americanus
Spiriferina anculata
Hustedia hessensis

Girty's collections from the "Delaware Mountain formation, Diablo Mountains" probably came from this horizon, for the forms which Girty described in that lot are in general common in this formation.

*The Bone Canyon black limestone.*—On Victorio Peak and to the north of it the Hess is overlain by the Bone Canyon black limestone member of the Leonard formation, averaging about 800 feet in thickness. On the south side of Victorio Peak the black limestone grades into alternating white limestone and calcareous shale of the Glass Mountains facies. In the Baylor Mountains, which lie east of the southern Sierra Diablo, the Hess limestone is overlain by alternating white limestone and shaly limestone of the Glass Mountains facies, quite like that of the southern Sierra Diablo. The Bone Canyon member in the Sierra Diablo and the Leonard in the Baylor Mountains are very fossiliferous at some horizons. The fossils are mainly brachiopods; no ammonoids have been found here. The following table indicates the relationships of the brachiopod forms:

Number of species—28
Number also in Pennsylvanian—1
Number also in Wolfcamp—4
Number also in Hess—20

[4]This section and that of the Hueco Mountains will be discussed in detail by P. B. King and R. E. King in a forthcoming bulletin of the Texas Bureau of Economic Geology.

Number also in Leonard—26
Number also in Word—19
Number also in Capitan—8
Number also in Bone Canyon of Delaware Mts.—15
Number also in Victorio Peak limestone—12
Present in Victorio Peak, not Leonard—1
Present in Leonard, not Hess—6
Present in Hess, not Leonard—0
Present in Leonard, not Word—8
Present in Word, not Leonard—0
Restricted to this unit—1

Among the important species of brachiopods are *Productus ivesi, P. occidentalis, Marginifera manzanica, M. cristobalensis, M. sublaevis, Aulosteges magnicostatus,* and *Spirifer marcoui infraplica.*

*The Victorio Peak white limestone.*—From Victorio Peak northward, the Sierra Diablo is capped by the massive white Victorio Peak member of the Leonard formation, over 1000 feet in thickness. The lower part of this member replaces laterally the upper part of the Bone Canyon member. *Fusulina* is abundant throughout the Victorio Peak limestone, but other fossils are abundant only sporadically. Ammonoids have not been found. The brachiopods are the most common forms in the fauna. An analysis of them gives the following results:

Number of species—19
Number also in Pennsylvanian—0
Number also in Wolfcamp—2
Number also in Hess—10
Number also in Leonard, Glass Mountains—12
Number also in Word—10
Number also in Capitan—2
Number also in Bone Canyon member—12
Restricted to this unit—0
Present in Leonard, not Hess—8
Present in Hess, not Leonard—1
Present in Word, not Leonard—0
Present in Leonard, not Word—9

*Productus ivesi, P. occidentalis, Buxtonia victorioensis, Waagenoconcha montpelierensis, Marginifera manzanica, Spirifer marcoui infraplica,* and *S. pseudocameratus* are the more important species.

*The Delaware Mountain formation.*—In the northern part of the Sierra Diablo a northeastward dip brings the Delaware Mountain sandstone below the summit line of the range, which is determined by the stripped pre-Comanchean surface. About 500 feet of the formation are exposed. Fossils are not abundant. From a layer of sandy limestone which caps several mesas southwest of Dos Alamos a few poorly preserved forms were collected. There were 12 species of brachiopods, of which 1 occurred in the Wolfcamp, 5 in the Hess, 6 in the Leonard, 12 in the Word, and 8 in the Capitan. Seven occurred in the Word but not the Leonard, none in the Leonard but not the Word, 5 in the Word but not the Capitan, and none in the Capitan but not the Word. Among the species are *Meekella skenoides, Linoproductus waagenianus, Marginifera o p i m u s, Aulosteges guadalupensis,* and *Composita emarginata affinis.*

### PERMIAN OF THE DIABLO PLATEAU

The Sierra Diablo is the eastern scarp of the Diablo Plateau. Except in Brackett Draw, a few miles west of the northern end of the Sierra Diablo, where the Victorio Peak member with its characteristic fossils is exposed, the plateau is underlain by a limestone and dolomite of practically homogeneous character, whose fossils are quite unlike any exposed on the escarpment. In the eastern half of the plateau the rock is largely dolomite, with a few fossils. Gypsum lenses appear toward the northeast, and in the northeasternmost part of the Plateau, on the west side of the Salt Flat (Crow Flat), opposite the Guadalupe Mountains, there is a rapid transition to the Chupadera facies of New Mexico, which is characterized by alternating beds of dolomite and gypsum. In the northeastern part of the Salt Basin quadrangle, in the hills south of a small basic igneous plug, there are alternating beds of dolomite and gypsum. The former contain abundant *Productus inca* (loc. 520), a fact which shows that these beds are faunally related to the upper part of the Gym limestone of the Hueco Mountains. The rocks here and elsewhere in the eastern part of the Diablo Plateau are apparently the northwestern facies of those exposed in Brackett Draw and on the Sierra Diablo scarp, and since there is no evidence of uplift which would bring to the surface beds of Hess age, they are probably of Leonard age. However, the faunas of the eastern Diablo Plateau are entirely unlike those of the Sierra Diablo, but closely allied to those of the Gym of the Hueco Mountains and the Chupadera of New Mexico. It therefore seems probable that there was an abrupt facies change along a northeast-southwest line passing a few miles to the northwest of the outcrops in Brackett Draw. Southeast of this line is the normal marine facies of the Sierra Diablo, while northwest of it are deposits of the Gym facies, which in turn change northward into the Chupadera facies.

In the western half of the Plateau and on its western scarp, the Hueco Mountains, the lower beds of the Permian section are again exposed. The dolomite of the eastern part of the plateau is underlain by limestone in the western part.

PERMIAN OF THE HUECO AND FRANKLIN MOUNTAINS

In the Hueco Mountains the Permian consists of massive grey limestone, which may be called the Gym formation because of its similarity faunally and lithologically to the Gym formation of the Florida Mountains, which lie about 90 miles to the west. The Gym is not as abundantly fossiliferous as many formations in the Permian of Trans-Pecos Texas, and its faunas are of a peculiar facies, unlike that of most of the formations of presumably the same age. The faunas indicate a transition to the Chupadera fauna of New Mexico. Three members are recognizable in the Gym of the Hueco Mountains. The lower one is about 500 feet thick; it is characterized by abundant *Schwagerina*. The middle Gym is a darker grey limestone containing many pelecypods and gastropods and a subordinate representation of *Schwagerina*. The upper Gym is characterized by an abundance of *Composita mexicana* and *Pugnoides texanus*.

The lower light grey limestone member of the Gym contains a great abundance of *Schwagerina*. The following is an analysis of the brachiopod fauna of this member:

Number of species—20
Number also in Glass Mountains Wolfcamp—12
Number also in *Uddenites* zone—7
Number also in Pennsylvanian—5
Number also in Hess—11
Number also in Leonard—6
Number also in Word—2
Present in Bolivia—9
Present in Bolivia, not in Pennsylvanian—4
Number occurring higher in Gym, not in Pennsylvanian—4
Number occurring in Pennsylvanian and higher Gym—2
Number in Wolfcamp and not Hess—8
Number in Hess and not Wolfcamp—7

Among the species present in the lower Gym are *Enteletes dumblei, Productus hessensis, P. huecoensis, P. semistriatus, P. wolfcampensis, Buxtonia peruviana, Avonia boulei, Marginifera capaci, Camarophoria deloi, Spirifer condor,* and *Spirifer huecoensis*. The Wolfcamp species appear to be dominant, though Hess elements also are well-marked. From our present knowledge it seems best to assign the lower Gym to the upper part of the Wolfcamp horizon.

The middle dark grey limestone member of the formation is about 250 feet thick. It is especially characterized by a great abundance of gastropods and pelecypods. Brachiopods are less common than in the lower Gym. Their analysis follows:

Number of species—8
Number also in Pennsylvanian—2
Number also in *Uddenites* zone—2
Number also in middle and upper Wolfcamp—3
Number also in Hess—3
Number also in Leonard—2
Number also in Word—2
Number also in Capitan—2
Present in Bolivia—4
Present in Bolivia, not in Pennsylvanian—2
Present in lower Gym—3
Present in upper Gym—3
Present in Wolfcamp, not Hess—2
Present in Hess, not Wolfcamp—3
Restricted to this unit—0

The most common species are *Meekella mexicana, Derbya buchi, Productus inca, Linoproductus cora, L. villiersi, Pugnoides texanus,* and *Composita mexicana.*

The upper light grey limestone member occurs along the summit of the range and in the northwestern part of the Diablo Plateau. It is about 800 feet in thickness. In its lower part there are about 150 feet of red beds, which are exposed in the vicinity of Deer Mountain. *Schwagerina* does not occur as high as this member, but the gastropod fauna of the middle member ranges up into this division. The following table expresses the relations of the brachiopod species:

Number of species—11
Number also in Pennsylvanian—0
Number also in *Uddenites* zone—2
Number also in middle and upper Wolfcamp—2
Number also in Hess—4
Number also in Leonard—4
Number also in Word—2
Number also in Capitan—0
Present in Bolivia—3
Present in lower and middle Gym—7
Present in Wolfcamp, not Hess—3
Present in Hess, not in Wolfcamp—3
Restricted to this unit—1

The most abundant and characteristic fossils of this member are *Composita mexicana* and *Pugnoides texanus. Derbya buchi, Productus inca,* and *Hustedia huecoensis* also are common. The association of *Composita mexicana* with *Pugnoides tex-*

*anus* appears to characterize a widespread faunal facies of this region. The two species occur together abundantly in the Gym at its typical locality in the Florida Mountains, in the Chupadera, in the Bone Canyon member of the Delaware Mountains, and in the uppermost Hess of the Glass Mountains, southwest of the Word Ranch.

On the west side of the Franklin Mountains, north of El Paso, the lower and middle members of the Gym are recognizable.

The correlation of the Gym with the Glass Mountains and Sierra Diablo sections is uncertain. Ammonoids do not occur in the Gym, and many of its brachiopods do not occur in the other faunas of Trans-Pecos Texas. The species which seem to be of significance point to the correlation of the lower Gym with the upper part of the Wolfcamp formation, and of the middle and upper members of the Gym with the Hess.

### PERMIAN OF THE FINLAY MOUNTAINS

South of the central part of the Diablo Plateau is the Finlay Dome, the center of which exposes Permian rocks. At the base of the section there is a conglomeratic limestone, containing *Prorichthofenia teguliferoides*, and therefore is to be correlated with the Hess. There follows about 125 feet of shale, then dark limestone. The limestone contains ammonoids of Leonard age, and the underlying shale has abundant Leonard fossils. Six species of brachiopods were collected from this locality. Of these, one occurs in the Wolfcamp, 2 in the Hess, 3 in the Leonard, 1 in the Word, and none in the Capitan. One form was restricted to that locality. One of the species occurred in the Leonard but not in the Hess, none in the Hess or Word that did not occur in the Leonard, and 3 in the Leonard that did not occur in the Word. Among the species occurring at this place are *Chonetes subliratus, C. molengraaffi?, Productus ivesi,* and *Marginifera sublaevis.*

Permian limestone and gypsum, said to contain Leonard fossils, occur below the Jurassic of the Malone Mountains, which lie about 5 miles to the south of the Finlay Dome.[5]

### PERMIAN OF THE CARIZZO AND WYLIE MOUNTAINS

In the Carrizo Mountains, near Dalberg (southwest of Van Horn), there are about 400 feet of grey Permian limestone, similar lithologically and faunally to the Gym. This limestone has the same stratigraphic position as the limestone of Hess age in the southern Sierra Diablo (which lies about 7 miles to the north), but is faunally quite different. The same facies occurs in the Wylie Mountains, about 10 miles to the east, where about 1400 feet of Permian limestone is exposed. *Productus hessensis* is the only common brachiopod.

### PERMIAN OF THE SHAFTER DISTRICT

Permian rocks are exposed in the Shafter district on the south, east, and north flanks of the Chinati Mountains. They were first studied by Udden* who measured sections along the valley of Cibolo Creek and gave to the whole the name Chinati series, which he subdivided into the Cieneguita, Alta, and Cibolo formations. The Cibolo Creek section was briefly examined by the writer in 1928. In 1929 Baker* published observations on the exposures in Upper Cibolo Basin and Pinto Canyon on the north flank of the Chinati Mountains.

According to Baker the Cieneguita clastic beds contain *Schwagerina* "from the bottom to the top" as well as other fossils which Beede informed him "belong in the base of the Permian or at least in the transition zone." Near Ojo Bonito I measured 2400 feet of shale, sandy shale, and thin limestones, probably belonging to the Alta formation, but with the base cut off by eruptive contact and the top by the Cretaceous overlap. The intercalated limestones contain *Productus ivesi, Perrinites vidriensis* (abundant near Ojo Bonito), and other Leonard fossils. Baker does not report fossils in the Alta formation north of the Chinati Mountains, though it has a great development in both the Upper Cibolo Basin and Pinto Canyon.

I collected Word fossils in the Cibolo formation out of the "thin-bedded zone" and the "yellow limestone" between the Shafter Mine and the Chinati Mountains. North of the Chinati Mountains Baker found *Waagenoceras* and *Medlicottia* in the "lower brecciated zone" of the Cibolo, and *Agathiceras girtyi, Adrianites marathonensis,* and *Stacheoceras gilliamense* in the "thin-bedded zone," also indicating a Word age for the strata. North of Ojo Bonito the Glen Rose limestone has a basal conglomerate in which are cobbles of a white limestone, probably

[5]See Baker, C. L.,: U. of Texas Bull. 2745, p. 11, 1927.

*Udden, J. A., Geology of the Shafter Silver Mine District, Univ Texas Mineral Surv. Bull. 8, 1904.

*Baker, C. L., Note on the Permian Chinati Series of West Texas, Univ. Texas Bull. 2901, pp. 73-84, 1929.

a higher Permian formation no longer exposed in the vicinity.

The following are analyses of my brachiopod collections from strata assigned to the Leonard and to the Word near Shafter:

Leonard

Number of species—26
Number also in Wolfcamp—3
Number also in Hess—16
Number also in Leonard—25
Number also in Word—13
Number also in Capitan—5
Number also in Pennsylvanian—0
Present in Leonard, not Hess—14
Present in Hess, not Leonard—0
Present in Leonard, not Word—12
Restricted to this locality—1

Word

Number of species—5
Number also in Wolfcamp—1
Number also in Hess—2
Number also in Leonard—2
Number also in Word—3
Number also in Capitan—2
Present in Word, not Leonard—2
Present in Leonard, not Word—1
Present in Word, not Capitan—1
Present in Capitan, not Word—0
Restricted to this locality—1

PERMIAN NEAR LAS DELICIAS, COAHUILA, MEXICO

About 275 miles south of the Glass Mountains, in southwestern Coahuila, a thick section of Permian rocks is exposed along the foot of the Sierra del Sobaco, northwest of the hacienda of las Delicias.[8] The lowest member of the Permian is exposed at Las Puerticitas—it is a bluish limestone containing few fossils. It is followed by a considerable thickness of shale, tuff, and basic lava flows; some layers of shale contain *Perrinites vidriensis.* Near the middle of the section are layers of bluish-grey limestone containing abundant fossils. This horizon is well exposed on Cerro Caballo. It is followed by more shale, tuff, and lava. Some layers of the upper shale contain abundant ammonoids, including *Waagenoceras dieneri.* The highest bed of the section is the limestone of Cerro Pichigüia, part of the fauna of which was described by Haack.[9] The limestone of Cerro Caballo has yielded a rather large number of brachiopods, the relationships of which are as follows:

Number of species—17
Restricted to this place—2
Present also in Wolfcamp—1
Present also in Hess—7
Present also in Leonard—10
Present also in Word—15
Present also in Capitan—5

These figures point to a Word age for the horizon. The limestone of Pichagüia gives the following analysis of its brachiopods:

Number of species—6
Restricted to this place—0
Present also in Wolfcamp—0
Present also in Hess—4
Present also in Leonard—4
Present also in Word—4
Present also in Capitan—2

This analysis gives no clue in itself to a definite age for the horizon, but the fact that it overlies the *Waagenoceras* ammonoid horizon indicates that it is of Word age or perhaps younger.

## CORRELATION OF THE PERMIAN SECTIONS OF TRANS-PECOS TEXAS WITH OTHER AMERICAN PERMIAN SECTIONS

### PERMIAN OF THE MID-CONTINENT REGION

*Permian of North-Central Texas.*—In north-central Texas the Cisco, the highest group of the Pennsylvanian, is divided into six formations, of which the highest is the Putnam. The Putnam formation is 125 to 150 feet thick, and consists largely of blue-grey and dark blue carbonaceous shale in the lower part grading up into buff, pink, and red shale; the shales contain three or four lentils of buff and dirty yellow limestone and one or two layers of brown calcareous sandstone. At the top of the Putnam is the Coleman Junction limestone, 3 to 15 feet thick, which has been placed by Drake at the base of the Permian, and by Plummer and Moore at the top of the Pennsylvanian. According to Dunbar,[10] the fusulinids of the Coleman Junction limestone are like those of the Cottonwood limestone of Kansas, which is regarded as the base of the Permian there, so that the Coleman Junction limestone is most properly regarded as of Permian age.

Overlying the Putnam in the Brazos River valley is the Wichita group, the lowest division of the

[8]See Böse, E.: On the Permian of Coahuila, Am. Jour. Sci., ser. 5, vol. 1, pp. 187–191.

[9]Haack, W.: Permfauna aus Nordmexiko, Zeit. d. deutsch. Geol. Gesell. Bd. 66, pp. 482–504, 1915.

[10]Personal communication.

Permian. It is 700 to 1200 feet in thickness, and consists of a series of thick limestone layers separated by thick marine shale and marl beds containing thin limestone lentils. It is divided into three formations, named in ascending order the Admiral, Belle Plains, and Clyde. Thirty-seven kinds of invertebrates have been listed from the Admiral formation, but except for ammonoids, which are said to be of Permian aspect, there is nothing which is particularly diagnostic (**135**, pp. 183–198).

The marine shale and limestone facies of the Wichita group grades northward into red beds, with minor members of bluish and grey-white sandstone and shale and thin layers of limestone, about 800 feet thick. The red bed facies of the Wichita and Red River valleys is therefore quite different from the normal marine facies of the Brazos valley, though of the same age. In this region the base of the Permian is not easily determinable, for the Coleman Junction limestone disappears in southwestern Archer County. The equivalents of the Admiral and Belle Plains formations are dominantly red beds, but the limestones of the Clyde formation continue from the Brazos valley northward into this region.

There are two ammonoid localities* in the Wichita of this region, the faunas of which have been described by C. A. White (**186**) and Emil Böse.[11] One locality, 4 miles south of Dundee, Archer County, is at the horizon of the Belle Plains formation. It is stated by Böse to be probably lower than the other, which is at the old Military Crossing of the Big Wichita River, about 6 miles north of Maybelle, Baylor County. Böse regards the fauna of the first locality as more primitive than that of the second, but the two horizons cannot be many hundred feet apart stratigraphically. The locality south of Dundee contains, according to Böse:

Medlicottia n. sp. (aff. M. artiensis Gruenew.)
Perrinites n. sp. (aff. P. cumminsi White.)
Stacheoceras (Marathonites?) n. sp. (aff. St. romanowskyi Karp.)
Agathiceras sp. ind. (aff. A. uralicum Karp.)

The following species were described from the old Military Crossing of the Big Wichita by C. A. White:

Paralegoceras baylorense (White)
Perrinites cumminsi (White)
Stacheoceras walcotti (White)
Medlicottia copei White

The other fossils, mostly molluscs, listed from the same and adjacent localities by White, are of little value for purposes of correlation. In the Glass Mountains *Perrinites cumminsi* occurs in the upper Wolfcamp ammonoid fauna of Dugout Mountain, and probably other species of the Wichita fauna will be found to be present there when it is completely studied. The presence of *P. cumminsi* is very important, species of *Perrinites* being of great value as index fossils. There can be little doubt that the Wichita ammonoid faunas are of approximately the same age as the upper Wolfcamp fauna. Since the Wichita group starts at the base of the Permian (as defined in central Texas), it therefore seems likely that the Wichita group is approximately equivalent in age to the Wolfcamp formation.

The vertebrate faunas of the Wichita have been reviewed by Romer,[12] who says that "some of the most specialized and advanced forms of the fauna are present in Texas at the very bottom of the Permian." Seventeen genera of amphibia, cotylosauria, pelycosauria, and "protosauria" occur in the Wichita group.

Near the same locality from which the middle Wichita ammonoids were collected, 4 miles south of Dundee and 4½ miles east of Fulda, there is a fossil plant locality, from which David White[13] has listed a large number of species of clearly lower Permian age. "The material in hand from the lower and middle portions of the Wichita puts beyond question the Permian age of that formation. . . . Besides the familiar Permian floras, including *Callipteris, Walchia, Comphostrobus* and *Dicranophyllum* and the abundant *Gigantopteris*, we find new migrants such as *Odontopteris neuropteroides, Odontopteris fischeri*, and *Annularia max-*

*Since this paper was completed further work has been done on the later Paleozoic ammonoids of Central Texas by F. B. Plummer and Gayle Scott. Their work will appear in a succeeding bulletin of the Bureau of Economic Geology, and brief abstracts have appeared in the Bull. Geol. Soc. Amer., vols. 40 and 41, 1929–1930 (Proceedings of the Paleontological Society). So extensive is the new information secured that no satisfactory revision of the present discussion can be attempted.

[11]Böse, E.: Permo-Carboniferous Ammonoids of the Glass Mountains, U. of Texas. Bull 1762, pp. 183–194. Böse stated that the locality south of Dundee was in the upper Wichita, and that the Military Crossing locality was in the lower Clear Fork, but in fig. 7 of U. of Texas Bull. 2801 it may be seen that the first locality is actually middle Wichita and the second upper Wichita. Dundee is not as stated by Böse, in Baylor, but in Archer County.

[12]**139**, pp. 66–108. Also see Schuchert, Charles: **150**, pp. 822–826.

[13]Proc. U. S. Nat. Mus., vol. 41, pp. 493–516, 1912.

*ima.* The odontopterids are Uralian plants, while the *Annularia*, like *Gigantopteris*, is Asiatic.[14]

Above the Wichita is the Clear Fork group, 1100 to 1900 feet thick, consisting of red and blue clays, sandy shales, white and red sandstones, clay-ball conglomerate, and layers of limestone, dolomite, and gypsum. There has been some controversy as to the place at which the Wichita-Clear Fork boundary should be drawn. Some workers have placed the Lueders and Arroyo formations in the Wichita and others in the Clear Fork. Plummer and Moore apparently follow Cummins in placing the Lueders in the Clear Fork group. In ascending order the Clear Fork comprises the Lueders, Arroyo, and Vale formations, the Bullwagon dolomite, and the Choza formation. The Bullwagon dolomite contains in Runnels County the following ammonoids, as identified by Böse[15]:

Perrinites n. sp.
Medlicottia n. sp. (aff. M. orbignyana Vern.)
Gastrioceras n. sp.

The other fossils from the formation are pelecypods and gastropods not specifically identified. The species of *Perrinites* is intermediate in complexity between *P. cumminsi* and *P. hilli*, but is not as complex as *P. compressus*, which occurs in the Hess formation of the Glass Mountains, and is much less highly developed than *P. vidriensis*. The most that can be said, then, concerning the ammonoid fauna of the Clear Fork group, is that it is intermediate between the Wolfcamp-Wichita and the Leonard-Double Mountain types. Because of this fact, as well as the stratigraphic position of the Clear Fork, it seems most logical to correlate the Clear Fork approximately with the Hess.

Romer lists 21 genera of amphibia, cotylosauria, pelycosauria, and "protorosauria" from the Clear Fork group, most of them from the Arroyo formation. Romer places the Lueders formation in the Wichita group, but states that its vertebrate fauna is more closely allied to that of the Clear Fork. In fact, two distinct vertebrate faunal zones are distinguishable, of which the lower, occurring in the lower and middle Wichita, and the upper Cisco, is characterized by the presence of *Cricotus*, while captirhinomorphs *(Captorhinus, Labidosaurus)*, *Seymouria*, and a number of other forms gives a criterion for the upper fauna. Small fossil plant collections have been made from the Clear Fork; they represent the lower Permian *Gigantopteris* flora.

Above the Clear Fork is the Double Mountain group, which varies in thickness from 200 feet in the vicinity of San Angelo to more than 1000 feet near the Red River. It is characterized by sandstones, thick gypsum beds, and some sandy shales and earthy limestones. At the base is the San Angelo formation, 125 to 175 feet thick, consisting of red sandstone, white sandstone, conglomerate, and shale, which rests unconformably on the underlying Choza formation of the Clear Fork. The only fossils are a few pelecypods and gastropods.

The San Angelo is followed by the Blaine formation, from 80 to 275 feet thick, which in the Colorado and Concho river valleys consists of cream-colored gypsiferous and calcareous sandstone, and farther north is composed of thick gypsum beds with some shale partings and thin dolomite. In Tom Green County it rests unconformably on the San Angelo formation. At three localities there are ammonoids in the Blaine. At the falls on Salt Crotan Creek in Kent County, *Perrinites hilli* (Smith) occurs with *Stacheoceras* sp. and *Medlicottia* sp. Smith has cited *Medlicottia copei*, the same species that occurs in the Wichita, from near San Angelo. Böse[16] lists large *Perrinites* n. sp. and large *Gastrioceras* n. sp. from Quanah, near the Red River. The only important species among those mentioned above, and the only one positively identified, is *Perrinites hilli*, which is exceedingly close to *P. vidriensis* of the Leonard formation of the Glass Mountains. Because of the absence of other significant fossils in the Double Mountain group in Texas, this species has a great deal of importance. The only similar species is *Perrinites vidriensis*, a form which does not occur above the Leonard. In the Word, *Perrinites* is unrepresented, but its descendant *Waagenoceras* is present. If the Blaine were of Word age, one would expect to find in it *Waagenoceras*, a fossil which does not occur at a lower horizon in the Glass Mountains section. Ordinarily, correlation on the basis of a single genus is not conclusive, but ammonoids of this type are very exceptionally good index fossils, and in view of the absence of other criteria for paleontologic correlation it is fortunate that such

[14]White, David, in The Permian of Western America from the Paleobotanical Stand-point. Proc. Second Pan-Pacific Congress, pp. 1060–1061, 1926.

[15]Op. cit., pp. 194–206.

[16]Op. cit., pp. 207–210.

a definite horizon marker is present. A check on the correlation of the Blaine with part of the Leonard is made possible by the subsurface tracing of the Blaine into the San Andres formation of New Mexico,[17] which in turn is very definitely correlated on stratigraphic and paleontologic evidence with the Leonard formation of the Glass Mountains. A further circumstance favoring correlation of the San Angelo and Blaine with the Leonard is the evidence of uplift at the base of the Double Mountain as shown by the unconformity at its base and the conglomerate in the San Angelo formation, which may correlate with the evidence of uplift of Llanoris in the lower part of the Leonard formation: the chert and quartz pebble conglomerate of the lower Leonard. It is highly probable that the uplift was contemporaneous in these two regions.

No vertebrate remains have been found in the Double Mountain group, and the only plants are small collections from the San Angelo formation of the *Gigantopteris* flora, of lower Permian age.

In north-central Texas the Blaine is unconformably overlain by the non-marine Triassic or by the Trinity sand.

*Permian of Oklahoma and the Texas Panhandle.*—In Oklahoma the Wichita group is divided[18] into the Stillwater and Wellington formations. The former, about 800 feet thick, has at its base the Cottonwood limestone, which probably is a time equivalent of the Coleman Junction limestone at the base of the Permian of north-central Texas. The typical Stillwater is a series of red and gray sandstones and red shales, but toward the north limestones appear and gradually thicken, and the red sandstone and shale grades into grey shale; this northward facies is the same as that of Kansas. Thus the northward change from non-red to red beds in the Wichita of Texas is duplicated by a change of the same kind in the opposite direction in Oklahoma. This fact is interpreted by Beede as due to the derivation of the Wichita red beds from the Wichita and Arbuckle mountains,[19] but these ranges seem to be hardly large enough to have supplied such a large amount of sediment. In southern and central Oklahoma the Wellington formation consists of red shale and sandstone and grey "mud-stone." Toward the north there is a transition to the grey and bluish shales with thin beds of "mud-stone," with a thickness of about 600 feet. This northern facies is, again, like that of Kansas. The Wichita group of Oklahoma does not contain fossils which enable correlations to be made with the Trans-Pecos sections, the fauna being largely of pelecypods and gastropods. Near Perry, Oklahoma, there are plants in the Wellington which are of definite lower Permian age. Lower Permian vertebrate remains also have been found in the Wichita.

The Clear Fork group is divided into two formations, the Garber below and the Hennessey above. The former, about 600 feet thick, consists of red clay shales, red sandy shales, and red sandstones. The latter, about 400 feet thick, is a red clay shale, without sandstone.

The Double Mountain group comprises a large number of formations.[20] At the base is the Duncan formation, consisting of two or three ledges of heavy white or buff sandstone separated by shale, varying from 100 to 300 feet in thickness. It traces southward into the San Angelo formation of Texas. Above the Duncan is the Chickasha gypsiferous red clay-shale, containing thin sandstones, varying up to 350 feet in thickness. The overlying Blaine gypsum, the ammonoid horizon of the Texas Double Mountain, is a zone of gypsum ledges with interbedded shales aggregating a thickness of about 50 feet. The next member, the Dog Creek shale, is a red clay-shale containing dolomite beds, from 30 to 400 feet thick. The Whitehorse sandstone is a friable red sandstone averaging 200 feet thick. The Day Creek dolomite is a widespread member from 1 to 5 feet thick. The Cloud Chief gypsum, containing interstratified red clay shale, varies up to 115 feet thick. The highest formation is the Quartermaster, consisting of 300 feet of soft, red sandstone, sandy clay, and shale.

The only fossils in the Double Mountain group of Oklahoma or the Texas Panhandle are in the Whitehorse sandstone, from which Beede (**5**, pp. 115–171) has described a fauna of pelecypods and gastropods, as well as a single species of brachiopod, *Dielasma schucherti* Beede. This has its

[17]Gould, C. N., and Willis, Robin: Tentative Correlation of the Permian Formations of the Southern Great Plains, Bull. Geol. Soc. Amer., vol. 38, pp. 435–438, 1927.

[18]Aurin, F. L., Officer, H. G., and Gould, Charles N.: Subdivisions of the Enid formation, Bull. Am. Assoc. Pet. Geol., vol. 10, pp. 786–799, 1926.

[19]Beede, J. W.: Origin of the Sediments and Coloring Matter of the Red Beds of Oklahoma, Science, n. s., vol. 35, pp. 348–350, 1912.

[20]Gould, C. N., and Lewis, F. E.: The Permian of Western Oklahoma and the Panhandle of Texas, Okla. Geol. Sur. Circular 13, 1926.

counterpart in the Glass Mountains faunas in *Dielasma schucherti minor* King, which differs from the Whitehorse sandstone species only in its smaller size. The presence of this form in the Word formation may indicate that the part of the Double Mountain group above the Blaine is of Word age. None of the other Whitehorse fossils is known at present to be of correlation value. The correlation made by Beede of the Whitehorse fauna with that of the Rustler limestone near Lakewood, New Mexico, is based more on similarity of facies and the presence of similar genera than on specific identities. The pelecypods of the Permian are characteristic of the limestones and sandstones intercalated with gypsum and red beds, and similar forms occur in this facies in rocks of various ages throughout the Permian. Where, on the other hand, a fossil occurring in the pelecypod facies occurs also with a normal marine fauna of brachiopods, corals, bryozoans, etc., more weight may be attached to it, for it permits the correlation of two unlike faunas. Since the classification is based on the succession of normal marine faunas, any species occurring in both it and the pelecypod facies is very important. Thus *Dielasma schucherti* is exceptionally significant because it occurs in the pelecypod facies and as a smaller variety in the normal marine facies (Word). Similarly, the ammonoids of the Blaine occur in the normal facies in the Leonard formation.

Neither plants nor vertebrates have been found in the Double Mountain of Oklahoma and the Texas Panhandle.

In conclusion, it may be pointed out that though the faunal evidence for correlating the Oklahoma section with that of Trans-Pecos Texas is meagre, fairly definite correlations are made possible indirectly by means of the equivalents of the Oklahoma formations in north-central Texas. Therefore, it is known that the Wichita is approximately equivalent to the Wolfcamp formation, the Clear Fork to the Hess, and the Double Mountain as high as the Blaine to the Leonard. The presence of related forms of *Dielasma* in the Whitehorse and the Word strongly suggests that the upper part of the Oklahoma-Texas Panhandle section is equivalent to part of the Word.[21]

*Permian of Kansas.*—The upper formation of the Pennsylvanian of Kansas is the Wabaunsee. The Permian types of fossils are there introduced. The total thickness of the Wabaunsee is 500 feet. Three hundred and fifty feet above the base is the Americus limestone. In the immediately succeeding lower beds of the Elmdale shale there is the first appearance of *Fusulina*, in the form of *F. longissimoidea.* This is rather a striking change, as the lower horizons are characterized by *Triticites*, while *Fusulina* is characteristically Permian. About 125 feet higher is the Neva limestone, which contains, according to Beede and Kniker (**6**, pp. 43–52), *Schwagerina kansasensis, Meekella mexicana* Girty, *Allorisma capax* Newberry?, and *Omphalotrochus?* sp., as well as other forms which do not occur at a lower horizon, many of them undescribed. The Eskridge shale, 30 to 40 feet thick, separates the Neva from the Cottonwood limestone, which is generally regarded as the base of the Permian, because at that horizon and above the Pennsylvanian elements drop out of the fauna, leaving a gastropod fauna of Permian aspect. *Fusulina emaciata* is common in the Cottonwood limestone, and *Triticites ventricosus* is a very characteristic species of the upper Wabaunsee and lower Permian (as the Permian is usually defined in that area).

The fusulinids in the Kansas faunas are the only fossils in that section which make possible correlations with Trans-Pecos Texas. *Schwakerina kansasensis* occurs also in the middle part of the Wolfcamp formation immediately above the *Uddenites* zone. *Triticites ventricosus* is abundant in the *Uddenites* zone. These facts would seem to indicate that the *Uddenites* zone is of the age of the beds immediately below the Neva limestone, the horizon of the Neva limestone occurring in the Glass Mountains directly above the *Uddenites* zone. If so, the base of the Permian in the entire mid-continent region has been drawn too high, and the contact should actually be placed approximately at the Americus limestone, with the first appearance of the Permian type of fusulinids.[22]

The evidence of fossil plants bears out that of the invertebrates.[23] At the level of the Neva limestone there is a plant association, embracing typical forms that can hardly be older than the Permian as recognized in western Europe. But still lower, in the Elmdale, there is a very late Odontopterid association "which would be referred to

[21]Schuchert **150**, p. 819 gives very similar correlations.

[22]Schuchert **150**, draws the line higher, at the base of the Neva limestone, in harmony with the conclusions of Drs. White and Beede.

[23]White, David: *Permian of Western America from the Paleobotanical* Standpoint, Proc. Second Pan-Pacific Congress, pp. 1057–1058, 1926.

the Permian by some paleobotanists, but which we regard as very latest Pennsylvanian. Also a few plants gathered by me from a level probably as low as the Americus limestone embrace fragments possibly representing Permian forms." Thus we see that the Permian floral elements were introduced at the same horizon as the genus *Fusulina*—at the Americus limestone.

It will now be shown that other evidence exists for placing the Pennsylvanian-Permian boundary below the Neva. In the Ardmore basin and north of the Arbuckle Mountains[24] the lowest formation resting on the folds of the Arbuckle Mountains is the Hart limestone, which forms the base of the Stratford formation. The Hart limestone traces northward probably into the upper part of the Buck Creek formation of northern Oklahoma, which in turn underlies the Elmdale shale. Therefore, the physical break representing the folding of the Arbuckles occurs at a position approximately equivalent to that of the appearance of *Fusulina* and the Permian flora, and to the horizon deduced as probably that of the *Uddenites* zone. Moreover, the upper part of the Vanoos formation, which underlies the Hart limestone, contains a flora in which David White identified *Walchia pinniformis*, a type generally regarded as diagnostic of the lower Permian. This evidence indicates the very late Pennsylvanian age of the beds immediately below the unconformity. If the beds below the unconformity are so high in the Pennsylvanian, it is the more likely that the first beds above the unconformity would be Permian. An early Permian age for the first beds above the unconformity accords very well with the geologic history of the Marathon region, and it is not at all improbable that the folding was simultaneous in the two areas.

A still further evidence favoring lowering of the Pennsylvanian-Permian boundary is the fact that red sedimentation begins in Texas with the Putnam formation, which is placed by Plummer and Moore tentatively in the Cisco, though it is physically more similar to the Wichita than to the underlying beds of the Cisco. The base of the Putnam would approximately correspond to the horizon of the Americus limestone of Kansas. Romer (**139**, p. 97) states that in the Putnam there are Permian types of vertebrates, closely allied to those of the overlying Wichita.

In summary, it may be noted that in Kansas, Oklahoma, and Texas there are facts which indicate that the Pennsylvanian-Permian boundary should be shifted downward to approximately the horizon of the Americus limestone.

The remainder of the Permian section of Kansas is of less interest in connection with the present study. The Wichita equivalent is known as the Big Blue group. The equivalents of the Stillwater formation of Oklahoma are the Council Grove, Chase, and Marion, aggregating about 550 feet in thickness, and consisting of shale with layers of limestone and cherty limestone. The fauna differs from that of the underlying beds in containing fewer brachiopods and a large number of pelecypods, some of which are new types of Permian affinity. The Wellington formation consists of about 500 feet of blue-grey shale containing thick salt beds. The beds above the Wellington are known collectively as the Cimarron. The first member of the Cimarron is the Harper red sandstone, about 325 feet thick, the lower part of which is equivalent to the entire Clear Fork and the upper to the Duncan-San Angelo, the basal part of the Double Mountain. The Chickasha formation of Oklahoma is equivalent to the Salt Plain, Cedar Hills, and Flower Pot members, which consist of red sandstone and shale with layers of gypsum. This is followed by the Blaine gypsum and the remainder of the section as in the Double Mountain of Oklahoma (**122**, pp. 47–73).

Dunbar (**36**, pp. 187–188) has pointed out that two very distinctive species of molluscs, *Stenopoceras dumblei* and *Myalina copei*, occur in both the Chase formation of Kansas and the Wichita group of Texas. In the basal part (Wreford) of the Chase group there are migrant European species of the plants *Callipteris*, *Taeniopteris*, *Walchia*, *Sphenophyllum*, and *Odontopteris*, everywhere recognized as characteristic of the Permian. In the upper Wellington there are a few Pennsylvanian survivors associated with species of *Callipteris*, *Taeniopteris*, *Ullmania*, *Voltzia*, *Araucarites*, and *Glenopteris*.[25] A very large fauna of insects of lower Permian type has been collected from a thin limestone in the Wellington shale (**36**, pp. 171–209).

[24]Morgan, George D.: Geology of the Stonewall Quadrangle, Oklahoma, Bur. of Geol. Bull. 2, 1924. Also Tomlinson, C. W.: Pennsylvanian System in the Ardmore Basin, Okla. Geol. Sur. Bull. 46, pp. 55–56, 1929.

[25]White, David: Permian of Western America from the Paleobotanical Stand-point, Proc. Second Pan-Pacific Congress, pp. 1057–1058, 1926.

PERMIAN OF ARIZONA AND NEW MEXICO

*The Permian of southwestern New Mexico and southeastern Arizona.*—The Permian rocks in this region are of the Gym facies, resembling the Permian of the Hueco Mountains escarpment. In the mountains of Luna County, near Deming, New Mexico (**15**, pp. 35–39), the massive light-grey Gym limestone is about 1000 feet thick. The type locality is Gym Peak, in the Florida Mountains, 17 miles south-southeast of Deming. The formation rests with considerable structural unconformity on the underlying strata. In places it rests on the lower Ordovician El Paso limestone, elsewhere on the upper Ordovician Montoya limestone, while in the Cooks Range north of Deming it overlies shale of Pennsylvanian age. The unconformity below the formation is therefore comparable in every respect to that below the Gym in the Hueco Mountains.

The following invertebrates have been listed by Girty from the Gym of this area (**15**, pp. 38–39):

Fusulinella sp.
Echinocrinus ornatus
Meekella mexicana?
Chonetes platynotus?
Productus aff. P. semireticulatus
P. occidentalis
Linoproductus cora
Marginifera splendens?
Pugnoides osagensis
Ambocoelia? sp.
Squamularia perplexa
Composita subtilita
C. sp.
C. mexicana?
Astartella? sp.
Parallelodon politus?
Edmondia sp.
Monopteria marian?
Myalina sp.
Schizodus sp.
Pinna peracuta
Pleurophorus sp.
Plagioglypta canna?
Bellerophon crassus
B. crassus wewokanus?
B. majusculus?
Bucanopsis modesta
Pleurotomaria texana
P. 3 sp.
Murchisonia 4 sp.
Euomphalus aff. E. pernodosus
Meekospira sp.
Discohelix? n. sp.
Rhynchomphalus obtusispira
Sphaerodoma aff. S. humilis
S. aff. S. primitenia
Cyclonema sp.
Glyptobasis? sp.
Orthonema socorroense?
O. sp.
Pseudomelania? 4 sp.
Zygopleura n. sp.
Loxonema? 2 sp.
Phymatipher n. sp.
Naticopsis sp.
Bulimorpha inornata

The uncertainty of some of the identifications of the species and the lack of specific determination of most of them makes their use difficult, but certain general comparisons may be made. The *Fusulinella* sp. is probably not a true *Fusulinella* but it is likely the same as the form occurring abundantly in the lower Gym of the Hueco Mountains. *Meekella mexicana* occurs in the middle Gym of the Hueco Mountains. The *Productus* aff. *P. semireticulatus* might well be *P. inca,* the common fossil of the middle and upper Gym of Texas, while *Linoproductus cora* is common also in the middle Gym. *Pugnoides osagensis* probably corresponds to *P. texanus,* one of the most characteristic Gym fossils in Texas. The species of *Composita* correspond to *Composita subtilita* and *C. mexicana,* two of the most common species in the Gym of Texas. The great abundance of pelecypods and gastropods is a further resemblance with the Gym faunas of the Hueco Mountains. The species in the Texas section have not as yet been identified. The only notable difference between the fauna of the Gym of Luna County and of the Hueco Mountains is that in the latter there is a great abundance of *Schwagerina,* which has not been reported from the Luna County section.

In southeastern Arizona the Permian has not been differentiated, in mapping, from the Pennsylvanian. Both systems are included in the Naco limestone, which is 1500 to 3000 feet thick. This formation is largely light-grey, compact limestone in beds of moderate thickness. The Pennsylvanian part of the formation has a fauna entirely like that of the Pennsylvanian of the Hueco Mountains, with *Chaetetes milleporaceus* a common fossil. The Permian part, on the other hand, is faunally very similar to the Gym. Girty has listed the following forms from it in the Bisbee district[26]:

Fusulina cylindrica
Michelinia? sp.
Lophophyllum cf. proliferum
Echinocrinus several sp.
Productus, semireticulatus type
Worthenia sp.
Murchisonia? several sp.
Euomphalus sp.
Omphalotrochus, several sp.
Productus cf. norwoodi
Marginifera cf. wabashensis
Martinia sp.
Composita subtilita
Plagioglypta cf. canna
Cyclonema sp.
Orthonema sp.
Polyphemopsis sp.
Bellerophon cf. crassus
Euphemus sp.

On plate 11, fig. 1, of Professional Paper 21 there is an illustration of the form identified as *Fusulina cylindrica.* It is a large robust fusulinid, having exactly the appearance of the *Schwagerina* of the lower part of the Hueco Mountains Gym. A coral similar to *Michelinia* is common in the lower Gym of Texas, as is *Echinocrinus.* One of the forms identified by Girty as *Productus semireticulatus* is

[26]Girty, in Geology and Ore Deposits of Bisbee Quadrangle, Arizona, U. S. Geol. Sur. Prof. Paper 21, pp. 50-54, pl. 11, 1904.

figured on Plate 11, fig. 4; it is indistinguishable from *P. inca* of the Texas Gym. One of the species of the semireticulate type of *Productus* is said to be related to *P. ivesi* and another to *P. occidentalis.* One species of *Omphalotrochus* is probably *O. whitneyi* Meek, while another is related to *O. obtusispira* (Shumard), first described from the Gym of the Hueco Mountains. The species of *Echinocrinus* are said not to be related to Pennsylvanian forms, but to be more suggestive of the Kaibab. Girty states definitely that the fauna is very closely similar to that of the Hueco Mountains.

The Permian part of the Naco limestone is evidently rather widely distributed in southeastern Arizona. About 50 miles west of Bisbee, near the Mowry Mine, 8 miles south of Patagonia, there is a limestone containing the same fauna, the fossils of which have been listed by Darton (**16**, p. 73). It succeeds a Pennsylvanian fauna. Northwest of Patagonia and about 40 miles south of Tucson, there is fossiliferous lower Permian in Sawmill Canyon in the Santa Rita Mountains near the Onyx Mine. A collection from there made by Professor Schuchert and kindly turned over to me for study contains *Productus occidentalis, Camarophoria deloi, Hustedia mormoni,* and *Composita mexicana,* which indicates a Gym age for this horizon. A collection which I have studied from Snyder's Hill, 9 miles southwest of Tucson, made by Professor Schuchert, contains *Productus occidentalis, Avonia boulei, A. subhorrida rugatula, Composita mexicana, Camerophoria deloi,* and *Squamularia guadalupensis.* Girty also has identified *Pustula* aff. *P. porrecta (=Buxtonia peruviana)* from this locality. All these fossils are very characteristic of the Gym. This horizon is underlain by typical Pennsylvanian. Permian fossils have been listed by Darton (**15**, p. 75) from the Waterman Mountains, 30 miles west-northwest of Tucson.

According to Stoyanow (**161**, pp. 320–324) the Gym gastropod fauna occurs in the Naco limestone of the Little Dragoon Mountains, about 50 miles east of Tucson. *Bellerophon crassus, B. percarinatus, B. carbonarius, Euconispira taggarti, Murchisonia terebra, Straparollus pernodosus,* and *S. subquadratus* are listed. Farther to the northwest, however, in the Galiuro Mountains near Winkelman, the lower Pennsylvanian is well exposed, and the Gym equivalents appear to be absent from this region.

The fauna of the highest part of the Naco limestone is of particular interest. From this horizon 5 miles south of Tombstone, Darton collected *Meekella pyramidalis?, Productus leei?, P. ivesi,* and *Marginifera* sp., evidently a Kaibab-San Andres-Leonard fauna. In the Chiricahua Mountains, in southeasternmost Arizona, near Paradise, Stoyanow (**161**, pp. 318–319) found that above the Naco limestone there is a solid quartzite over 100 feet in thickness, overlain by an indistinctly bedded light-grey limestone, about 500 feet thick, containing the Kaibab fauna. *Productus ivesi, P. occidentalis, Meekella pyramidalis, Derbya* sp., *Squamularia guadalupensis,* and *Aviculopecten coloradoensis* are present in this fauna. From these facts, it appears that the upper Naco limestone, equivalent to the Gym of the Hueco Mountains and southwestern New Mexico, is here overlain by beds of Leonard-San Andres-Kaibab age. This makes still more probable the correlation of the Gym with the Yeso formation of central New Mexico and with the Hess of the Glass Mountains section. One is tempted to believe that the quartzite underlying the Kaibab fauna in the Chiricahua Mountains is equivalent to the Coconino of the Grand Canyon region and the Glorieta of central New Mexico.

It is to be hoped that future workers in southeastern Arizona will differentiate the Permian from the Pennsylvanian. If so, it will be appropriate to extend the name Gym to these Permian limestones, because their lithology and fauna are the same as that of the typical Gym and its equivalent in the Hueco Mountains.

Permian limestones are known to occur in northeastern Sonora in the mountains south of Bisbee and Douglas, but their stratigraphy and faunas have not been made known.

*Permian of central New Mexico.*—North of the Hueco and Guadalupe mountains, in the central and south-central parts of New Mexico, there are about 2500 feet of Permian sediments of a quite different lithologic character from the Permian of either the Guadalupe or the Gym facies.[27] The New Mexican section has been called the Manzano group, but it is best to discard this name, for at the type locality of the group, the Manzano range, only the lower member of it, the Abo sandstone, is present.

The Abo red sandstone, 400 to 800 feet thick, rests unconformably on the pre-Permian rocks,

[27] See Darton, N. H.: The Permian of Arizona and New Mexico, Bull. Am. Assoc. Petr. Geol., vol. 10, pp. 833–852, 1926.

lying on limestone of the Magdalena group (Pennsylvanian) except in the Zuni Mountains and part of the Nacimento range, where it overlaps upon the granite, and in the Pedernal Mountain region, where it abuts against pre-Cambrian quartzite. In most places the basal contact of the Abo is marked by little structural discordance, but by sharp change in the nature of sedimentation from that in the Magdalena, namely local conglomerates, and some channeling. Southward from central New Mexico toward the Texas boundary the Abo thins, and several miles north of the state line it has entirely wedged out between the Pennsylvanian and the overlying Permian.

The invertebrate fossils of the Abo are largely pelecypods and gastropods. A collection of ammonoids, said to be from the Abo, made by Baker east of Tularosa and studied by Böse (**9**, pp. 57–60), are of Pennsylvanian type, but according to Darton they may have been obtained from a red sandy member in the upper part of the Magdalena. The brachiopods described from the Abo by Girty (**98**) are mostly Pennsylvanian species, but Darton states that many of these actually came from the Magdalena. Fossil plants from the lower conglomeratic part of the Abo near Pecos City and Glorieta are said by David White to disclose a rather limited basal Permian flora. The upper Abo near Coyote has yielded many vertebrate fossils, comprising the genera *Diadectes, Spehenacodon, Ophiacodon, Clepsydrops, Elcabrosaurus, Diasparactus, Aspidosaurus,* and *Limnocolis.* They have been regarded as Permian types, but Romer (**139**, p. 98) doubts this. Few of the genera are present in the late Paleozoic vertebrate faunas of Texas, and those which do occur there are inconclusive. Romer's argument that the vertebrate fauna is probably of pre-Wichita age seems rather unconvincing. There is thus little direct evidence as to the age of the Abo, but since it contains Permian plants and is conformably overlain by the Yeso formation of Hess age while separated from the underlying Magdalena by a considerable unconformity, it is probably of Wolfcamp age. There can be little doubt that the pre-Abo unconformity is to be correlated with the Marathon disturbance in the Marathon region, the Pennsylvanian-Permian unconformity in the Hueco and Diablo mountains, and the break below the Gym in Luna County.

The Chupadera group of Darton comprises several members. The lower 500 to 800 feet of these beds was called by Lee the Yeso formation; it consists of red shale, pink and yellow sandstone, and thin limestone and gypsum layers. The upper part of this formation is generally termed the Glorieta sandstone, after its exposure on Glorieta Mesa. Above is the San Andres limestone of Lee, a massive white formation 500 or more feet in thickness. Darton states that these members are not continuously traceable, and therefore not natural groups; he prefers to refer to them collectively as the Chupadera group. The faunas of the Yeso and the San Andres as described by Girty (**98**) are, however, quite distinct from each other.

The following Brachiopods have been described from the Yeso:

| | |
|---|---|
| Meekella mexicana | Pugnoides osagensis pusilla |
| Productus ivesi | Squamularia perplexa |
| P. mexicanus? | Composita mexicana |
| P. nebraskensis (=Buxtonia sp?) | C. subtilita? |

*Meekella mexicana* is abundant in the middle Gym of the Hueco Mountains. *Productus nebraskensis* of Girty is probably a species of *Buxtonia,* perhaps *B. peruviana,* which is characteristic of the Gym. *Pugnoides osagensis pusilla* was not identified from the Gym, but may be related to *P. texanus. Composita mexicana* and *C. subtilita* are common Gym fossils. The only similarity to the Leonard formation is the presence of *Productus ivesi,* which occurs also in the Hess. The Yeso is therefore rather definitely of Gym and Hess age. The outcrop of the Yeso formation is continuous with that of the Gym, and near the Texas-New Mexico boundary there is a transition from the lithology of one to that of the other, the red beds and gypsum of the Yeso being replaced southward by limestone of the Gym facies. According to White,[28] the Yeso contains abundant *Taeniopteris, Walchia, Gomphostrobus,* and *Callipteris,* with several forms of *Sphenopteris* and *Pecopteris* identical with those of the Permian of western Europe.

The San Andres limestone has yielded the following brachiopods, described by Girty:

| | |
|---|---|
| Meekella mexicana | Marginifera manzanica |
| Productus ivesi | M. cristobalensis |
| Productus inca (=P. leei) | Squamularia perplexa? |
| Productus occidentalis | Composita mexicana |
| Productus mexicanus? | C. subtilita? |

[28]Permian of Western America from the Paleobotanical Standpoint, Proc. Second Pan-Pacific Sci. Congress, p. 1064, 1926.

*Productus ivesi, P. occidentalis, Marginifera manzanica, M. cristobalensis, Squamularia guadalupensis (=S. perplexa* as identified by Girty), *Composita mexicana,* and *C. subtilita* occur in the Leonard formation of the Glass Mountains. *Productus occidentalis* and *Squamularia guadalupensis* occur also in the Word and the Capitan, but none of the species occurring in the San Andres is restricted to higher beds than the Leonard, with the possible exception of *Productus mexicanus,* which is doubtfully identified. *Productus ivesi* does not occur above the Leonard, and its zone of abundance is in the Leonard, while the two species of *Marginifera* are definite indices to the Leonard in Trans-Pecos Texas. *Productus inca* occurs in Texas in the middle and upper Gym, which is certainly not younger than Leonard in age. In view of these facts, a correlation of the San Andres limestone with the Delaware Mountain or Capitan seems to be rather conclusively disproved, and its correlation with the Leonard fairly certain.

An actual physical connection between the Manzano and Guadalupe groups exists in southeastern New Mexico in the country between the Sacramento and Guadalupe mountains, and considerable geologic work has been done in this area. Unfortunately, there has been no general agreement as to the correlations between the two districts, though the interpretation which has been most widely advanced is that the Chupadera formation is the equivalent to the Guadalupe group* and, more specifically, that the San Andres is laterally continuous with the lower lagoonal member of the Capitan along the western side of the Guadalupe Mountains†. So confidently has this assertion been made, indeed, that Willis‡ is led to conclude that, in spite of the "disagreement between the paleontologists and the men who have studied the Guadalupian Permian in the field . . . . it seems reasonable to correlate the San Andres with the lower Capitan."

Many geologists find, with Lloyd,§ an explanation to these apparent contradictions of stratigraphy and paleontology by the easy recourse to the assumption that the Capitan and other reef barriers shut off the Guadalupian fauna from the interior Manzano seas, where faunas of another facies existed. Though this assumption is superficially reasonable, it is made without understanding the nature and distribution of the fossils concerned. We have already shown how the majority of the San Andres brachiopods occur far below the level of the Capitan formation in Texas, in the Leonard formation, to which many of them are restricted. In addition, most of the Phosphoria brachiopods occur in the Delaware Mountain and Word formations in Trans-Pecos Texas, and, as will be shown below, the Phosphoria formation clearly overlaps the Kaibab (northwestern extension of the San Andres) in central Utah.

In view of the lack of adequate maps of the northern Guadalupe Mountains, the meagerness of published geologic observations, and the almost complete absence of fossil determinations, it is difficult to evaluate the evidence on which the various stratigraphic conclusions are based. For these reasons the following analysis of the literature is attempted with some hesitancy.

We have shown in the description of the Delaware and Guadalupe mountains how the great Capitan reef mass passes northwestward into three lagoonal members: thin-bedded limestones below (not satisfactorily named), medial sandstone and gypsum (Queen and Seven Rivers members) and thin-bedded limestone above (the Carlsbad member)*. The Seven Rivers and Carlsbad members extend northeast down the back slope of the Guadalupe Mountains to the Pecos River north of Carlsbad, where the Carlsbad member has largely disappeared and the medial gypsums and red beds have attained considerable thickness. Since these two members do not extend into the northwest part of the mountains they are of no further concern to us. Most of the northwest part of the mountains is spread over by a great sheet of the lower lagoonal limestones, which are also downfaulted on the west. Limestones extend from here northwestward into the Sacramento Mountains. Part of these belong to the San Andres formation, for fossils characteristic of that formation were collected by

*Darton, N. H., and Reeside, J. B., Guadalupe Group, Bull. Geol. Soc. Amer., Vol. 37, p. 419, 1926.

†Blanchard, W. G., and Davis, M. J., Permian Stratigraphy and Structure of Parts of Southeastern New Mexico and Southwestern Texas, Bull. Amer. Assoc. Petrol. Geol., Vol. 13, pp. 971–972, 1929.

‡Willis, Robin, Preliminary Correlation of the Texas and New Mexico Permian, Bull. Amer. Assoc. Petrol. Geol., Vol. 13, p. 1022, 1929.

§Lloyd, E. Russel, Capitan limestone and Associated Formations of New Mexico and Texas, Bull. Amer. Assoc. Petrol. Geol., Vol. 13, p. 665, 1929: "The San Andres . . . . contains a marine fauna which is . . . . quite different from the Guadalupian fauna of the Capitan. The difference in the fauna may be explained by differences in environment."

*For areal distribution see Darton and Reeside, *op. cit.,* figure 1, or Darton **17,** Plate 49.

Darton* at Russel Gap, 40 miles north of the state line, and farther north by Richardson* on the lower Penasco River. A large and abundant San Andres fauna is also known from near the summit of the Sacramento Mountains, at Cloudcroft,* which is definitely within the Manzano area.

Since the upper limestones are down-faulted along the western face of the Guadalupe Mountains, the lower beds are hidden in some of the most critical places and less is known of their stratigraphy. Our information is based on fortuitous local exposures resulting from structural accidents or deep erosion.

We have already remarked on the peculiar stratigraphic relations along the west side of the Guadalupe Mountains in Texas, where the Delaware Mountain formation rests with marked erosional unconformity on the Bone Canyon and Victorio Peak members of the Leonard formation. The limestone of probable Victorio Peak age thickens northward and at the state line Darton and Reeside say it has a "Chupadera character" (fig. 4). At the same time these lower beds rise northward in altitude with the Delaware Mountain sandstone layers disappearing one at a time by overlap against them, until that formation is only 100 feet thick at the state line (Darton and Reeside). Beyond this are no continuous exposures.

Darton and Reeside have collected fossils farther north from strata in the bottom of Last Chance Canyon and Rocky Arroyo, on the eastern slope of the Guadalupe Mountains, where dark slabby limestones underlie partly sandy limestone of "Chupadera character," evidently the northern extension of the Capitan lagoonal beds. They state that the fossils indicate a level "somewhat lower" than the upper dark limestone of the Delaware Mountain, "but by no chance the lower dark limestone number" (our Bone Canyon). That the latter interpretation is, however, rather likely is suggested by the citation of *Productus* aff. *ivesi* and *Composita mexicana* which range no higher than the Leonard, and it is not improbable that the Capitan lies unconformably here on rocks older than the Delaware Mountain sandstone. Various isolated sections are described by Blanchard and Davis on the west side of the Guadalupe Mountains from two to nineteen miles north of the state line. At Tank Canyon they find "San Andres" (lower Capitan) resting directly on "Bone Springs" (Victorio Peak?), while to the south the Delaware Mountain formation intervenes and to the north "Yeso" underlies the "San Andres." These authors cite no fossils except fusulinids.

If the Guadalupian and Manzano are separate entities, then the Capitan limestones must pass out by overlap against the unconformably underlying Victorio Peak-San Andres limestones between Tank Canyon, and the Russel Gap locality where Darton found the southernmost Manzano fossils. Such an interpretation has, however, not been offered by any of the field geologists except Baker.*

The stratigraphic anomalies of the northwestern Guadalupe Mountains would be simply a local field problem were it not that this district is the link between two widespread and uniform stratigraphic sequences, the Guadalupian on the south and the Manzano on the north. By following the widely advanced field interpretations, Girty* concludes that "evidence is such as to demand a consideration, if not an adoption, of the hypothesis that the facies of the Guadalupian fauna is a regional matter denoting not time relations but geographic relations." If this is the case, the highly distinctive Guadalupian fauna is then simply an unusual local facies of the lower Permian, and is not the highly developed upper Permian fauna indicated to us by the long evolutionary sequence in the Glass Mountains.

From the above discussion it may be concluded that the Abo is probably of Wolfcamp and Wichita age, the Yeso approximately of Gym, Hess, and Clear Fork age, and the San Andres of Leonard and lower Double Mountain age. The correlation of the San Andres with the lower Double Mountain is checked by the subsurface tracing of the Blaine into the San Andres.[30]

*Permian of northern Arizona.*—In northern Arizona[31] the Redwall limestone is overlain by the Supai red sandstone and shale. In the Grand Canyon, the Supai is from 800 to 1100 feet thick, and rests on Redwall limestone of Mississippian age.

*Darton, N. H., **17**, p. 221.

*Cited by Girty, G. H., The Guadalupian Fauna and New Stratigraphic Evidence, Annals of New York Acad. Sci., Vol. 19, p. 141, 1909.

*Girty, *op. cit.*, p. 139.

*Baker, C. L., Contributions to the Stratigraphy of Southeastern New Mexico, Amer. Jour. Sci., (4), 49, pp. 99–126, 1920.

Also, Discussion of Permian Symposium, Amer. Assoc. Petrol. Geol., Vol. 13, pp. 1060–1062, 1929.

*Girty, *op. Cit.*, p. 141.

[30]Gould, C. N., and Willis, Robin: Tentative Correlation of the Permian Formations of the Southern Great Plains, Bull. Geol. Soc. Amer., vol. 38, pp. 435–438, 1927.

[31]Darton, N. H.: **16**, pp. 78–104. Also The Permian of Arizona and New Mexico, Bull. Am. Assoc. Pet. Geol., vol. 10, pp. 824–833, 1926.

Along the southern escarpment of the Arizona Plateau, especially on the scarp of the Mogollon Plateau and in the Holbrook Dome, there are 600 to 800 feet of red beds, which lie on Pennsylvanian limestone to the south and east but overlap upon pre-Cambrian granite to the northeast. The Pennsylvanian limestone was formerly included in the Redwall formation, but is now classified by the U. S. Geological Survey as the lower member of the Supai. This classification seems rather unnatural, for the remainder of the Supai is lower Permian in age, and is separated from the Pennsylvanian by a considerable time break. On the other hand the Supai red beds of the Mogollon Plateau and Holbrook Dome contain near the top rather thick limestones, which have yielded the following fossils, as listed by Girty:

| | |
|---|---|
| Echinocrinus cratis | Allerisma capax? |
| Rhombopora lepidodendroides | Deltopecten coreyanus |
| | D. occidentalis? |
| Productus cora | Myalina subquadrata? |
| P. leei | M. perattenuata? |
| Pustula nebraskensis | Monopteria marian |
| Pugnoides utah | Plagioglyptus canna? |
| Spirifer triplicatus | Pleurotomaria alamillana |
| Composita subtilita | Phymatifer cornudanus, |
| Hustedia mormoni | etc. |

According to Girty[32] this is a Manzano fauna, and a similar Permian one occurs in a limestone member of the Abo formation in the Zuni uplift of New Mexico. Because of the similarity faunally, lithologically, and stratigraphically of the Supai to the Abo, it can hardly be doubted that the Supai is the Arizona representative of the Abo. In the Defiance uplift of northeastern Arizona the Supai is present again, and there it contains *Walchia,* a plant diagnostic of the Permian. Darton states that "the evidence of these various fossils shows that the Supai formation as originally defined (i.e., excluding the limestone formerly placed in the Redwall) is all of Permian age."[33]

Above the Supai is the Hermit shale, varying from 80 to over 500 feet in thickness. It is exposed along the upper slopes of the Grand Canyon, and does not occur on the south side of the Arizona Plateau. It is unconformable upon the Supai. Permian plant remains occur in it, including *Callipteris conferta, Walchia* cf. *W. gracilia, Gigantopteris* cf. *G. sphenophyllum, Alethopteris?,* and *Pecopteris* (**148**, pp. 353–360).

The succeeding unit, the Coconino sandstone, is very widespread over the Arizona Plateau. It is generally from 300 to 600 feet thick, and consists of massive light-grey to pale buff sandstone of remarkably uniform character. The only fossils that have been found in it are footprints. In places in the Kaibab Plateau the Coconino is not distinguishable, for in that region the entire Coconino is replaced at some localities by limestone of the Kaibab, which overlaps the Coconino by replacement.

The highest member of the Arizona Permian section is the Kaibab limestone, which covers a very large area in northern Arizona. To the north of the Grand Canyon the Kaibab is 1000 or more feet in thickness, and here the Coconino is not present. To the east and southeast the Coconino becomes distinguishable, and thickens, replacing the lower Kaibab so that in eastern sections the Kaibab may be but a few feet thick. The Kaibab varies in thickness also as a result of pre-Triassic erosion. Approximately the upper 150 feet of the Kaibab is distinguished as the Harrisburg gypsiferous member, the upper part of which is known as the "Bellerophon limestone."

The brachiopods occurring in the main part of the Kaibab are, as identified by Girty, with some modifications by myself:

| | |
|---|---|
| Orbiculoidea convexa | Avonia subhorrida rugatula |
| Derbya aff. D. nasuta | Pugnoides osagensis var. |
| Meekella pyramidalis | Squamularia guadalupensis |
| Chonetes quadratus | Spiriferina campestris? |
| Productus ivesi | Composita mexicana? |
| P. occidentalis | C. subtilita |
| Waagenoconcha montpelierensis | Dielasma sp. |
| | Heterelasma n. sp. |

Of the forms definitely identified, *Orbiculoidea convexa* and *Meekella pyramidalis* do not occur in Trans-Pecos Texas. *Chonetes quadratus* occurs in the upper Word, *Productus ivesi* is most abundant in the Leonard, and the others occur in several formations, all of them being present in the Leonard and some as low as the Wolfcamp and others as high as the Capitan. This evidence would seem to be conflicting, but it is a striking fact that the most abundant of all the Kaibab species, *Productus ivesi,* is also abundant in the Leonard and does not occur in Texas at a higher horizon. The other species, though mostly longer ranged, also occur in the Leonard, so that the Kaibab fauna as a

[32]In Darton and Reeside, Bull. Am. Assoc. Pet. Geol., vol. 10, pp. 825–826, 1926.

[33]Op. cit., p. 89.

whole has much in common with that of the Leonard. Unfortunately, no ammonoids occur in the Kaibab, but the formation correlates definitely both faunally and lithologically with the San Andres of New Mexico, which in turn also has been correlated by us with the Leonard.

The fauna of the Bellerophon limestone is quite unlike that of the main part of the Kaibab, consisting almost wholly of molluscs, of which the characteristic species is *Bellerophon majusculus.*

With the age of the Kaibab definitely fixed as that of the Leonard and San Andres, the age of the beds below it may be approximately placed. The Coconino may represent the same phase of deposition as the Glorieta sandstone of New Mexico and the Hermit part of the Yeso.*

### PERMIAN OF THE GREAT BASIN AND NORTHERN ROCKIES

*Permian of southeastern and central Utah and southwestern Colorado.*—In southeastern Utah[34] the lowest formation in the section is the Goodridge, consisting of dense, medium to massively bedded limestone, red and greyish-white sandstone, and red and dark-grey shale, with a total thickness of 1500 feet. In the lower 1200 feet there is a typical Pennsylvanian fauna with almost no molluscs; the upper 300 feet bears almost entirely a molluscan fauna, which is of lower Permian type and closely similar to that of the Rico formation of southwestern Colorado, the Supai of the south side of the Arizona Plateau, and the Abo. The Goodridge grades upward into Supai red beds, which are here about 400 feet thick. The upper Goodridge probably equals the lower Supai of the Grand Canyon region, while the Supai of southeastern Utah is probably of upper Supai and in part perhaps Hermit age. The overlying Coconino sandstone here attains a thickness of 1000 feet. The Kaibab is not over 150 feet thick, and it thins progressively northward, with a corresponding thickening of the Coconino sandstone. The fauna here is of the normal Kaibab type rather than that of the Bellerophon limestone.

In southwestern Colorado[35] and the Moab region of east-central Utah the Pennsylvanian Hermosa limestone is overlain by the Rico formation, which consists of sandstone and conglomerate with intercalated shale and sandy fossiliferous limestone, 250 to 600 feet in thickness. The fauna is almost wholly molluscan, and contains types similar to those of the Kansas Permian. The Rico grades upward without break into the unfossiliferous Cutler red shale, sandstone, grit, and conglomerate, having a total thickness of about 2000 feet. In places the upper Rico is laterally replaced by the Cutler. Girty believes that the fauna of the Rico is of Permian type. It correlates with the upper Goodridge, while the Hermosa correlates with the lower Goodridge. In east-central Utah the Rico has been traced westward by measured sections into the Supai (the upper Goodridge not being present there), and the Cutler has been traced into the Coconino.[36] None of the Rico or upper Goodridge fossils is of use in making correlations with the Trans-Pecos Texas section of Permian.

In the San Rafael Swell of central Utah[37] there are 700 feet of Coconino overlain by 85 feet of Kaibab. There are two faunas here, one of which is that of the Bellerophon limestone, and the other a fauna having Phosphoria affinities, shown by the presence of *Waagenoconcha montpelierensis* and *Pustula nevadensis. Productus ivesi* and *P. occidentalis* are absent, and there is a large number of molluscs. This perhaps indicates a transition to the Phosphoria from the typical Kaibab facies. The presence of the Bellerophon limestone fauna in only 85 feet of the formation indicates that only the upper Kaibab is present here, the lower probably having been replaced by the Coconino.

*The Phosphoria formation and its equivalents.*—Throughout northern Utah, western Wyoming, southeastern Idaho, and western Montana the Permian is represented by the very widespread and remarkably uniform Phosphoria formation. In

*Since this paper was written, new observations have been published by A. A. Baker and J. B. Reeside on the Permian of northeastern Arizona and its surroundings, in which special emphasis is placed on interfingering and lateral replacements of different facies. These in general confirm our own briefer conclusions (Baker and Reeside, Correlation of the Permian of Southern Utah, Northern Arizona, Northwestern New Mexico, and Southwestern Colorado, Bull. Amer. Assoc. Petrol. Geol., Vol. 13, pp. 1413–1448, 1929).

[34]Longwell, C. R., Miser, H. D., Moore, R. C., Bryan, Kirk, and Paige, Sidney: Rock Formations in the Colorado Plateau of Southeastern Utah and Northern Arizona, U. S. Geol. Sur. Prof. Paper 132A, 1923.

[35]Cross, W., and Hale, Allen D.: Engineer Mountain Folio, Colorado, U. S. Geol. Sur. Folio 171, pp. 6–7, 1910. Also Baker, A. A., Dobbin, C. E., McKnight, E. T., and Reeside, J. B.: Notes on the Stratigraphy of the Moab Region, Utah, Bull. Am. Assoc. Pet. Geol., vol. 11, pp. 785–796, 1927.

[36]Prommel, H. W. C., and Crum, H. E.: Structural History of Parts of Southeastern Utah from Interpretation of Geologic Sections, Bull. Am. Assoc. Pet. Geol., vol. 11, pp. 809–815, 1927.

[37]Gilluly, James: Geology and Oil and Gas Prospects of Part of the San Rafael Swell, Utah, U. S. Geol. Sur. Bull. 806, pp. 80–83, 1928.

northernmost Utah and southeastern Idaho the Phosphoria rests on the Wells formation of Pennsylvanian age, in western Wyoming on the Tensleep sandstone, and in western Montana on the Quadrant.

The Phosphoria formation consists of from 80 to 450 feet of dark-grey cherty quartzite containing layers of nodular chert and shale, and one or more beds of phosphate rock. In different localities it is divisible into various members. In southeastern Idaho Mansfield **(108,** pp. 75–81) has divided it into the phosphatic shale member below and the Rex chert member above, the former 75–180 feet thick, and the latter 110–550 feet thick. The fauna of the phosphatic shale member has been described by Girty **(70)**, and that of the limestone and chert members of the formation by Meek and Girty in various places.[38]

The fauna of the phosphatic shale is considerably different from that of the remainder of the formation, but together the faunas of the two members form a rather varied aggregation of forms of similar stratigraphic affinities. In the limestone and chert division of the formation the following are characteristic brachiopods:

Productus multistriatus
Linoproductus waagenianus (=Productus eucharis)
Pustula nevadaensis
Avonia subhorrida
Aulosteges hispidus
Camarophoria sp.
Spirifer pseudocameratus
Spiriferina pulchra
Composita mira
C. persinuata

*Spiriferina pulchra* is such a common form that this fauna has come to be known as the *Spiriferina pulchra* fauna, and it is much more widespread than that of the phosphatic shale, for the biota occurs also throughout central and eastern Nevada (where most of the type specimens of its species were collected by the King survey) and as far south as the Inyo Mountains of southeastern California, where the Owenyo limestone[39] 125 feet thick, carries an abundant fauna of this type.

The characteristic species of the phosphatic shale fauna are:

Lingula carbonaria?
Orbiculoidea missouriensis
Chonetes ostiolatus and vars.
Productus geniculatus
Linoproductus waagenianus (=Productus eucharis)
Waagenoconcha montpelierensis
Linoproductus phosphaticus
Liorhynchus weeksi
Pugnoides swallovianus (Pugnax osagensis var. occidentalis)
Ambocoelia guadalupensis (Ambocoelia arcuata)
Leda obesa
Plagioglypta canna
Omphalotrochus ferrieri
O. conoideus
Gastrioceras simulator
Hollina emaciata occidentalis

In the Uinta Mountains the phosphate-bearing Permian was originally known as the "Bellerophon limestone," after one of its characteristic fossils; the presence of *Bellerophon* may indicate faunal relationship to the "Bellerophon limestone" at the top of the Kaibab, which, as shown below, is probably of the same age. The Permian of the Uinta Mountains is underlain by the Weber quartzite, which may be the equivalent of the Coconino.

Considering the species themselves that occur in the Phosphoria fauna, there is little to indicate their age. Many of them are related to species of the *Schwagerina* horizon and the Artinsk of the Russo-Arctic region. Such are *Chonetes ostiolatus, Productus multistriatus, P. geniculatus, Linoproductus waagenianus, Waagenoconcha montpelierensis, Linoproductus phosphaticus,* and species of *Omphalotrochus* and *Gastrioceras.* Other forms are closely related to Pennsylvanian species, and apparently have descended from them; these are *Pustula nevadaensis, Spirifer pseudocameratus, Ambocoelia guadalupensis, Composita mira, C. persinuata,* and *Pugnoides swallovianus. Aulosteges hispidus* and *Avonia subhorrida* are Permian types, but so far as their structure would indicate, they might occur in any part of the Permian system. As a result of the nondescript nature of the Phosphoria fauna, there has been much controversy as to its age. It was long regarded as Pennsylvanian, but in 1920, in the "Ore Deposits of Utah," Girty classed it as Permian, because of its affinity to the lower Permian of Russia, though admitting that the evidence was inconclusive. In 1927, Girty[40] discussed the age of the Phosphoria at some length. It was pointed out that the Phosphoria fauna was very similar faunally to the lower Permian of Alaska.

[38]Girty, G. H.: **73**, pp. 644–647. Also, Meek, F. B.: **118**.
[39]Kirk, Edwin: U. S. Geol. Sur. Prof. Paper 110, p. 43, 1918.
[40]In U. S. Geol. Sur. Prof. Paper 152, pp. 79–81, 1927.

The stratigraphic evidence of the age of the Phosphoria, the fact that the formation appears to trace southward into the upper part of the Kaibab in the San Rafael Swell, is in opposition to the correlation of the Phosphoria with the lower Permian, for the Kaibab is very certainly of middle Permian age. The stratigraphic evidence is conclusively supported by the correlation of the Phosphoria fauna in the Trans-Pecos Permian section. Most of the Phosphoria species occur in the Texas fauna; they are *Orbiculoidea missouriensis, Chonetes ostiolatus, Productus multistriatus, Linoproductus waagenianus, L. phosphaticus, Avonia subhorrida, Waagenoconcha montpelierensis, Leiorhynchus weeksi, Pugnoides swallovianus, Spirifer pseudocameratus, Ambocoelia guadalupensis, Composita mira,* and *C. persinuata.* Only *Pustula nevadaensis, Productus geniculatus, Aulosteges hispidus,* and *Spiriferina pulchra* have not been found in Texas. Of the Phosphoria species occurring in the Texas section, 7 are restricted to the Word, 0 to the Leonard, 4 occur in the Leonard and Word, and 2 in the Hess, Leonard, and Word. Furthermore, *Gastrioceras simulator* of the Phosphoria is very closely similar to *G. roadense* of the Word. The evidence of numbers, therefore, favors a Word age for the Phosphoria.

It is obvious that in two faunas containing species in common with one another, some species would have migrated from the region of the first fauna into that of the second while others would have migrated from the second into the first. In this instance it seems reasonable to suppose that the species occurring in both the Leonard and Word originated in Texas and in Word time migrated northward into Idaho. *Composita mira, Spirifer pseudocameratus,* and *Ambocoelia guadalupensis* almost certainly originated in Texas, for their immediate ancestors occur in the lower part of the Texas section.

If the Phosphoria is of Word age, the question arises as to how the Phosphoria and the Word succeeded in interchanging their invertebrate species when the rocks intervening between the Phosphoria and the Word-Delaware Mountain do not contain these species. The only Phosphoria brachiopod species occurring in the Kaibab is *Waagenoconcha montpelierensis,* while the common Kaibab species are entirely lacking in the Phosphoria. In the San Rafael Swell, however, the upper or "Bellerophon" limestone of the Kaibab contains *Waagenoconcha montpelierensis* and *Pustula nevadaensis,* and does not have the Kaibab species. This fact suggests that the Phosphoria fauna was contained in an uppermost member of the Kaibab, probably above the *Bellerophon* fauna, and has now been eroded away in most places. The "Bellerophon" limestone itself was generally eroded away before the Triassic, so that it is quite possible that still higher beds, no longer preserved, were once present. The same circumstance might have occurred in southern Arizona, where even the Kaibab equivalents are seldom seen between the Naco limestone and the Mesozoic, because of the large amount of erosion following the retreat of the Permian sea from the region. It therefore seems probable that a marine connection, traces of which are lost, existed between the Phosphoria of the northwestern United States and the Word of Texas.

The reason for the lower Permian aspect of the Phosphoria fauna remains to be explained. A large proportion of the Phosphoria species have close affinity to Russo-Arctic lower Permian species, and some of the resemblances may prove to be identities. Girty has shown[41] that the Phosphoria fauna is very similar to the Alaskan "Artinsk" fauna. There can hardly be any doubt that there was direct marine connection between the Phosphoria and the Alaskan Permian seas. It seems likely, therefore, that the species of Russo-Arctic affinity migrated into the Phosphoria and thence to Texas by way of the Alaskan Permian seas. Whether the Alaskan sea derived its fauna from the northeast alone, by way of Spitzbergen, north Greenland, and Ellesmere and Heiberg Islands, or from both the northeast and the west, connecting also with the lower Permian of Ussuriland, is a matter for conjecture; at any rate, both the Permian of Ussuriland and that of Ellesmere and Heiberg Islands is undoubtedly lower Permian, of the age of the *Schwagerina* horizon and lower Artinsk. Therefore, the lower Permian faunas must have entered the Alaskan sea from the north, and remained there with little change until middle Permian time, when marine connection opened up toward the south. The marine connections with the Russian seaways having in the meantime been cut off, middle Permian forms did not gain access to Alaska, and when the sea spread southward into the Phosphoria region the accompanying fauna

[41] U. S. Geol. Sur. Prof. Paper 152, pp. 79-81, 1927.

was merely a slightly modified lower Permian fauna lacking middle Permian types.*

*Permian of the Canadian Rockies.*—There is little record of Permian rocks in the Canadian Cordillera, but they are known to exist in several places. In the section near Banff, Alberta, the Rocky Mountain quartzite,[42] about 600 feet thick, consists of an alternation of light grey quartzite and light grey limestone; in the upper part there is a phosphate bed associated with chert, quartzite, and shale. It contains an abundance of *Euphemus carbonarius arenarius* and *Plagioglypta canna,* besides *Orbiculoidea arenaria, Productus semireticulatus, Paraphorhynchus obscurum, Bakewellia parva, Myalina wyomingensis,* and *Deltopecten occidentalis latiformis. Plagioglypta canna* occurs also in the Phosphoria, while *Euphemus carbonarius arenarius* is stated by Shimer to be similar to *E. subpapillosus* of the Phosphoria, and the two may be identical. *Paraphorhynchus obscurum* would unhestitatingly be referred to *Leiorhynchus weeksi,* were it not for the statement by Shimer that it is radially lirate. One is tempted to think that this feature may be the result of some peculiarity of preservation. At any rate, there can be little doubt that the Rocky Mountain quartzite is a northern equivalent of the Phosphoria.

Kindle has described[43] from French Mountain in northern British Columbia an occurrence of *Lyttonia* associated with other Permian types, all imperfectly preserved. Strangely enough, this fauna has no resemblance to either the Phosphoria or the Alaskan faunas.

### PERMIAN OF NORTHERN CALIFORNIA

In northern California[44] the upper Mississippian Baird shale is overlain by the McCloud grey limestone, containing *Schwagerina robusta, Fusulina gracilis, F. californica, Clisophyllum gabbi, Omphalotrochus whitneyi* (**6**, pp. 55–56), and *Hustedia compressa.* It is followed by the Nosoni sandstone and tuff, in Shasta and Plumas counties, with *Fusulina elongata.* Smith has listed *Productus* cf. *spiralis* from this horizon; it is quite possible that this is actually *P. invesi.* His *Camarophoria* cf. *purdoni* may be *C. venusta,* while *Productus* cf. *abichi* may be *Waagenoconcha montpelierensis.* The highest Paleozoic formation is the Wildwood limestone of Trinity County, with a Guadalupian fauna, including *Stacheoceras.* It is unfortunate that we know nothing further concerning this interesting section, as it might throw considerable light on the origin of some of our Permian species.

### PERMIAN OF ARCTIC AMERICA

*Permian of Alaska.*—The Permian faunas of Alaska are little known as yet, and the best known ones are in Kuiu Island in southeastern Alaska, and on the west bank of the Yukon opposite the mouth of Nation River. The rock at the first locality is a white cherty limestone, unconformably overlain by Triassic. The fauna is, as has been stated above, similar to that of the Phosphoria, and closely related to that of the *Schwagerina* horizon and the Artinsk of the Russo-Arctic region.

*Permian of the Parry Islands.*—In Banks Island, Victoria Island, and Melville Island there is a considerable thickness of Lower Carboniferous non-marine plant bearing beds. This is succeeded in the northern parts of the Parry Islands by limestone containing fossils which were described by Haughton.[45] Haughton's descriptions did not, however, appear to be adequate, and later workers[46] have shown that his species have different relations from what Haughton supposed. His *Productus sulcatus* var. *borealis* has been shown to be identical with *Horridonia timanica* (Stuckenberg), one of the most common fossils in the Permian faunas of Ellesmere Island and northeast Greenland. The name *P. borealis* considerably antedates *P. timanicus,* and perhaps should be the proper name for that species. So little is known of the faunas of this region that all that may be said definitely is that there is here a connecting link between the "Artinsk" of Alaska and the Permian of Ellesmere and Heiberg Islands, which, it will

*After this manuscript was written a paper appeared in which the Phosphoria fauna is described (Branson, C. C., Paleontology and Stratigraphy of the Phosphoria Formation, Univ. Mo. Studies, Vol. V, 1930). In the light of the facts presented above, Branson's conclusion that the formation lies on the boundary between the Pennsylvanian and Permian must be considered as unlikely.

[42]Shimer, H. W.: Upper Palaeozoic Faunas of the Lake Minnewanka Section, near Banff, Alberta, Canadian Geological Sur. Bull. 42, pp. 1–85, 1926.

[43]Trans. Royal Soc. Canada, ser. 3, vol. 20, pp. 109–111, 1926.

[44]See Smith, J. P.: Geological Record of California, Jour. Geol., vol. 18, p. 217, 1910. Also Mesozoic Changes in the Faunal Geography of California, Jour. Geol., vol. 3, pp. 372–374, 1895.

[45]Reminiscences of Arctic Ice-Travel in Search of Sir John Franklin and His Companions. By Captain F. L. McClintock. With Geological Notes and Illustrations by Samuel Haughton. Jour. Royal Dublin Soc., vol. 1, pp. 183–250, 1856–1857. Not seen.

[46]See Grönwall, K.: The Marine Carboniferous of North-East Greenland and Its Brachiopod Fauna, Medd. om Grønland, vol. 43, pp. 599–604, 1917.

be shown, are connected by way of northeast Greenland to the lower Permian of Spitzbergen.

*Permian of Ellesmere and Heiberg Islands.*—The lower Permian fauna of Ellesmere and adjacent islands has been made known as a result of the work of Whitfield and of Tschernyschew and Stepanow.[47] It has been collected from a white cherty limestone at Store Bjornekap, southern Ellesmere Island, from a dark grey limestone at Svarte Veg, at the north end of Heiberg Island, and from limestone in northeastern Ellesmere Island at Fielden Peninsula and Cape Sheridan. Tschernyschew and Stepanow have described 61 species of fossils from the first two localities, of which 34 are of brachiopods. Practically all of these species are identical with forms in the Russian *Schwagerina* horizon and Artinsk, and most of them occur in the lower Permian of Spitzbergen. Many are close to Phosphoria and Word-Delaware Mountain species, as the following table shows:

| Species in Ellesmere and Heiberg Islands | Related Phosphoria or Word forms |
|---|---|
| Productus arcticus (=P. inflatus T. & S.) | Productus arcticus |
| Avonia pseudoaculeata | Avonia subhorrida |
| Waagenoconcha irginae | Waagenoconcha montpelierensis |
| Linoproductus aagardi | Linoproductus waagenianus |
| L. simensis | L. nasutus |
| L. cancriniformis | L. phosphaticus |
| Productus weyprechti | Productus multistriatus |
| Horridonia timanica | Horridonia texanus |
| Derbya grandis | Derbya nasuta |
| Squamularia guadalupensis | Squamularia guadalupensis |
| Spirifer lyra | Spirifer sulcifer |
| Rhynchopora nikitini | Rhynchopora taylori |
| Spiriferina cristata | Spiriferina laxa |

Other species have relatives only at lower horizons than the Word, as *Buxtonia porrecta (=B. peruviana)*, *Spirifer fasciger* (related to *S. costella)*, and *S. marcoui* (related to *S. marcoui infraplica)*. The *Spirifer condor* figured by Tschernyschew and Stepanow is very obviously incorrectly identified, for it has a very high cardinal area. The nature of this fauna strongly suggests that it lived in the seaway connecting the Russian Permian basin, by way of the Spitzbergen Permian sea, with the Alaskan embayment, for most of the Alaskan Permian forms as identified in a preliminary way by Girty correspond to species from Ellesmere and Heiberg Islands. Moreover, the presence of a large number of similar and some identical species in the Word indicates that those species must have migrated into the Alaskan basin and lived there until middle Permian time, when the Arctic faunas invaded Trans-Pecos Texas by way of the Phosphoria seaway. Some of the species such as *Productus arcticus* and *Horridonia texana*, occur in the Word and in the Arctic fauna but are not known in the Phosphoria. Perhaps further collecting will eventually reveal these forms in the Phosphoria.

*Permian of northeast Greenland.*—Grönwall has described in detail the "Upper Carboniferous" faunas of northeast Greenland and their relations.[48] In Holm's and Amdrup's Lands, between lat. 80° and 81°, there are at the base of the section 200–500 meters of sandstone and shale with Lower Carboniferous plant fossils. This formation is obviously equivalent to the other widespread Lower Carboniferous terrestrial formations of the Arctic, such as that above-mentioned which occurs in Banks Island and the southern parts of the Parry Islands, and the Culm sandstone of Spitzbergen. Overlying this unit unconformably are 200–300 meters of limestone interstratified with sandstone and shale. The fossils in this formation are quite unlike those in the overlying limestone. Most of them are not particularly diagnostic, but the presence of *Spirifer supramosquensis* indicates that the fauna is earlier than that of the *Schwagerina* horizon of Russia, and the absence of *Horridonia timanica* and *Productus arcticus (=P. inflatus* Grönwall, non McChesney), which occur in the overlying limestone, also indicates a pre-Permian age. On the other hand, the Permian species *Chonetella nasuta* is figured from this horizon, and its occurrence, in view of the other

[47]Whitfield, R. P.: **187**, pp. 51–58.

[48]Grönwall, K.: The Marine Carboniferous of North-East Greenland and Its Brachiopod Fauna, Medd. om Grønland, vol. 43, pp. 511–618, 1917.

evidence, seems to be anomolous. Yet it seems most reasonable to suppose that this unit is equivalent to the *Cyathopyllum* limestone of Spitzbergen, a conclusion which is borne out by the lithologic similarity of the two. The upper member of the Carboniferous of northeast Greenland consists of 500 to 1000 meters of fine-grained light limestones and white detrital limestone with "pebbles" of fragments of organic origin, and dolomites. Fossils are common. *Schwagerina princeps* is abundant in places. Most of the other fossils are not particularly diagnostic. *Horridonia* and *Productus arcticus* are, as in the Permian faunas farther west in the Arctic archipelago, common and characteristic.

It seems logical to conclude that the *Schwagerina* limestone of this section is equivalent to the lower Permian limestones of the Parry Islands and Ellesmere and Heiberg Islands. The *Spirifer supramosquensis* horizon, however, is probably of Upper Carboniferous age and is unrepresented farther to the west. The connection with the Carboniferous and Permian sections of Spitzbergen and Russia is even closer, as is shown by the following table, taken from Wiman (**188**, pp. 9–22):

| Series | Ne. Greenland | Spitzbergen | | Bear Island | Russia |
|---|---|---|---|---|---|
| Artinsk<br>*Schwagerina* horizon | Upper limestone | *Productus*-bearing siliceous limestone | | Absent | Artinsk |
| | | *Spirifer* limestone | | *Spirifer* limestone | *Schwagerina* horizon |
| | Break? | Break | | Decided unconformity | Break? |
| Uralian | Lower limestone | *Cyathophyllum* limestone | *Fusulina* ls.<br>Cora limestone with *Fusulina* | Cora limestone<br>Coral sandstone<br>Break<br>*Fusulina* ls. | Cora horizon<br>*Omphalotrochus* hor. |
| Moscovian | Absent? | | Limestone with *Spirifer mosquensis*<br>Basal conglomerate | Unfossiliferous sandstone<br>Sandstone with *Composita ambigua* | *Mosquensis* limestone |
| | Break | Break? | | Break | Break |
| Lower Carboniferous | Sandstone and shale | Culm sandstone | | Absent | Lower Carboniferous |

Following Beede and Kniker and Schuchert,[49] the *Schwagerina* horizon is regarded as Permian, the greatest faunal change in the Russian section taking place between the *Cora* and *Schwagerina* horizons, rather than between the *Schwagerina* horizon and the Artinsk. *Schwagerina* is regarded as an index to the lowest Permian. The notable unconformity between the *Cora* limestone and the *Spirifer* limestone (*Schwagerina* horizon) in Bear Island seems to be physical evidence for placing the *Schwagerina* horizon in the Permian.

From the facts above brought out, it may be seen that a sheet of limestone was deposited in lower Permian time in a seaway extending from the Urals by way of the Timan ranges, Novaya Zemlya, Bear Island, Spitzbergen, northeast Greenland, Ellesmere and Heiberg Islands, and the Parry Islands to Alaska. In this seaway lived a rich invertebrate fauna, with very little difference in composition from Russia to Alaska. The sea may have withdrawn from most of this Arctic region at the close of lower Permian time, but in Alaska it lingered until the middle Permian, when an Arctic invasion of western North America (the Phosphoria sea) carried the Russo-Arctic faunas as far south as Trans-Pecos Texas.

### "UPPER CARBONIFEROUS" OF THE ANDES

The fauna of the "Upper Carboniferous" limestone of Bolivia and Peru was first made known by d'Orbigny in 1842,[50] but remained imperfectly known until the publications of Kozlowski[51] and Meyer (**120**, pp. 590–652) appeared in 1914. In 1914 Douglas (**34**, pp. 30–37) outlined the stratigraphy and palaeontology of the "Carboniferous" of the Lake Titicaca region, and in 1929 Steinmann (**158**, pp. 34–53) summarized our knowledge of the "Upper Carboniferous" stratigraphy and

[49] **6**, pp. 60–71. Also **150**, pp. 806–807.

[50] Voyage dans l'Amérique méridionale.

[51] Kozlowski, Roman: Brachiopodes du Carbonifére supérieur de Bolivie, Ann. de Pal., t. 9, 1914.

faunas of Peru. The faunas of the "Carboniferous" of the Lake Titicaca region appear to come from a limited thickness of siliceous limestone, probably not representing a very great lapse of geologic time.

Kozlowski described 33 species of brachiopods from this horizon, of which 15 occur in the Pennsylvanian of North America and 21 in the Permian of Trans-Pecos Texas. Another species, *Enteletes andii*, is very close to, if not identical with the Texas species, *E. dumblei*. Of the species occurring in the Texas Permian, 9 do not occur in the Pennsylvanian, namely, *Productus inca*, *Linoproductus* cf. *cancriniformis*, *Buxtonia peruviana*, *Waagenoconcha irginae (=W. montpelierensis)*, *Avonia boulei*, *Derbya buchi*, *Orthotichia kozlowskii (=O. morgani* Kozlowski non Derby), *Spirifer condor*, and *Camarophoria thevenini*. Two other species, *Marginifera capaci* and *Linoproductus villiersi*, do not certainly occur in the Pennsylvanian. Only 3 of the Bolivian species occur in the Pennsylvanian of the United States but not in the Permian of Texas; these are *Chonetes geinitzianus (=C. glaber* Kozlowski non Hall), *Spiriferina campestris*, and *Pugnoides utah*. None of these is particularly diagnostic of any horizon, and any might occur in the lower Permian. On the other hand, several of the forms restricted to the Permian are generally diagnostic of the Permian, such as the genus *Waagenoconcha* and *Buxtonia peruviana*. Therefore, it is very probable that the Andean "Upper Carboniferous" fauna is of Permian age. We will now consider to what part of the Permian section the Andean faunas belong. Some of the species occur in the *Uddenites* zone and the middle and upper Wolfcamp formation, and many of the same species and some others of the Andean fauna are present in the Gym formation. *Productus inca* is one of the most characteristic species of the middle and upper Gym, while other forms are equally characteristic of the Wolfcamp. The explanation of this may be that the Andean fauna is of both Wolfcamp and Gym (=Hess) age, or that it represents a relatively short time, such as that of the Wolfcamp, and that certain species, such as *Productus inca*, originated in South America and did not reach Texas until later than the remainder of the fauna, which was indigenous to Texas, or originated in neither Texas nor the Andean region.

The Amazonian Carboniferous fauna described by Derby has commonly been regarded as of the same age as the Andean fauna, and some specific identities have been pointed out. The differences are, however, far more striking than the similarities, and it is a notable fact that the Amazonian fauna contains lower Pennsylvanian species, such as *Spirifer rockymontanus*, and lacks such Permian types as *Waagenoconcha* and *Buxtonia peruviana*. For two faunas in the equatorial region of the same continent these are remarkably dissimilar. The true relations of the faunas are probably demonstrated in Trans-Pecos Texas and southeastern Arizona. Near Winkelman, Arizona, in the Galiuro Mountains, there is a fauna in the Naco limestone with very close affinity to that of Brazil (**161**, pp. 324–327). *Orthotichia morganiana* and *Derbya tapajotensis* occur there in association with such lower Pennsylvanian fossils as *Productus morrowensis* and *Spirifer rockymontanus*. The lower Pennsylvanian age of this horizon can hardly be questioned. In the Gym formation, the lower part of the Permian, overlying the beds containing this fauna, there occurs the fauna of the Andean "Upper Carboniferous." It is probable, therefore, that there are two different ages of late Paleozoic faunas in northern South America.

## PALEONTOLOGY

### SUMMARY OF THE PERMIAN FAUNAS

*General character.*—The Permian section of Trans-Pecos Texas is the most complete of that age in North America, and for a long, unbroken sequence it is perhaps not exceeded in any other Permian sequence in the world. A study of its faunas not only reveals, in a way which has not been possible before, the evolution of the Pennsylvanian faunas into the highly specialized Permian ones, but also makes possible the accurate chronological placing of many late Paleozoic faunas in other widely separated regions, the age relations of which until now have been uncertain. It is therefore one of the most important standard sections of Permian rocks in the world, and is comparable to the sections in the Salt Range of India, the island of Timor, and the Urals.

The Permian faunas of Trans-Pecos Texas are a remarkable contrast to those of the mid-continent region and of New Mexico, Arizona, and northwestern North America, which, though abundant in individuals, are poor in numbers of species. In Trans-Pecos Texas almost all the forms of invertebrate life of the time are found, a fact which

indicates the presence there in Permian time of a sea in which conditions were very favorable to invertebrate life, having such free communication with the other Permian epicontinental seas that there was considerable intermigration of faunas throughout most of Permian time. An analysis of the Texas fauna shows a derivation from many different sources, so that there was a mingling of types from the Andean region, the Arctic, and the Tethyan seaways with those which seem to have been indigenous and to have descended from American Pennsylvanian species.

In the present report there are described 185 species and varieties of brachiopods in 47 genera; of these, 62 species, 12 varieties, and 4 genera are new.* Of the species and varieties described by myself or by Girty in "The Guadalupian Fauna," 22 were described originally from the Pennsylvanian, 17 by Shumard from the Guadalupe Mountains, 67 by Girty in "The Guadalupian Fauna," 3 by Girty from the Permian of New Mexico, 2 by Newberry from the Kaibab limestone, 7 by Girty from the Phosphoria, 4 by Meek from the Phosphoria, 9 by d'Orbigny and Kozlowski from the Andean "Upper Carboniferous," and the remaining 14 by various authors from scattered sources. The total of 219 species and varieties does not recognize the questionable forms noted by Girty and some of the species named by Girty which here are regarded as synonyms.

The Trans-Pecos Permian faunas differ markedly from those of the Pennsylvanian in their greater number of species and genera. In the large collections studied by Plummer and Moore from the Pennsylvanian of central Texas, for example, 95 species and varieties of brachiopods, in 31 genera, were recognized, while in the present paper 185 species and varieties in 46 genera are described from the Permian. Contrary to the general belief that Permian life was greatly restricted, this fauna compares favorably with that of other Paleozoic periods. In his "Mississippian Brachiopoda," Weller describes 308 species of Mississippian brachiopods in 63 genera.[52] Although this number considerably exceeds that of the present fauna, it should be remembered that the forms handled by Weller were collected throughout the entire northern Mississippi valley over a period of more than 40 years, while the Permian faunas now described represent the collections made in a much smaller area during a few field seasons. There can be no doubt that further collecting will bring forth a large number of new species, and probably genera, heretofore unrecognized.

The species here described are mostly very short ranged. The number restricted to the Permian is 158. Of these, 74, or 40%, are limited to one formation, 72, or 39%, to two, and 28, or 15%, to three, while only 8, or 4%, range through four formations and 3, or 1.5%, through five.[53]

*Biological summary of the brachiopods.*—The Atremata are, so far as known, unrepresented in this fauna. The Neotremata are not much better known, for only two genera were found, each represented by a single specimen.

Of the Orthacea only the groups typical of the late Paleozoic are present. There are 5 species of *Rhipidomella*, a larger number than has been recognized in other Permian faunas. Except for *R. hessensis*, which is closely related to the Asiatic species *R. subquadrata*, the species of this genus seem to be of American origin. The genus *Orthotichia*, derived from *Schizophoria*, is characteristic of the Pennsylvanian and lower Permian, and is present in the lower part of the Glass Mountains section. *Enteletes* is represented by 5 species. One of these, *E. dumblei*, is very close to the Andean "Upper Carboniferous" species *E. andii*, while *E. plummeri* finds its closest affinity with species of the Tethyan region. *Parenteletes* is an upper Pennsylvanian and lower Permian genus of rather wide distribution in rocks of the Mediterranean faunal facies. The orthids are much more diversified in

*After the present manuscript had been completed and submitted for publication, and after the writer had departed for work in South America, a short paper was published by Girty, describing seven species of brachiopods from the Diablo Plateau. Professor Dunbar has made a preliminary comparison and suggested the following probable relationships:

*Chonetes victorianus* Girty = *Chonetes spinoliratus* King
*Chonetes consanguineus* Girty = *Chonetes verneuilianus* King (non-Norwood and Pratten)
*Chonetes quadratus* Girty = [?]*Chonetes quadratus* King
*Schizophoria huecoensis* G. = [?]*Orthotichia kozlowski* King
*Camerophoria inaequalis* G. = *Camerophoria venusta* King (in part)
*C. huecoensis* Girty = *C. deloi* King (in part)
*Pugnoides mesacostalus* Girty = *Pugnoides transversus* King

In view of the author's absence, it has seemed undesirable to alter these names in his finished manuscript.—Editor's note.

[52]Probably at least 10 additional genera would be recognized in the productids, etc., if the standard of the present report were followed.

[53]In computing these figures, the species occurring in the Gym were disregarded except where a species was restricted to it, because of the uncertainty as to its correlation. Species occurring also in the Pennsylvanian were regarded as occurring in one additional formation. If the species was recognized also by Girty, the information given by him regarding its range was incorporated with my own data.

the present fauna than in the American Pennsylvanian and Mississippian. The same genera are present in the former, but in the Permian they contain many more forms, while the Mississippian contains a good many species but only two genera.

Most of the late Paleozoic genera of the Orthothetinae are present. Six species of *Streptorhynchus* have been found by Girty and myself. *S. lamellatus* is a very peculiar species without close relation in either America or Europe. The group of *S. pelargonatus*, so common in Europe, is represented by *S. pyramidalis*. The plicated type of *Streptorhynchus*, which has been called *Kiangsiella*, has not been found in this country, though it is rather common in Asiatic Permian formations. The new genus *Orthotetella* is without known relative. *Orthothetina*, a type common in the Permian faunas of LoPing and Djulfa, is represented by a few specimens in the Wolfcamp and the Capitan, the latter reported by Girty. There are 9 species of *Meekella*, only one of which, *M. irregularis texanus*, shows foreign affinity, so that this group, like *Rhipidomella*, may be regarded as largely of American origin. This genus is unrepresented in the Salt Range faunas and is rare in all other Permian faunas but that of the Trogkofel formation and the *Schwagerina* horizon of Russia. *Geyerella*, a genus occurring also in the Trogkofel and Sosio limestones, is present in the Hess and the Capitan. Of the 5 species of *Derbya*, *D. crenulata* probably evolved from *D. crassa* of the Pennsylvanian, *D. cymbula* occurs also in the Pennsylvanian, and *D. buchi* and *D. nasuta* are very similar to some Eurasian forms. The genus *Orthothetes*, which was found by Girty to be rather common in the Capitan, was not identified from the present collection. It is rare above the Mississippian, and present in few Permian faunas. On the whole, the Orthotetinae are more varied than in the Pennsylvanian. They are like the Pennsylvanian forms but with the addition of several foreign elements such as *Geyerella* and several species of *Streptorhynchus*. There is about the same number of species as in the Mississippian, but the generic types are quite different, the genera *Schuchertella*, *Schellwienella*, and *Leptaena*, which are important in the Mississippian, being quite absent in the Permian, while *Meekella*, *Geyerella*, and *Derbya* are absent from the Mississippian.

The representation of the genus *Chonetes* is rather diversified, as in most other Permian faunas, 11 species being present. Two of these occur also in the Pennsylvanian, 5 have Eurasian affinities, and the remainder are descended from American species or are so different from other described types as to be of uncertain origin. The striate and smooth types of ornamentation are the only ones present in these species. Forms bearing large costae, which are common in some Asiatic faunas, are absent, as well as the genera *Chonetella* and *Chonetina*, which likewise occur in Asia. Several of the species of *Chonetes* in this fauna are modifications of Pennsylvanian species, but many additional types are introduced. A large number of species of *Chonetes* is present in the Mississippian, but they are mostly quite unrelated to those in the Permian faunas.

The genus *Isogramma* ranges up into the Wolfcamp formation, but is unknown elsewhere in Permian faunas except in Mongolia.

About a third of the entire fauna consists of the species formerly placed in the genus *Productus*. Ten genera in that group are here recognized. Nineteen species of *Productus* s. s. were distinguished. No other Pennsylvanian or Permian faunas have yielded as many species of this genus except the Uralian fauna described by Tschernyschew. For example, only 7 species of *Productus* s. s. are present in the Salt Range fauna. Of the species in the Texas Permian, only 2 occur in the Pennsylvanian. *Productus inca*, common in the Gym fauna, is present in the Andean "Upper Carboniferous." Four species are identical with northwestern North American and Arctic forms, while 4 have very close affinity with Eurasian species. The remaining species probably are purely American types.

The genus *Linoproductus* is represented by 6 species, 2 of which belong to the group of *L. cancrini (Cancrinella)*. *L. cora* is abundant in the lower beds of the section and a variety of it extends as high as the Leonard formation. *L. waagenianus* is the most abundant Permian species of the genus, being one of the most common fossils of the Leonard, Word, and Capitan formations. It is related on the one hand to *L. aagardi*, a common Russo-Arctic species, and on the other is apparently identical with forms from the upper Permian of the Himalayas. In the *L. cancrini* group, *L. phosphaticus* is close to if not identical with *L. cancriniformis*, while *L. villiersi* is closely related to sev-

eral Upper Carboniferous and lower Permian species of the group in Eurasia and America.

The genus *Striatifera*, which is generally a less common type than *Linoproductus*, is represented by a single form. A species apparently belonging to this genus occurs also in the Gaptank formation, but the group is unrepresented elsewhere in the Pennsylvanian.

*Pustula*, so common in the Mississippian and Pennsylvanian, is represented solely by the Pennsylvanian species *P. semipunctata*. There are three species of *Buxtonia*, one of them a Pennsylvanian one, and another, *B. peruviana*, identical with Andean and Uralian forms. Two species of Buxtonia, *B. occidentalis* and *B. longa*, are present in the American Pennsylvanian. *Waagenoconcha*, the group of *Productus humboldti*, is represented by two species, one of Andean and Uralian affinities and the other related to a Salt Range species. It is probably almost wholly a Permian type. *Overtonia* is doubtfully identified on the basis of its resemblance to an Andean species in which the generic characters were definitely determined. It is present in the Cisco and possibly in the Mississippian.

*Avonia*, a genus almost absent from the American Pennsylvanian, is represented by 7 species, two of which are related to Eurasian forms and one identical with an Andean species, most of the others being of undeterminable origin. The group of *Productus horridus* and *P. timanicus*, *Horridonia*, though very common in the Eurasian boreal realm during Upper Carboniferous and Permian times, here contains but a single species, of which only one specimen was found.

There are 13 species of *Marginifera*, which is a very characteristic member of Permian faunas in other parts of the world. Two forms are very similar to those of the Tethyan region, another is identical with an Andean species, while two others occur in the Pennsylvanian. Several groups may be recognized in this genus. The typical group has a semireticulate ornamentation, as in *M. capaci*, *M. dugoutensis*, *M. lasellensis*, and *M. reticulata*. *M. cristobalensis* has a tuberculose decoration. *M. manzanica*, *M. whitei*, and *M. wordensis* have pustules and wrinkles in the early stages, which are replaced later by costae. *M. opimus*, *M. popei*, and *M. texana* are costate only, while *M. sublaevis* is nearly smooth. Independent evolution appears to have taken place during the Permian along these different lines.

On the whole, the productids of this fauna are very much like those of other Permian assemblages. The main difference is the small representation of *Horridonia*, but in this respect the Texas fauna resembles those of the Tethyan region, where *Horridonia* is seldom present. A definite Permian aspect is given by the presence of *Waagenoconcha*, *Linoproductus waagenianus*, *Avonia signata*, and the abundance of *Marginifera*. There are striking differences from the Pennsylvania faunas. The species of *Productus* s. s. which do not have a wrinkled surface over the visceral area (as *P. guadalupensis)* are foreign to the Pennsylvanian, as well as *Striatifera*, *Waagenoconcha*, *Avonia*, *Horridonia*, and the tuberculose, costate, wrinkled, and smooth types of *Marginifera*. On the other hand, the genera *Pustula* (group of *P. semipunctata)* and *Juresania* (group of *P. nebraskensis)* are practically unrepresented in the Permian, but are very abundant in the Pennsylvanian. The productids of the American Mississippian are somewhat less diversified generically than those of the Permian, only *Productus*, *Linoproductus*, *Striatifera* (in the Cordilleran region), *Avonia*, *Pustula*, and possibly *Overtonia* being present, of the genera occurring in the Permian. The Mississippian genera *Productella*, *Diaphragmus*, and *Kensuella* (group of *P. giganteus*, present in the Cordilleran region) do not occur in the Pennsylvanian or Permian.

The genus *Scacchinella* is a characteristically Permian form occurring also in the Trogkofel and Sosio limestones, the Permian of Ferghana, and the Akasaka limestone of Japan. The related genus *Tschernyschewia* has not been found in the Permian of North America.

There is an exceptional representation of *Aulosteges*, 8 species having been recognized. *Aulosteges guadalupensis* and *A. medlicottianus* are related to or identical with Salt Range species. The genus does not occur in the Pennsylvanian, and may be regarded as characteristic of the Permian.

*Strophalosia* is rare, in contrast to its abundance in the Permian of Eurasia. It is likewise uncommon in the Pennsylvanian, but perhaps more abundant in the Mississippian.

The Richthofenidae are abundant in the Permian of west Texas, but present a peculiarly American aspect, to some extent unlike that seen in Europe

and Asia. Primitive species of the genus *Teguliferina* occur in the Pennsylvanian in North America, but apparently not in Eurasia. It appears from this fact that the family is of American origin. The Wolfcamp species *T. bösei* is very similar to the genotype *T. deformis*, of the Trogkofel limestone. This species gave rise to *Prorichthofenia teguliferoides* of the Hess formation, and that in turn to the three more advanced species of *Prorichthofenia*. It is very likely that the Eurasian genus *Richthofenia*, which has not been found in America, arose from the American genus *Prorichthofenia* by specialization of the muscle-bearing organs. The genera *Gemmellaroia* and *Tectarea* are further Eurasian specializations of this group and have not been found in America. Probably the early part of the evolution of this family took place in America, only occasional migrants being present in Eurasian faunas until early middle Permian time, when the American genus *Prorichthofenia* sent its modified descendant *Richthofenia* into the Tethyan region, the later modification and specialization of the family taking place there. The presence of *Prorichthofenia* in the Permian is a striking difference from the Pennsylvanian, while neither *Prorichthofenia* nor *Teguliferina* occurs in the Mississippian.

Unlike the Richthofenidae, the Lyttoniidae appear to have been of Eurasian origin, for the complete chain of evolution of that family can be reconstructed from Eurasian faunas, but only disconnected links are present in America. The most primitive genus of this family, *Keyserlingina*, occurs in the *Schwagerina* stage of the Uralian, but is not present in America. Somewhat lower stratigraphically is the occurrence in the Cisco of *Poikilosakis*, but as that genus is more specialized than *Keyserlingina* it is very probable that the latter actually made its first appearance at a lower horizon. In Russia, *Poikilosakos* occurs above the first appearance of *Keyserlingina*. *Parakeyserlingina*, probably the direct descendant of *Keyserlingina*, occurs in the lower Permian of Asia and in the Wolfcamp formation. *Lyttonia*, derived from *Parakeyserlingina*, is of worldwide distribution in middle and upper Permian faunas between lat. 40° north and 10° south. It is abundant in the Hess, Leonard, Word, and Capitan formations. The genus *Oldhamina*, which is common in the upper Permian of India and China, has not been found in America. The presence of the group of lytoniids is a very striking contrast to the faunas of the Mississippian and Pennsylvanian.

The Rhynchonellidae are more diversified than those of the Pennsylvanian, but contain practically the same genera. The genus *Leiorhynchus*, which is represented also in the lower and middle Pennsylvanian by *L. rockymontanus*, has two Permian species, one of which occurs in the Phosphoria, and the other is related to the Phosphoria species. *Leiorhynchus* cannot be definitely identified from Eurasian Permian faunas. *Pugnoides* contains 7 species, most of them very close to the Pennsylvanian *P. osagensis*. *Pugnoides* is almost wholly an American type, being very rare in most of the Upper Carboniferous and Permian faunas of Eurasia. There are two species of *Rhynchopora*, which is a very characteristic genus of the European Permian. *R. illinoisensis*, which occurs rarely in the Wolfcamp formation, is present, perhaps somewhat modified, in the Upper Carboniferous and Permian faunas in North America, the Andean region, and Europe. *R. taylori*, the Word species, is probably more closely related to *R. illinoisensis* than to the European Permian forms; it probably is an Arctic modification of the *R. illinoisensis-R. nikitini* type, which entered Texas by way of the Phosphoria seaway. There is but one abundant species of *Camerophoria*, *C. venusta*, which is very close to the common Eurasian species, *C. purdoni*. *C. thevenini* is present in the "Upper Carboniferous" of Bolivia, while *C. deloi* is a form occurring in the Gym facies of southern Arizona. The new genus *Uncinuloides* is probably an American representative of a widespread Eurasian group which has been called *Uncinulus*. It is one of the most characteristic forms of the Permian of the Tethyan geosyncline. One genus which is common in Eurasian Permian faunas, *Terebratuloida*, has not been found in America. Thus it may be said that the rhynchonellids are, on the whole, very similar to those of the Permian faunas of Eurasia, but differ in the presence of *Leiorhynchus*, the abundance of *Pugnoides*, and the absence of *Terebratuloidea*. In the Pennsylvanian, *Camarophoria* and *Uncinuloides* are not present, and the group of rhynchonellids is much less diversified than in the Permian. The Mississippian, on the other hand, is remarkable for the abundance and variety of its rhynchonellids. While the Permian contains the genus *Uncinuloides*, that does not

occur in the Mississippian, the latter deposits contain a large number of genera, most of them represented by many species, which do not occur in either the Pennsylvanian or Permian.

There are 10 species of *Spirifer*, of which 8 show the typical late Paleozoic expression, *Neospirifer*, having fasciculate costae. *Spirifer condor* occurs in the Andean "Upper Carboniferous" and in the Uralian, and *S. marcoui infraplica* is very close to the common Eurasian species *S. marcoui*. *S. costella* is the American variant of the Eurasian species *S. fasciger*. The most abundant *Spirifer* is *S. pseudocameratus*, which clearly descended from *S. triplicatus*, the common Pennsylvanian species. *S. mexicanus*, which is apparently the most common form of the Capitan, is closest to *S. oldhamianus* of the Salt Range faunas. An important member of the fauna is *S. sulcifer* Shumard, which is abundant in the middle and upper Word. It is extremely close to *S. rajah*, a very common species in the upper Permian of Asia, and to *S. lyra*, a Russo-Arctic species. The genus *Spiriferella* common in the Russo-Arctic realm of late Paleozoic faunas, is completely lacking in the Texas Permian.

The Pennsylvanian species *Squamularia perplexa* occurs in the basal Permian; it is succeeded in higher beds by *S. guadalupensis*, the longest ranged species of the Glass Mountains section, which is close to the Indian species *S. indica*. The Pennsylvanian *Ambocoelia planoconvexa* also occurs in the basal Permian; in the upper part of the Glass Mountains section there is another species of the genus, which occurs also in the Phosphoria. *Ambocoelia* is all but absent from Eurasian Permian faunas. It occurs only in the Zechstein fauna. *Martinia*, a genus not present in the American Pennsylvanian, is abundant in the basal Permian of the Glass Mountains. The nearest relative of the Wolfcamp species is *M. triqueta* of the *Schwagerina* horizon of the Uralian. *M. rhomboidalis* of the higher Permian is likewise of Eurasian affinity. The genus *Martiniopsis*, which is very common in Eurasian Permian faunas, does not occur in Texas.

As a whole, the Spiriferidae are similar to those of the Pennsylvanian, but with some different elements. As in the Pennsylvanian, the neospiriferoid type predominates in the genus *Spirifer*. The Permian fauna is, however, very strikingly different from that of the Mississippian. In that system, the genus *Spirifer* is represented by quite different specific groups. *Syringothyris, Pseudosyrinx, Spiriferella, Brachythyris, Delthyris*, and *Cyrtia*, which are rather common in the Mississippian, do not occur in the Permian. The representation of species, as well as of genera, is much greater in the Mississippian than in the Pennsylvanian or Permian.

As in the Mississippian and Pennsylvanian, the *Spiriferina* group comprises *Spiriferina* s. s. and *Punctospirifer*. The species of these genera in the American Permian are very similar to those in the Tethyan region. The genus *Cyrtina*, which occurs in the Mississippian and in the Permian Sosio limestone, is unrepresented in the American Permian.

The Rhynchospirinidae are represented solely by *Hustedia*, which contains 6 species, one of which, *H. meekana*, is close to or identical with the common Salt Range species, *H. indica*. The genus *Uncinella*, which occurs in the Permian of India, is not found in Texas. In the Pennsylvanian the only genus belonging to this family is *Hustedia*, but in the Mississippian *Hustedia* is uncommon, while *Eumetria, Ptychospira*, and *Acambona* are well represented.

The Meristellidae and Atrypidae have each one genus in the Mississippian, but none in either the Pennsylvanian or Permian.

Among the Athyridae, only *Composita* is present, but this genus shows considerable diversity. Most of the species are related to the Pennsylvanian *C. subtilita*, but *C. emarginata affinis* is very close to *C. xetra* of the Himalayan Upper Permian. The genera *Spirigerella* and *Cliothyridina*, which are conspicuous elements of Eurasian Permian faunas, are quite absent from the Texas Permian. This is the only family in which a common Pennsylvanian genus is unrepresented, for *Cliothyridina*, an abundant fossil in places in the lower and middle Pennsylvanian, does not occur higher in this country. The Athyridae of the Mississippian resemble those of the Permian in having *Composita*, but contain also the genera *Camarophorella, Rowleyella, Athyris*, and *Cliothyridina*.

The terebratulids are rare in the present collection, and are represented by few species and genera, and two genera *(Notothyris* and *Heterelasma)* described by Girty were not recognized. As in other late Paleozoic faunas, *Dielasma* is the most important terebratulid genus. One species of it is very close to several forms of the Salt Range Permian, while *D. bovidens* occurs in the lower part of the Texas Permian section. There are one

or two species of *Dielasmina*. *Notothyris* and *Heterelasma* have been identified by Girty from the Capitan; the former is rather doubtful. The genus *Hemiptychina*, which is rather common in the Permian of Eurasia, is entirely absent from the present collection. The Permian of West Texas resembles the Pennsylvanian in the rarity of the terebratulids, and contrasts strikingly with the Mississippian, where there are 6 genera, of which only *Dielasma* occurs in the Permain.

In summary it may be said that the Permian faunas resemble in a general way those of the Pennsylvanian, but contain practically all the Pennsylvanian genera with the addition of a good many that are either restricted to the Permian, or occur in pre-Pennsylvanian faunas but not in the American Pennsylvanian. The presence of the Richthofenidae and Lyttoniidae and the large number of Productidae are among the most striking features of the Permian faunas. In contrast to the Mississippian, the Permian probably will prove to be as rich when fully collected, but has some notable differences. The Orthacea and Orthotetinae are more diverse in the Permian, and are represented by almost entirely different genera. There is a much larger number of productids, including several different genera. *Scacchinella*, *Aulosteges*, the Richthofenidae, and the Lyttoniidae, though common in the Permian, are quite unlike anything in the Mississippian. On the other hand, the Permian contains but few rhynchonellids and spiriferids, in contrast to the large number of these stocks in the Mississippian. The Rhynchospirinidae and Athyridae are much less varied than they are in the Mississippian, and the Terebratulidae are rare and of different types than those of this period.

## DESCRIPTIONS OF BRACHIOPOD GENERA AND SPECIES

### Order NEOTREMATA

### Family DISCINIDAE

#### Genus ORBICULOIDEA d'Orbigny

*Diagnosis* (after Weller): Shell subcircular or subelliptical in outline, inequivalved. Pedicle valve nearly flat or depressed-convex, with apex excentric and inclined slightly toward the posterior margin; a pedicle groove originates at the apex of the valve, passes backward for a short distance, beyond which point it is continued as a pedicle tube which pierces the substance of the valve very obliquely, opening to the interior near the posterior margin. Brachial valve depressed conical, with the apex excentric and inclined backward. Surface of the valves smooth, or more commonly marked by concentric lines, and more rarely by fine radiating costae.

Genotype: *Orbicula morrisi* Davidson.

*Geologic range:* Ordovician to Cretaceous.

#### ORBICULOIDEA MISSOURIENSIS (Shumard)?

Pl. I, Fig. 1

1858 *Discina Missouriensis.* Shumard, St. Louis Acad. Sci., Trans., vol. I, p. 221.

1910 *Lingulidiscina missouriensis.* Girty, Fauna of the Phosphate Beds of the Park City Formation, U. S. Geol. Surv. Bull. 436, pp. 22–24, pl. 1, figs. 6–10. (See for synonymy.)

Phosphate beds of the Park City formation: Woodruff Creek, Utah; Sublette Range, Wyoming; Thomas Fork, Wyoming; Montpelier, Idaho.

*Description:* Shell phosphatic, subcircular in outline. The apex of the brachial valve lies at about a fifth the diameter from the posterior margin, and is distinctly curved downward, so that the highest point is a little anterior to the apex. There are numerous fine concentric growth lirae. Width, 8 mm.

*Observations:* This identification is made on the authority of Girty's discussion in the "Fauna of the Phosphate Beds," and is not based on study of typical representatives of the species from the Pennsylvanian. There is but a single specimen, which is, however, well preserved, as it came from a limestone concretion.

*Distribution:* Waagenoceras horizon (Las Delicias) at section 2, bed 4.

### Family CRANIIDAE

#### Genus CRANIA Retzius

*Diagnosis* (after Weller): Shells subcircular in outline, generally more or less unsymmetrical in growth, inequivalved. Pedicle valve without perforation for the passage of a fleshy pedicle, attached by its apex or by its entire external surface to some external object. Brachial valve depressed-subconical, with a subcentral, erect, or backward directed apex. The interior of each valve is marked by two pairs of large adductor muscle scars, the posterior pair being close to the margin and widely separated, the anterior pair being much closer together and subcentrally located. The external surface of the brachial valve marked by more or less irregular concentric lines of growth, by regular radiating costae, or by fine setae or pustules.

Genotype: *Anomia craniolaris* Linne.

*Geologic range:* Ordovician to Recent.

#### CRANIA? sp.

1908 *Crania* sp. Girty, The Guadalupian Fauna, U. S. Geol. Surv. Prof. Paper 58, p. 156.

Middle Capitan, Capitan Peak (sta. 2926); dark limestone, Pine Spring (sta. 2930).

*Observations:* Under this heading is placed a single brachial valve, 7 mm. in diameter, deprest conical, with a somewhat excentric apex. The surface is marked by concentric lirae. It agrees well with the description by Girty of two specimens of Crania from the Guadalupe Mountains.

*Distribution:* Middle Word formation (Glass Mountains) at 239.

Superfamily ORTHACEA

**Genus RHIPIDOMELLA Oehlert**

*Diagnosis:* Shells subcircular, biconvex, brachial valve somewhat deeper than the other. Hinge-line short. Surface covered by fine, rounded subequal lirae* which are hollow and open out to the surface. Pedicle valve bearing two strong diverging teeth; from the base of each there extends forward a more or less strongly defined curving ridge, bordering the muscular area, which is large, deeply impressed, and flabelliform. Adductor scars small and completely enveloped by the diductors. Cardinal process erect, strongly arched on its posterior face, posterior surface trilobate. Dorsal muscular area small, quadruplicate, divided longitudinally by a broad, low median ridge extending forward from the base of the cardinal process.

Genotype: *Terebratula michelini* L'Eveille.

*Geologic range:* Silurian to Permian.

**RHIPIDOMELLA CARBONARIA (Swallow)**

1858 *Orthis carbonaria.* Swallow, Trans. Acad. Sci. St. Louis, vol. 1, p. 218.

1877 *Orthis pecosii.* White, U. S. Geol. Expl. w. 100 Mer., vol. 4, p. 125, pl. 9, figs. 5a-e.

*Observations:* Except for two specimens that are exceptionally large and one from loc. 90 having rather coarse lirae, these specimens are quite like the average form of the Pennsylvanian species in so far as their characters may be seen in their fragmentary condition. The specimens identified by Waagen, Schellwien, and Tschernyschew as *R. pecosi* are almost certainly not this species. Fliegel* was evidently wrong, however, in believing the Salt Range and Alpine forms to be identical with the Chinese *R. subquadrata,* a species which is much more closely related to *R. hessensis.*

*Distribution:* Gaptank formation at 75 and 273. Wolfcamp formation (Glass Mountains) at 86, 87 low?, 88, 90, 93, 189x, 196?, and 198. Not common.

**RHIPIDOMELLA HESSENSIS King. n. sp.**

Pl. I, Figs. 2–4

*Description:* Shell medium-sized, strongly biconvex, brachial valve very convex, pedicle valve smaller and less convex, or even flattened anteriorly. Greatest width at about a third of the length behind the anterior margin, which is straight or very slightly convex, and not at all emarginate. Cardinal area of the pedicle valve triangular, well-defined, rising at about 50° from the dorsal area, then incurving so as to lie parallel to it; delthyrium an equilateral triangle, nearly filled by the projecting cardinal process. Cardinal area of the brachial valve lower than the other, slightly incurved ventrally. Cardinal process large, projecting ventrally through the delthyrium and divided posteriorly into three diverging lobes, of which the median one is much the largest and most elevated. Muscle scars deeply impressed.

Surface marked by fine, rounded lirae, separated by narrower striae. On the anterior lateral parts of the pedicle valve and on most of the surface of the brachial valve there are large, equal-sized, forward-projecting tubules arising from the lirae; they are closely spaced near the anterior margin on large shells, but are widely scattered over the posterior halves of the valves. Growth lamellae present anteriorly on mature shells. The shell is very thick.

Dimensions in mm. of three specimens from locality 107:

| | 1 | 2 | 3 |
|---|---|---|---|
| Length | 16 | 17 | 20 |
| Width | 17 | 17 | 21 |
| Thickness | 8 | 9 | 11 |
| Height of ventral area | 1.5 | | 2 |

*Comparisons:* This species differs from most other species of *Rhipidomella* having similar configuration and size (such as *R. penniana, R. nevadaensis, R. oweni,* and *R. michelini)* in the large size and the abundance of tubules. It is closest to *R. subquadrata* (Fliegel) as figured by Frech; both species have abundant and large tubules and have similar convexity, but the Chinese species differs in having a broader posterior part and a slight sinuosity of the front margin.

*Distribution:* Upper Hess formation (Glass Mountains) at 1, 3, 8, 15, 98, 105, 106, 107 (abundant, types), T107, T108, and 222. Upper Hess formation (Sierra Diablo) at 478 and 479.

**RHIPIDOMELLA LEONARDENSIS King n. sp.**

Pl. I, Figs. 5–7

*Description:* Shell small, pentagonal in outline, with the greatest width at mid-length, hinge-line a little more than half the greatest width of the shell, anterior and lateral margins nearly straight but curving evenly into each other.

Pedicle valve deepest at about a third the length from the beak; median part of the shell flattened or slightly depressed; beak not incurved. Cardinal area a third as high as wide, lying at about 75° to the plane of the valve margin, flattened but curving toward the beak; delthyrium a fourth to a fifth as wide as the hinge, higher than wide. Dental lamellae strong and somewhat diverging; a strong but low median septum arises close under the beak and extends about half the length of the shell; muscle scars very elongate, with nearly parallel borders.

Brachial valve equally convex with the other, with the greatest convexity at about the middle, beak short and extended, with the cardinal area nearly in the plane of the valves.

Surface marked by lirae of very irregular size, each strong one being generally separated from the next of the same size by from one to three finer ones which do not reach back to the beak, but are intercalated between the coarse lirae; lirae 3 to 4 in the space of one mm. at the anterior margin. Striae punctate, so that the lirae have a crenulated appearance. The lirae open out into tubules fairly closely spaced in the posterior part of the shell but

*In this and the following descriptions, the word "lirae" will be used for the radial ridges on a brachiopod shell and "striae" for the grooves.

*Palaeontographica, Bd. 48, p. 126.

sparse in the anterior part. Strong concentric widely-separated growth lines are present.

Dimensions in mm. of two specimens:

| | 1 (loc.123) | 2 (loc.240) |
|---|---|---|
| Length | 8 | 11 |
| Width | 8 | 12 |
| Height of ventral area | 1.5 | 2.5 |

*Observations:* This species differs from *R. carbonaria* by its flattened median part of the pedicle valve, higher cardinal area, narrow delthyrium, strong dental lamellae, and different type of liration. The last character is not easily seen in badly silicified specimens.

*Distribution:* Leonard formation (Glass Mountains) at 12, 26S, ?82, 120, 123 (types), 232?, T240, and 484? This species is not abundant at any of the localities named, but is fairly common at 26S and 123. Word formation? (Glass Mountains) at 44–45?

### RHIPIDOMELLA MESOPLATYS KING, n. sp.

Pl. I, Figs. 10–11

*Description:* Shell small, subovate, with the greatest width at mid-length, hinge-line equaling a third the greatest width of the shell, anterior margin rounded.

Pedicle valve depressed at about a third of the length from the beak; beak slightly incurved. Cardinal area a fourth as high as wide, lying at about 80° to the plane of the valve margin, strongly incurved near its end. Delthyrium higher than wide.

Brachial valve equally convex with the other, the greatest convexity being near the center. Beak as high as that of the pedicle valve, incurved at the end. A broad, low median depression or flattening originates about 4 mm. in front of the beak and expands gradually toward the anterior margin.

Surface marked by even lirae, 2 or 3 in the space of 1 mm. at about mid-length, opening out on the anterior part of the shell into unevenly distributed tubules.

Dimensions in mm. of two specimens:

| | 1 (loc. 123) | 2 (loc. 256) |
|---|---|---|
| Width | 14 | 13 |
| Length | 14 | 13 |
| Thickness | 9 | 8 |
| Height of ventral area | 1.75 | — |

*Observations:* This species differs from *R. leonardensis* in the different character of its liration, its shorter hinge, the generally more rounded outline, and probably also by a lower convexity. It differs from *R. hessensis* in its constantly smaller size, higher cardinal area and much coarser liration. The two species are, however, very closely related, and it is likely that *R. mesoplatys* descended from *R. hessensis.*

*Distribution:* Leonard formation (Glass Mountains) at 20, 26s?, 123 (types), T233?, D3, D5, D8, D12, and D22. Word formation (Glass Mountains) at T41, 54, T144a, 241, and 256.

### RHIPIDOMELLA MESOPLATYS BAYLORENSIS King, n. var.

Pl. I, Figs. 8–9

*Description:* This variety differs from typical *R. mesoplatys* in having a slightly lower ventral area, which diverges at about 56° from the plane of the shell margin, then curves up against the dorsal area, which is inclined ventrally at an angle of about 30°. The ventral beak is higher than the other. Apparently more tubules arise from the lirae than in the typical form of the species.

*Distribution:* Leonard formation (Baylor Mountains) at 502 and 503 (type). Bone Canyon member (Sierra Diablo) at 485.

### RHIPIDOMELLA TRANSVERSA King, n. sp.

Pl. I, Figs. 12–13

*Description:* Shell small, transversely subovate, rather strongly biconvex. Ventral cardinal area narrow and high, its width being approximately equal to about two-fifths that of the shell, and its height about half as great as its width. Delthyrium narrow. The ventral area lies at an angle of about 60° to the shell margin, and is evenly curved upward toward its tip. The dorsal cardinal area is less than half as high as the other. Surface marked by narrow radial lirae separated by broad striae, and on one shell, bearing concentric growth undulations. One specimen has a shallow, narrow dorsal sinus. Shell finely punctate.

Dimensions in mm. of two specimens:

| | 1 (loc. 123) | 2 (loc. T15a) |
|---|---|---|
| Length | 10 | 9 |
| Width | 12 | 11 |
| Height of ventral area | 2 | — |

*Observations:* This species is distinguished by its greater width than length, its high, narrow ventral cardinal area, and its narrow lirae with broad striae.

*Distribution:* Hess formation (Glass Mountains) at 4 and 17a. Leonard formation (Glass Mountains) at T15a, 123 (type), and D5.

### Genus ORTHOTICHIA Hall and Clarke

*Diagnosis:* Externally like *Schizophoria.* Ventral muscular area not impressed in the shell, divided longitudinally by a thin, elevated median septum and bordered laterally by prominent dental lamellae, which lie at a low angle to the septum; at the anterior end of the muscular area the dental lamellae are united to the septum by a thickened margin. Dorsal interior similar to that of *Schizophoria.*

Genotype: *Orthis? morganiana* Derby.

Range: Pennsylvanian and Permian.

*Observations:* This genus is generally regarded as representing an intermediate stage between *Schizophoria* and *Enteletes.* Schellwien[1] has argued that it is not a natural group, but is composed both of species intermediate between *"Orthis"* and *Enteletes* and of species of *Enteletes* which have arisen from an evolution of plicated to unplicated forms, supposedly in the early Permian. However, most Permian species of *Enteletes* are even more strongly plicated than the Pennsylvanian ones, and those Permian species which have faint plications do not approach the *Schizophoria*-like external form of a true *Orthotichia.* Even more significant is the fact that the arrangement of muscular areas in the pedicle valve of *Enteletes*

[1]Die Fauna der Trogkofelschichten, pp. 3–5.

(but not *Paraenteletes)* is quite unlike that in *Orthotichia* and *Schizophoria*—in the former the internal plates lie parallel, while in the latter they diverge at a greater or less angle. It would be surprising if, in losing its plications, a regressive *Enteletes* would also assume primitive internal characters. *Enteletes derbyi demissa* Schellwien appears clearly not to be a variety of *Orthis derbyi* Waagen, for the dental lamellae are nearly parallel to the median septum in Schellwien's figure,[2] while in Waagen's figure[3] the typical internal characters of *Orthotichia* are shown. Girty* has discussed Schellwien's opinions regarding *Orthotichia,* but he fails to emphasize the importance of internal characters in permitting easy distinction between smooth-shelled orthids transitional between *Schizophoria* and *Enteletes,* and those arising from the loss of plications in *Enteletes.*

**ORTHOTICHIA KOZLOWSKII King, n. sp.***
Pl. I, Figs. 14–15

1914 *Orthotichia Morgani.* Kozlowski (non Derby), Brachiopodes du Carb. Super. de Bolivie, Ann. de Pal,. t. 9, p. 62, pl. 3, figs. 11–12.

*Description:* Shell medium-sized, wider than long; brachial valve much deeper and more inflated than the other. Pedicle valve sinuate.

Pedicle valve longitudinally of flattened curvature, and transversely of low convexity. Hinge-line equal to half the width of the shell; cardinal area high, lying at about 70° to the plane of the shell margin, slightly concave. Beak pointed, slightly incurved. Median sinus originating about 10 mm. in front of the beak, extending forward and slightly indenting the margin. Lateral parts smooth. In the interior the median septum and dental lamellae extend more than half the distance to the anterior margin; the dental lamellae are connected at the front to the median septum by transverse plates.

Brachial valve much more strongly convex in the posterior than in the anterior part. Beak strongly incurved over the cardinal area. A very low and broad median fold appears to be present corresponding to the sinus of the other valve. The crural plates diverge from the median septum at an angle of 40°.

Surface marked by fine subequal lirae. Certain ones are slightly stronger than others, and give rise to tubules.

Dimensions in mm. of two specimens:

| | 1 (loc. 92—type) | 2 (loc. 456) |
|---|---|---|
| Length of brachial valve | 22 | 15 |
| Length of pedicle valve | 20 | 14 |
| Width | 24 | 19 |
| Thickness | 19 | — |
| Height of ventral area | 3.5 | — |

*Observations:* This species differs from *O. morgani* (Derby) in being much smaller and less strongly sinuate, and from *O. schucherti* Girty, from the Hermosa of Colorado, in being much more strongly convex and having a larger cardinal area. *O. kozlowskii* was identified from Bolivia by Kozlowski as *O. morgani,* but seems clearly to be conspecific with the lower Permian form of Texas. *O. derbyi* (Waagen) is very similar to *O. kozlowskii,* but has a nearly erect beak, while the beak of the Texas form is strongly incurved. The sinus is on the brachial valve in *O. indica* (Waagen), and *O. marmorea* (Waagen) differs by having no sinus at all.

*Distribution:* Wolfcamp formation, *Uddenites* zone (Glass Mountains) at T88 and 175. Wolfcamp formation (Glass Mountains) at 90, 92 (type), and 196–197. Upper Wolfcamp formation (Sierra Diablo) at 456. Gym formation (Hueco Mountains) at 395? Leonard formation (Glass Mountains) at D1 and D16.

**Genus ENTELETES Fischer von Waldheim**

*Diagnosis:* Shells biconvex, the brachial valve being the deeper. Hinge-line short. Anterior part of the shell more or less plicated. If, as is generally the case, the median plication is a fold or sinus, the fold is in the brachial valve and the sinus in the pedicle valve. Ventral interior bearing two strong elongated teeth on both sides of the delthyrium, supported by strong dental lamellae, which extend forward, first approaching each other, then lying parallel and close together. Median septum lying between the dental lamellae, highest near its anterior end, where it rises to a sharp crest. Crura sickle-shaped, directed laterally; crural plates strong, diverging at a high angle antero-laterally. Cardinal process a short narrow ridge.

Genotype: *Enteletes choristites* Fischer de Waldheim.

Range: Pennsylvania and Permian.

**ENTELETES DUMBLEI Girty**
Pl. II, Figs. 4–8

1908 *Enteletes dumblei.* Girty, The Guadalupian Fauna, U. S. G. S. Prof. Paper 58, p. 295, pl. 26, figs. 4-4b. Hueco formation, Diablo Mountains (sta. 3764).

1908 *Enteletes angulatus.* Girty, The Guadalupian Fauna, U. S. G. S. Prof. Paper 58, pp. 295–296, pl. 26, figs. 3-3a. Hueco formation, Diablo Mountains (sta. 3764).

*Description:* Shell large, wider than long, the brachial valve very convex, the pedicle valve a little less so, with the greatest convexities at about mid-length.

Cardinal area of the pedicle valve well-defined, about half as long as the greatest width of the shell and about one-fourth as high as long, its basal part flat and lying at approximately right angles to the plane of the shell margin, with its apical part curving sharply over it under the beak. Ventral beak incurved over the delthyrium, which occupies nearly a third of the area. Sinus originating about 10 mm. in front of the beak, fully twice as large as the lateral sulci; lateral plications coarse, subrounded to angular, numbering 4 on each side of a shell 3.5 mm. wide. The three septa on the inside of the valve are straight and slightly diverging. They extend half the length of the shell. The median septum rises to its highest point about two-thirds the distance from the beak to its anterior end; this point is sharply angular, the top of the septum curving from it to each end.

Brachial valve rather evenly convex, bearing a subrounded median fold and 3 or 4 plications on each lateral

[2]Ibid., p. 1, fig. 7.

[3]Waagen: Pal. Indica, ser. 13, vol. 1, pl. 56, fig. 2.

*U. S. Geol. Sur. Prof. Paper 58, pp. 291–293.

*This may be the species described by Girty as *Schizophoria huecoensis,* Jour. Wash. Acad. Sci.

slope; the median fold has a similar form to but fully twice the size of the adjacent plications. Entire surface covered with fine lirae, which are narrower than the striae.

Dimensions in mm. of typical specimens:

| | Girty's type | 1 (loc. 222) | 2 (loc. 264) | 3 (loc. 240) | 4 (loc. 479) |
|---|---|---|---|---|---|
| Length | 22 | 30 | 22 | 29 | 17 |
| Width | 26 | 33 | 29 | 29 | 22 |
| Thickness | 26 | 25 | — | — | — |
| Height vent. area | 4.5 | 3.5 | — | — | 4.5 |

*Observations:* The only distinguishing characters given by Girty between *E. dumblei* and *E. angulatus* are the high arching of the ventral valve and the angularity of the plications of the latter, characters which are somewhat variable in this species. The fact that the types of both species were from the same lot (sta. 3764) strengthens the conclusion that *E. angulatus* is a synonym. Even if *E. angulatus* Girty were not a synonym, Girty's name cannot be used, as it is preoccupied by the name *Rhynchonella angulata* Geinitz (probably synonymous with *E. hemiplicatus).*

*E. dumblei* differs from *E. hemiplicatus* in being much larger, in its stronger plication, and in its higher cardinal area. According to Girty, the two are distinguished also by a different type of liration, *E. dumblei* having supposedly narrower lirae and broader striae, but this character appears to vary on different parts of a single specimen. Girty believed *E. dumblei* to be close to *E. elegans* Gemmellaro, but that form has much smaller plications than *E. dumblei* has at the same distance from the beak. However, the Texas species is very close to *E. waageni* Gemmellaro, but does not attain nearly as large a size. *E. dumblei* can hardly be distinguished from the specimen of *E. andii* d'Orbigny figured by Kozlowski,[1] the greater depth of the median sinus of the Bolivian shell being the only distinctive feature noticeable. The Bolivian species has been figured also by Douglas,[2] who comments on the close similarity to species from the Hueco formation *(=E. dumblei).* To judge from his figures, they are indeed very similar.

*Distribution:* Wolfcamp formation (Glass Mountains) at 93? Hess formation (Glass Mountains) at 35, 107 (common), 108 (abundant), T108, 164, 191, and 222 (abundant). Probable Hess formation (Glass Mountains) at T9, T15a, and Tba. Hess formation (Sierra Diablo) at 465, 476, 478, and 479. Bone Canyon member (Sierra Diablo at 484 and 485. Gym formation (Diablo Plateau) at 435? and 443?. Gym formation (Hueco Mountains) at B392 and 389. Word formation (Glass Mountains) at 137, 141, 154, T239, 241, T249, T264, Tx148, and Tx. Leonard? formation (near Shafter) at 515, Tj, and Tq.

*Enteletes dumblei* is apparently nearly confined to the Hess and Word formations and their equivalents. Its absence from the typical Leonard would seem to be good evidence for the supposition that the Hess and the Word specimens are actually different species. Since, however, no distinguishing characters were discerned, it is not possible to separate them.

[1]Ann. de Pal., 1914, pl. 3, figs. 10a-c.

[2]Quart. Jour. Geol. Soc., vol. 70, pp. 35–36, pl. 8, figs. 2a-2c.

### ENTELETES LEONARDENSIS King, n. sp.

Pl. III, Figs. 10–12

*Description:* Shell medium-sized, with a highly inflated brachial valve and a pedicle valve of low convexity, the mid-part of the valve being nearly flat. In an old shell the pedicle valve is only half as long, measured along the curvature, as the brachial, but in most specimens the disparity is less. In all, however, there is a greater difference than in other species of *Enteletes* in this fauna. The anterior margin is so flattened by the strong curvature of the brachial valve that the valves meet at a very obtuse angle. Ventral cardinal area high and not much incurved at its end. The median fold and sinus originate about 10 mm. anterior to the beaks, and are not much larger than the lateral plications and sulci. The plications are low and rounded, and number three to four on each side of the fold and sinus.

Dimensions in mm. of several specimens:

| | 1 (loc.128) | 2 (loc.37) | 3 (loc.288) |
|---|---|---|---|
| Length of pedicle valve | 19 | 19 | 20 |
| Length of brachial valve | 23 | 24 | 21 |
| Width | 24 | 23 | 24 |
| Thickness | 22 | 22 | 19 |
| Height of ventral area | 4 | — | 4 |

*Observations: E. leonardensis* is closely related to *E. wordensis,* having, like it, relatively small fold and sinus and a similar number of lateral plications. The low convexity of the pedicle valve and the large, strongly convex brachial valve, however, distinguish it from that and other species of *Enteletes.* It differs from *E. dumblei* in its smaller size, lesser width in proportion to length, greater difference between the size of the two valves, and narrower fold and sinus.

*Distribution:* Leonard formation (Glass Mountains) at T15, 16, 26s, 28s, 37, 120, 128, T128, D1, and D4.

### ENTELETES LIUMBONUS King, n. sp.

Pl. II, Figs. 9–11; Pl. III, Figs. 1–2

1908 *Enteletes* sp. c. Girty, The Guadalupian Fauna, U. S. G. S. Prof. Paper 58, pp. 297–299, pl. 24, figs. 4-5a, pl. 26, figs. 1-2b.

Basal black limestone, Guadalupe Point (stas. 2920 and 2967). Diablo Mountains (sta. 3764).

*Description:* Shell medium-sized or large when mature, but mature forms are less common than the immature ones. Since, however, all gradations exist from small, nearly unplicated shells to the mature ones, there is evidently but one species. Brachial valve considerably larger than the other, mature shells very strongly inflated, while the pedicle valve is rather strongly convex but much lower.

The cardinal areas and beaks resemble in every respect those of *E. dumblei.* The internal characters likewise are similar. The umbonal regions for about 13 mm. anterior to the beaks are unplicated. Plications low and rounded, numbering, on mature shells, three, with a faint fourth, on each side of the fold and sinus. One shell from loc. 381

has the fold and sinus divided respectively into two lesser folds and sinuses. Lirae rounded, separated by narrower striae.

Dimensions in mm. of several specimens:

| | 1 (loc. 17) | 2 (loc. 17) | 3 (loc. 106) | 4 (loc. 17) |
|---|---|---|---|---|
| Length of brachial valve | 14 | 19 | 22 | 25 |
| Length of pedicle valve | 15 | 17 | 20 | 21 |
| Width | 17 | 21 | 26 | 29 |
| Thickness | 12 | 12 | 22 | 25 |
| Height of ventral area | 1.5 | 2 | 2.5 | 2.5 |

*Observations:* This species is clearly very close to *E. dumblei*. It is distinguishable by its more strongly inflated brachial valve when mature, by its unplicated umbonal region, and by its low, rounded plications. The immature individuals collected by Girty were almost unplicated, but his description leaves no doubt that this is the form designated by him as *E.* sp. c. The great number of half-grown shells is very striking; the fact that all gradations in size may be collected from the same piece of rock shows that there is but one species. Girty compared *E.* sp. c with many faintly plicated foreign species, but since his faintly plicated shells were immature, such comparison is unnecessary here.

*Distribution:* Hess formation (Glass Mountains) at 1, T15, 17 (abundant types), 105, 106, and 210. Probably also immature specimens from 3 and 8. Leonard? formation (Glass Mountains) at 245. Gym formation (Hueco Mountains) at 381, 383?, and 384? Hess formation (Sierra Diablo) at 467 and 479. *E Liumbonus* occurs at the same horizon in the Hess as *E. dumblei*, and in some places the two occur together.

### ENTELETES PLUMMERI King, n. sp.

Pl. III, Figs. 3–8

*Description:* Shell small, slightly wider than long, valves evenly convex, the brachial valve being somewhat larger than the other. Ventral cardinal area equal to or somewhat less than half the greatest width of the shell, and about a third as high as wide; beak strongly incurved. There are three pairs of angular lateral plications on the pedicle and four on the brachial valve of most specimens. The fold and sinus are distinctly larger than the lateral plications and sulci; they originate about 8 mm. in front of the beaks.

Dimensions in mm. of several specimens:

| | 1 (loc. T9) | 2 (loc. T9) | 3 (loc. 231) | 4 (loc. 104) |
|---|---|---|---|---|
| Length | 18 | 17 | 19 | — |
| Width | 19 | 18 | 20 | 27 |
| Thickness | 16 | 17 | — | 26 |
| Height of ventral area | 3.5 | 3.5 | 4 | 6.5 |

The specimen from loc. 104 is exceptionally large, the others being about normal size.

*Observations:* *E. plummeri* is most nearly related to *E. hemiplicatus*, but may be distinguished from it by the relatively smaller size of the plications, and the relatively smaller fold and sinus. It is very similar to the group of Eurasian forms which have been divided into the species *E. oehlerti* Gemm., *E. contractus* Gemm., *E. elegans* Gemm., *E. meridionalis* Gemm., *E. subaequivalvis* Gemm., *E. haugi* Gemm., and *E. tschernyscheffi* Diener. As Diener[1] has pointed out, these species are so narrowly restricted that there is considerable doubt as to their validity. If, as is quite possible, they all are members of one variable species, then it would not be possible to distinguish *E. plummeri* from it. *E. plummeri* is similar in every detail to *E. oehlerti* as figured by Schellwien.[2]

*Distribution:* Upper Hess formation (Glass Mountains) at 8. Lower Leonard formation (Glass Mountains) at T9, 104, T205, 231, 232, T232, 128? (having smaller plications than typical specimens), and Tb. *E. plummeri* is common at nearly all these localities; it is apparently restricted to the highest beds of the Hess and the lowest of the Leonard.

### ENTELETES WOLFCAMPENSIS King n. sp.

Pl. III, Fig. 9

*Description:* Shell small, globose. The posterior parts of a mature shell have a very low curvature; at about three-fifths the length of the shell from the beak there is a rounded geniculation, anterior to which the curvature is very low to the front margin, where the two valves meet at an angle of about 120°. This is seen only on mature specimens; some immature forms are too small to have reached the stage of geniculation. The brachial valve is somewhat deeper than the other. The flattened posterior regions show the beginnings of a fold and sinus on the brachial and pedicle valves respectively, but the lateral plications originate at the geniculation: they are low and rounded and are two to three in number on each side. One specimen shows strong growth undulations. Lirae strong and even. The shell is finely punctate. The internal plates of the pedicle valve are arranged as is characteristic of *Enteletes*.

Dimensions in mm. of two specimens:

| | 1 (loc. 196) | 2 (loc. B88) |
|---|---|---|
| Length | 16 | 17 |
| Width | 17 | 18 |
| Thickness | 18 | 19 |

*Observations:* This species is easily distinguished by the geniculation of both valves, the slight curvature anterior and posterior to the geniculations, and the absence of plications over the entire posterior three-fifths of the shell. *E. contractus* Gemm. is similar in having a geniculated profile, but has much stronger plications.

*Distribution:* Wolfcamp formation (Glass Mountains) *Uddenites* zone at B88, middle Wolfcamp at 93 and 93s, upper Wolfcamp at 196. The species is rare at each locality.

### ENTELETES WORDENSIS King, n. sp.

Pl. III, Figs. 13–15

1908 *Enteletes* sp. a. Girty, The Guadalupian Fauna, U. S. G. S. Prof. Paper 58, pp. 296–297, pl 30, figs. 2 and 2a.

"Comanche Canyon," Glass Mountains.

[1]Pal. Indica, ser. 15, vol. 1, part 5, pp. 28–30.

[2]Die Fauna der Trogkofelschichten, vol. 16, pl. 1, figs. 11–13.

1908 *Enteletes* sp. b. Girty, The Guadalupian Fauna, U. S. G. S. Prof. Paper 58, p. 297.
"Comanche Canyon," Glass Mountains.

1908 *Enteletes* sp. d. Girty, The Guadalupian Fauna, U. S. G. S. Prof. Paper 58, pp. 299–300, pl. 21, fig. 23.
Delaware Mountain formation, Guadalupe Point, Guadalupe Mountains (sta. 2919). Delaware Mountain (sta. 3501). "Delaware Mountain formation," Glass Mountains (sta. 3763?).

*Description:* Shell medium-sized to large, subcircular in outline, evenly biconvex, the brachial valve being somewhat the larger. Cardinal area of the pedicle valve flat for most of its length, strongly incurved at the beak; its length equals half the width of the shell and its height a little more than a fourth its length. Fold and sinus only slightly larger than the lateral plications and sulci; in some specimens they are of nearly equal size. Plications originate about 8 mm. anterior to the beaks; there are three to four on each side of the fold and sinus. Internal septa of the pedicle valve parallel, the median septum reaching its highest point near its anterior end, as described for *E. dumblei.*

Dimensions in mm. of several specimens:

| | 1 (loc. 264) | 2 (loc. 251) | 3 (loc. 265) |
|---|---|---|---|
| Length | 21 | 32 | 27 |
| Width | 23 | 28 | 26 |
| Thickness | 20 | — | — |
| Height of vent. area | 2.5 | — | — |

*Observations:* This species is distinguishable by its relatively narrow and numerous plications and the proportionately much smaller fold and sinus than in *E. dumblei* and *E. hemiplicatus.* It is constantly larger than *E. plummeri,* but has about the same number and relative size of plications at the growth stage when *E. wordensis* is the size of mature *E. plummeri. E. wordensis* is unlike *E. leonardensis* in having even convexity and no great difference in the size of brachial and pedicle valves. Several imperfect specimens of *Enteletes* were described by Girty. *E.* sp. a is, with little doubt, to be assigned to this species, the size and number of plications being the same as those of *E. wordensis. E.* sp. b Girty is less certainly identical, but it is likely that that shell was flattened from its normal profile. The character of the liration, cited by Girty as a distinction, is so easily obscured on silicified specimens such as he had that any differences between the ornamentation of one shell and another are probably to be regarded as due to silicification.

*Distribution:* Word formation (Glass Mountains) at 154?, 241, 251, 255, 264, 265, and Tc. Delaware Mountain formation (Sierra Diablo) at 504. It is not common. At localities 241 and 264 it is associated with *E. dumblei.*

### Genus PARENTELETES King, novum

*Diagnosis:* Plicated and finely lirate Enteletinae in which the fold is in the pedicle valve and the sinus in the brachial. The oldest species of the genus, *P. suessi* and *P. cooperi,* have, as in *Orthotichia,* divergent dental lamellae united anteriorly to the median septum by a thickening of the shell at the anterior ends of the adductor muscle scars. Middle and late Permian species described by Waagen have sub-parallel dental lamellae as in *Enteletes.*

Genotype: *Parenteletes cooperi* King, n. sp.

*Geologic range:* In America the earliest representative of *Parenteletes* is *P. cooperi,* which first appears in the lower Gaptank, in beds of middle Canyon age. The same form occurs again in the *Uddenites* zone of the basal Wolfcamp, of very earliest Permian age. The only other American species is *P. globosus* (Girty), which probably was collected from the Word formation. The earliest Eurasian representative is *P. suessi acuticosta* (Schellwien), from the Auernig formation of Upper Carboniferous age. The overlying Trogkofel formation contains typical *P. suessi* (Schellwien). In the Salt Range the earliest *Parenteletes* is *P. ferrugineus* (Waagen) from the Katta; then follow successively *P. sublaevis* (Waagen) and *P. pentameroides* (Waagen) from the Virgal, *P. latesinuatus* (Waagen) from the Kalabagh, and *P. acutiplicatus* (Waagnen) from the Khundghat or Jabi. *P. akasakensis* (Ozawa) is a species similar to *P. suessi* which occurs in Japan in the lower Akasaka limestone of lower Permian age.

*Origin and relations of the genus:* It has been suggested by Waagen and Girty that the "dorsisinuate *Enteletes" (Parenteletes)* evolved from the "ventrisinuate *Enteletes" (Enteletes)* through the development of a median plication in the sinus. Three arguments are used in favor of this: first, that *Schizophoria* and *Orthotichia* are ventrisinuate, and that it would be unlikely for a dorsisinuate shell to evolve from such a form without an intermediate stage in which a plication appeared in a ventral sinus, second, that the "dorisinuates" appeared only in the late Carboniferous or Permian, while the "ventrisinuates" appeared earlier; and third, that *E. lamarcki* and *E. hemiplicatus* show transitionary forms toward "dorsisinuate" shells in the appearance in the fold and sinus of a plication and a sulcus. While it is possible that the first argument is valid, the second and third are quite wrong. Actually, the "dorsisinuate group" *(Parenteletes)* was the first to appear. In North America *Parenteletes* appears in the middle Canyon (lower Gaptank), while *Enteletes hemiplicatus* is first found in the Thrifty (Cisco) and Lansing (Missouri). In Europe it is impossible to determine the sequence, for *Parenteletes* occurs first in the Upper Carboniferous of the Alps (Auernig formation) and *Enteletes lamarcki* in the *Spirifer mosquensis* zone of Russia; probably these formations are of approximately the same age. Further, the presence of a plication in the sinus and a sulcus in the fold of a true *Enteletes* is of common occurrence, but in every instance the plication and sulcus are distinctly smaller than the fold and sinus and could not be mistaken for it. But by far the most convincing fact is the *Orthotichia*-like arrangement of septa and muscle scars in the early *Parenteletes (P. suessi* and *P. cooperi),* in contrast to the specialized narrow muscular areas bounded by parallel dental lamellae and median septum in the earliest species of *Enteletes hemiplicatus* and *E. lamarcki.* It would be a reversal of evolution for the primitive internal characters to have appeared in a species which had developed from *Enteletes,* with its

specialized internal characters. Apparently, then, *Parenteletes* appeared before *Enteletes* in the Pennsylvanian, and both genera evolved independently from *Orthotichia.*

It must be made clear that specimens of *Enteletes* which contain a plication in the sinus and a sulcus in the fold are not to be included in *Parenteletes.* It is possible that some species included in *Parenteletes* did evolve from *Enteletes* in the manner suggested by Girty, but certainly *P. suessi* did not, and probably the rest of the genus is monophyletic.

#### PARENTELETES COOPERI King, n. sp.

Pl. I, Figs. 16–20; Pl. II, Figs. 1–3

*Description:* Shell medium-sized to large, wider than long, the brachial valve much larger than the other and strongly convex, the pedicle valve very slightly convex. Pedicle valve transversely elliptical, beak erect, the curvature being only slight along the median line of the valve. Ventral cardinal area deeply concave, erect, equaling about two-fifths the width of the shell and about one-half as high as wide. Delthyrium large, equaling in width slightly more than a third that of the area. Dorsal cardinal area smaller than the other and lying at somewhat less than right angles to it. The pedicle valve bears a large, high, sharp median fold, originating about 8 mm. anterior to the beak, the curvature being so slight that its crest lies nearly parallel to the plane of the shell margin. Lateral plications number three on each side of the fold, the first one being about half the size of the fold; their crests are much lower than that of the fold. The brachial valve has a deep median sinus originating about 5 mm. anterior to the beak; it curves evenly downward to meet the fold of the other valve. Lateral plications three or four on each side of the sinus, the sulci being about half as large as the sinus. Lirae fine and even, separated by very narrow striae.

In the pedicle valve the dental lamellae diverge from the median septum at an angle of about 15°; at their anterior ends they are connected by a ridge extending backward to join the end of the median septum.

Dimensions in mm. of several specimens:

| | 1 (loc. 199) | 2 (loc. 199) | 3 (loc. 273) |
|---|---|---|---|
| Length of brachial valve | 22 | 29 | 25 |
| Length of pedicle valve | 19 | 25 | 24 |
| Width | 26 | 37 | 32 |
| Height of ventral area | 6 | 6 | — |

The specimen from 273, as all the other Gaptank specimens, has been crushed, so that the original ratios of dimensions cannot be determined.

*Observations:* This species is very close to *P. suessi* (Schellwien) and its variety *acuticosta* (Schellwien). It differs from them in having a narrower fold and sinus and larger lateral plications. It is likewise similar to *P. latesinuatus* (Waagen), but has a narrower fold and sinus, and the dental lamellae diverge and are united anteriorly to the median septum, rather than lying subparallel as in Waagen's species.[1]

[1]This species has been erroneously identified by Beede (U. of Tex. Bull. 1753, p. 39) as *E.* aff. *waageni* Gemm. and by Schuchert (A. J. S., ser. 5, vol. 14, pp. 385 and 391) as *E. hemiplicatus* and *E. angulatus.* All these names are those of species of true *Enteletes* rather than *Parenteletes.*

*Distribution:* Lower Gaptank formation at 273 and T65 (common at both). *Uddenites* zone of Wolfcamp formation at 88, 89, 94, 95, 95s, 199 (abundant), 201, and Tba.

#### Genus STREPTORHYNCHUS W. King

*Diagnosis:* Shell biconvex, hinge-line short, lateral and anterior margins rounded. Pedicle valve high and generally distorted, its cardinal area high, with a large delthyrium which is closed to the apex by the deltidium. Internally the delthyrial supporting plates are reduced to mere thickenings of the inner surface of the cardinal area at the margins of the deltidium; median septum absent. Brachial valve with a very rudimentary area or no area at all. Cardinal process high and without a very transverse base, the height depending on the elevation of the umbonal region of the pedicle valve.

Genotype: *Terebratulites pelargonatus* Schlotheim.

*Geologic range:* Mississippian to Permian.

#### STREPTORHYNCHUS LAMELLATUM King, n. sp.

Pl. IV, Figs. 1–3

*Description:* Shell small, hinge-line shorter than the greatest width of the shell. Pedicle valve high, conical. Cardinal area of the pedicle valve lying at about 120° to the plane of the shell margin, flat, somewhat higher than wide, with a convex deltidium occupying about the median fourth. Brachial valve convex, slightly wider than long. Along the hinge is a broad cardinal area lying parallel to that of the pedicle valve, with a slightly convex chilidium.

Surface marked by small, rough costae, about 6 occurring in the space of 5 mm. on the anterior part of the shell but finer in the posterior part. The costae are truncated at intervals by closely spaced concentric lamellae, which give the shell a scalloped appearance, as in *Atrypa.*

Internal characters obscured; cardinal process apparently large.

Dimensions in mm. of two specimens:

| | 1 (loc.245—type) | 2 (loc.105) |
|---|---|---|
| Length of brachial valve | 10 | 8 |
| Width | 12.5 | 9 |
| Height of ventral area | 5 | — |
| Height of dorsal area | 1.5 | — |
| Length of hinge | 10 | 8 |

*Observations:* This species differs from the other Orthotetinæ in this fauna by its ornamentation and by the presence of a large dorsal area. Though the internal characters are not definitely known, they appear to be like those of *Streptorhynchus.*

*Distribution:* Upper Wolfcamp formation (Glass Mountains) at 105 and 211. Leonard formation (Glass Mountains) at 83, 228, and 245 (type).

#### STREPTORHYNCHUS PYGMAEUM Girty

Pl. IV, Figs. 4–6

1908 *Streptorhynchus pygmaeum.* Girty, The Guadalupian Fauna, U. S. Geol. Sur. Prof. Paper 58, pp. 178–180, pl. 30, figs. 3–6b.

Delaware Mountain formation, "Comanche Canyon," Glass Mountains; Delaware Mountain formation, southern Delaware Mountains (sta. 2962?).

1908 *Streptorhynchus?* sp. a. Girty, *idem*, pp. 178–180, pl. 30, fig. 7.
Delaware Mountain formation, "Comanche Canyon," Glass Mountains.

*Description:* Shell very small, unsymmetrical, outline polygonal, the greatest width being near mid-length. Pedicle valve larger than the brachial, with its beak strongly twisted to one side. The cardinal area lies at 90 to 140° to the margin, and its height is equal to about .4 the width; it is defined by an angulation from the main part of the shell. Deltidium convex in the median part, its width equal to 0.3 that of the cardinal area. Internal characters not determinable. According to Girty, "the progression of the hinge teeth is marked by two prominent ridges, which are not, however, prolonged as plates, while anything in the nature of a median septum is quite absent."

Brachial valve somewhat wider than long, the hinge-line shorter than the greatest width below. Cardinal process of a small brachial valve from loc. 26s lying at right angles to the plane of the margin, relatively small, marked by 4 ridges perpendicular to the hinge; crural plates diverging from each other at right angles, straight, short.

Surface marked by thin, high, more or less irregular and alternating, widely spaced and somewhat crenulated lirae. Undulations of growth present near anterior margin. Scars of attachment were not seen on the specimens examined, though all three of Girty's showed scars.

Dimensions in mm. of several specimens:

| | 1 (loc. 26s) | 2 (loc. 26s) | 3 (loc. T240) | 4 (loc. T148) |
|---|---|---|---|---|
| Length of brachial valve | 7 | 6 | 7 | 13 |
| Width | 9 | 8 | 9.5 | 17 |
| Length of hinge | 8 | 6 | 7 | 13 |
| Height of cardinal area | 3 | — | 3.5 | 4 |

The specimen from loc. T148, as well as one from Tx148, is unusually large.

*Observations:* Girty divided his specimens of this type from the Glass Mountains into two groups, one of which, represented by a single specimen, he described as *S.* sp. a. He notes the fact that the latter is possibly immature, as indicated by its thin shell. The differences between *S.* sp. a. and *S. pygmæum* were the thinner and higher lirae of *S.* sp. a, the absence of crenulations and varices of growth, and the smallness of the cardinal process. The specimens above described show very little crenulation of the lirae, the lirae being probably more like those of *S.* sp. a than of typical *S. pygmæum*, but varices of growth are present as in typical *S. pygmæum*. The cardinal process of the small, thin valve described above also is like that of *S.* sp. a. These features point to a greater similarity to *S.* sp. a than to *S. pygmæum*, but the nature of the internal characters is probably affected by the immaturity in both *S.* sp. a and the small brachial valve above described. The differences in the character of the liration are so slight, and this feature is so variable even on different parts of the same shell, that it can hardly be regarded as of specific importance. The likelihood that they are the same species is further indicated by the fact that they occur together at the place from which they were originally described.

*S. pygmæum* shows no close similarity to any foreign species of the genus, nor is there any American species except *S. undulatus* that has such coarse liration.

*Distribution:* Leonard formation (Glass Mountains) at T9 and 26s. Word formation (Glass Mountains) at T148, Tx148, and T240. Hess formation (Sierra Diablo) at 478. Upper Gym formation (Hueco Mountains) at 440a.

### STREPTORHYNCHUS PYRAMIDALE King, n. sp.

Pl. IV, Figs. 7–10

*Description:* Shell medium-sized, with very elongate ventral beak. Pedicle valve with a very large cardinal area, beak not sharply pointed or twisted to any great extent. Cardinal area a little wider than the shell beneath it, so that it forms a flange-like projection at its sides; the area is inclined at about 135° to the valve margin and is more than twice as high as wide. Median septum and dental lamellae absent.

Brachial valve subcircular, hinge-line equaling in length about a third the width of the shell. Cardinal process long, bifid, curving backward until nearly parallel to the valve margin. Anterior face of the process with a deep, broad depression extending from the end nearly to the center of the valve, where it broadens out into the deeply impressed muscular area, which is divided by a low median septum; the posterior face of the process divided by three shallow grooves radiating from the beak, separating 4 low ridges; process united anteriorly with broad, diverging socket plates, from which slight ridges are prolonged around the muscular area.

Surface marked by strong radial lirae, 9 to 11 occupying the space of 5 mm., separated by very narrow striae; the lirae are unequal because of the intercalations of small ones which do not reach complete equality with the older lirae for some distance anterior to the point of origin. Interruptions of growth are indicated by periodic, widely spaced, thin concentric striae. Shell very thick.

Dimensions in mm. of a brachial valve from loc. 93: length, 12; width, 14; length of hinge, 10. One pedicle valve must be more than 25 mm. long. A valve from loc. 93, evidently a pedicle valve, which appears to belong to this species, has a slight mesial depression and a rather incurved beak. These characters do not appear to agree with those of the other pedicle valves of this species from the same locality. The specimen from loc. 511 has a strong deflection of the margin in a ventral direction; this may be in part due to crushing. This specimen agrees well with the Wolfcamp ones, though from a much higher horizon. The type material is fragmentary, the specific characters being determined from several specimens. The range in size and in form could not be determined.

*Observations:* This species belongs to the group of *S. pelargonatus*, the common Eurasian Permian species, but differs from that form in having a much higher and flatter cardinal area, more strongly defined from the rest of the shell, and by lacking the mesial depression of the brachial valve so prominent in the foreign species. The ornamentation is very much alike. *S. lenticularis* Waagen is similar, but it differs in the same respect as *S. pelargonatus*, and may be identical with the latter species. *S. pseudopelargonatus* Broili, from Timor, is related, but even more different

than the others, especially differing in the nature of its sculpture. *S. subpelargonatus* Fliegel from Lo Ping is evidently similar, as it has no sinus, but as it was not figured, satisfactory comparison cannot be made.

The most similar American species is *S. pygmaeum* Girty, which is much smaller at maturity, has its cardinal area lower and less well defined from the antero-lateral margins, and has a much shorter cardinal process.

*Distribution:* Wolfcamp formation (Glass Mountains) at 88? and 93 (type). Leonard formation (Baylor Mountains) at 503?. Leonard formation (Finlay Mountains) at 511. Limestone of Cerro Caballo (Las Delicias), at section 1, bed 11?.

**STREPTORHYNCHUS? UNDULATUM King, n. sp.**

Pl. IV, Figs. 11–13

*Description:* Shell rather large, with a high, conical pedicle valve distorted by very strong irregular undulations. Cardinal area lying at right angles to the plane of the margin as high as or higher than wide, flat except for an upward curving at the beak, which is twisted to one side. Deltidium narrow, occupying about a fourth of the area, but where the shell has been exfoliated exposing the interior mold it appears to be much broader. The surface of the area is vertically striated. Within the pedicle valve there are thickened ridges under the edges of the delthyrium. Internal plates probably not present, as sections across the beaks show no such structures.

Surface marked by fine, uneven radial lirae, about 18 of which occur in the space of 5 mm., separated by broad striae.

Dimensions in mm. of several specimens:

| | 1 (loc. T9) | 2 (loc. T9) | 3 (loc. 211) |
|---|---|---|---|
| Length of pedicle valve | 35+ | 32 | 39 |
| Length of brachial valve | 32 | 24 | 30 |
| Width | 18+ | 20 | 15 |
| Height of cardinal area | 22 | 21 | 24 |
| Length of hinge-line | 30 | 26 | 30+ |

*Observations:* This species is very similar externally to *Orthothetina* sp. but differs in having a higher cardinal area and more undulating surface. It resembles also some of the species of *Orthotetes* described by Girty. The cardinal area is high as in *O. distortus,* but the shell is larger and has a less even surface. The specimen figured by Schellwien as *Streptorhynchus* aff. *operculatus* Waagen is very similar, and, indeed the two may prove to be identical.

*Distribution:* Hess formation (Glass Mountains) at T9, 17, 105, and 211 (type).

**Genus ORTHOTETELLA King, novum**

*Diagnosis:* Shell similar externally to *Orthotetes,* the pedicle valve being flat in longitudinal profile, the brachial valve slightly convex. Surface radially lirate, unplicated. Cardinal area very wide, rather high, divided into a primary and secondary area as in *Orthotetes.* Internally the deltyrial supporting plates are produced forward and inward, joining to form a cone or spondylium of subcircular cross-section, its apex at the beak. No median septum present.

Genotype: *Orthotetella wolfcampensis* King, n. gen., n. sp.

*Geologic age:* Lower Permian (Wolfcamp formation).

*Observations:* The determination of the internal characters, on which the above description was based, was made from a single excellently preserved large pedicle valve. The other specimens referred to the type species were so classified because of their external features.

The presence of a spondylium would appear to indicate that *Orthotetella* is a pentamerid, but the other characters make it clear that it is one of the Orthotetinae, and the presence of uniting delthyrial supporting plates show that it is closely related to *Orthotetes,* the only difference being the absence of a median septum and the complete merging of the two plates without a line of division at their junction. It is likely, therefore, that this is a further specialization of the camerate Orthotetinae, the group of *Orthotetes,* in which the median septum is lost and the delthyrial supporting plates fuse to form a spondylium. *Orthotetella* would therefore be one of the most highly specialized genera of the Orthotetinae.

**ORTHOTETELLA WOLFCAMPENSIS King, n. sp.**

Pl. IX, Figs. 27–28

*Description:* Shell large, symmetrical, wider than long, the hinge-line equaling or slightly less than the greatest width of the shell. Pedicle valve flat or slightly concave in longitudinal profile; beak pointed but low; cardinal area .25 as high as wide, rising nearly perpendicularly from the plane of the margin, with a deltidium occupying about a sixth of the width of the area which is convex and transversely striated. The lateral parts of the area are divided in two parts by a line extending diagonally from the beak, on the outer part of which the striae run at right angles to the hinge. Delthyrial supporting plates join to form a conical spondylium, with its point at the beak, firmly joined to the posterior part of the shell by a thickening around the top of the cone. Median septum absent. Muscular scars large and flabellate, causing the outer surface of the spondylium to be grooved.

Brachial valve very slightly convex, with strong crural plates, and the cardinal process marked on its posterior face by four ridges.

Surface marked by low, thin lirae, about 6 to 8 in the space of 5 mm., separated by very broad, flat striae, about 3 times as broad as the lirae. The lirae increase by intercalation, and as a result are somewhat alternating.

Dimensions in mm. of two specimens from loc. 93:

| | 1 | 2 |
|---|---|---|
| Length of pedicle valve | 40+ | 25+ |
| Width | 100 | 50? |
| Height of cardinal area | 29 | 9 |
| Width of cardinal area | 100 | 50 |
| Width of deltidium | 15 | 6 |

*Observations:* This species may easily be recognized even without the knowledge of the internal characters by its narrow lirae and very broad, flat striae.

*Distribution:* Wolfcamp formation, *Uddenites* zone (Glass Mountains) at 88, T88, 95?, 175?, 195, and 201? Wolfcamp formation (Glass Mountains) at 2?, 85?, 87,

89?, 90, 92, and 93 (type). Lower Gym formation (Hueco Mountains) at 394?. Middle Gym formation (Hueco Mountains) at 421?.

### Genus ORTHOTHETINA Schellwien

*Diagnosis:* Shell biconvex, similar in form to *Orthotetes* or *Meekella.* Surface lirate but not plicated. Within the pedicle valve the delthyrial supporting-plates are, as in *Meekella,* long, and reach to the floor of the valve, which they meet along the median line. Other internal structures probably also as in *Meekella.*

Genotype: *Streptorhynchus crenistria* var. *eusarkos* Abich.

*Geologic range:* Upper Mississippian to Permian.

*Observations:* As various authors have pointed out, especially Thomas,* this genus is probably heterogeneous, including shells both transitional from some primitive form to *Meekella,* and shells which have descended from *Meekella* by the loss of their plications. Since the only real difference from *Meekella* is the absence of plications, it may be best to regard *Orthothetina* as a synonym of *Meekella.* On the other hand, some species that have been placed in *Orthothetina,* such as *O.* cf. *mutabilis* of the present collection, have much more the form of *Orthotetes* than that typical of *Meekella.* It is possible that these species have arisen independently of *Meekella* from *Orthotetes,* a change which would not be great, as it would necessitate only the splitting apart of the delthyrial supporting plates near the floor of the shell so that they would not lie together to form a septum as in *Orthotetes.* Because of this uncertainty it seems well to make use of the name *Orthothetina* for such forms.

### ORTHOTHETINA cf. ORTHOTHETES MUTABILIS Girty

Pl. IV, Figs. 14–15

*Description:* Shell medium-sized, wider than long, with the greatest width at mid-length; length of the hinge equal to 0.7 the width of the shell. Pedicle valve high, larger than the brachial, with a flat cardinal area extending at right angles from the plane of the shell margin; height of the area equal to from half the length of the hinge to as high as the hinge-line is long. The deltidium occupies about a third the width of the area. No median fold or sinus is present. Growth very irregular, the pedicle valve being undulated transversely and with the beak more or less twisted. Brachial valve moderately convex, with the greatest depth at about a third the length; beak low. Within the pedicle valve the delthyrial supporting plates converge from the inner side of the area to meet along the median line of the floor of the shell. Median septum absent.

Surface marked by very fine subequal radial lirae, which do not have straight courses, but swerve very slightly from side to side; about 20 lirae occur in the space of 5 mm.

Dimensions in mm. of the specimen from loc. 91: length of pedicle valve, 24; length of brachial valve, 22 width, 28; height of cardinal area, 12?; length of hinge, 23; width of deltidium, 7?.

*Observations:* Externally the specimens in the present collection are quite like the types from the Pennsylvanian of Idaho of *Orthotetes mutabilis**. However, three of the Texas specimens show clearly that the delthyrial supporting plates do not converge as in *Orthotetes* before reaching the floor of the shell, forming a median septum, but lie apart as in *Meekella.* Girty did not figure the part of the shell showing the arrangement of the internal plates, but one cannot help but think that an apparent camerate structure was given his specimens by deformation, possibly by the crushing together of the lower parts of the delthyrial supporting plates to form an apparent septum. The presence in the Pennsylvanian of trans-Pecos Texas of other forms in common with faunas of the same age in the northern Cordilleran region indicates the similarity of the faunal association in Idaho and Texas.

An unnamed form of *Orthothetina* was found by Girty in the Capitan. It cannot be compared adequately with the forms here under consideration. No foreign species of *Orthothetina* shows great similarity, probably the most closely similar is *O. circularis* Fliegel, of the Lo Ping fauna.

*Distribution:* Gaptank formation (Marathon Basin) at 75. Strawn group (Hueco Mountains) at 354. Wolfcamp formation (Glass Mountains) at 71?, 91, and 196?.

### Genus MEEKELLA White and St. John

*Diagnosis:* Shell biconvex, commonly sub-pyramidal. Hinge-line shorter than the greatest width of the shell. Ventral cardinal area elevated and generally distorted; deltidium occupying approximately a third of the area, its median part strongly convex. Hinge-teeth prominent. Delthyrial supporting plates relatively very elongated, reaching nearly to the center of the shell, sub-parallel to parallel in direction, converging downward. Brachial valve gibbous in the umbonal region; dorsal cardinal area rudimentary or linear. Crural plates strongly divergent, extended forward more than a third the length of the valve, supporting a high, erect cardinal process, which may extend nearly to the apex of the umbonal cavity of the other valve; the process is a somewhat curved, thin crescentric plate, bearing on its summit two slender median apophyses, below which are two stouter lateral projections having the form of extended, rounded lobes, which form the bases of the crura. Surface marked by more or less strong radial plications and by fine lirae.

Genotype: *Plicatula striatocostata* Cox.

*Geologic age:* Middle Pennsylvanian to Upper Permian (the citation from the Morrow by Mather is regarded as doubtful).

*Observations:* Unplicated shells having the internal characters of *Meekella* are here split off for convenience as *Orthothetina,* not, however, without some doubt as to the importance of this distinction.

### MEEKELLA ATTENUATA Girty

Pl. V, Figs. 2–7

1908 *Meekella attenuata.* Girty, The Guadalupian Fauna, U. S. G. S. Prof. Paper 58, pp. 205–206, pl. 24, figs. 7–9a, pl. 25, figs. 4–4d.

*Thomas, Ivor: British Carboniferous Orthotetinae, Mem. Geol. Sur. of Gt. Britain, Pal., vol. 1, pt. 2, pp. 91–92.

*Girty, U. S. Geol. Surv. Prof. Paper 152, p. 433, pl. 27, figs. 9–15.

Delaware Mountain formation (probably actually Hess), Diablo Mountains (sta. 3764). Basal black limestone, Guadalupe Point (sta. 2967).

*Description:* Shell medium-sized to large. Pedicle valve high, subconical, inclined backward; hinge-line equal to about half the greatest width of the shell. Beak somewhat contorted; one specimen has a truncated beak, as if due to attachment. The cardinal area makes at the base an angle of 90–150° with the plane of the shell margin, but curves upward toward the beak until parallel or nearly so with the plane of the valve margin. It is well-defined, flat transversely and concave longitudinally. Deltidium equal to about .4 the width of the area, slightly convex, poorly defined. Delthyrial supporting plates converge toward the bottom of the shell, where they lie subparallel. One specimen from loc. 144 shows a double angulation, resulting in lateral offset, similar to the structure described by Girty for *M. difficilis*, but since other specimens fail to show it, this character may be accidental.

Brachial valve subcircular, widest near mid-length, regularly rounded; very convex near the beak and slightly convex anteriorly. Cardinal process long, with two extended median apophyses and a small lateral projection on each side, forming the base of the crura.

Surface marked by 11–28 strong, sub-angular plications, which subside toward the beak and resolve more or less abruptly into a prominent unplicated region, on which the lirae are conspicuous, or (as is most common in the Hess and Leonard specimens) they may extend to the beaks, near which each plication bears only one or two lirae. A specimen from loc. 38 has very fine plications to the number of about 38; were it not that intermediate forms to the normal type are present it would surely be regarded as distinct. Lirae number 4–5 in the space of 1 mm., counting the smaller incipient lirae intercalated between the larger ones; they show some tendency on the anterior part to curve up toward the crests of the plications.

Dimensions in mm. of several individuals:

| | 1 (242) | 2 (144) | 3 (Tx148) | 4 (123) | 5 (481) | Girty's type |
|---|---|---|---|---|---|---|
| Length of pedicle v. | 24 | — | 21 | 16 | 24 | 14 |
| Length of brachial v. | 18 | — | 17 | 12 | 20 | 10 |
| Width | 18 | 32 | 19 | 15 | 23? | 10 |
| Thickness | 16 | — | 16 | 12 | — | — |
| Length of hinge | 11 | 24 | 10.5 | 10 | — | 7 |
| Height of card. area | 7 | 18 | 7.5 | 11 | 13 | 6.5 |

*Observations:* The posterior part of large specimens of this species agrees exactly with the small specimen taken by Girty for the type of *M. attenuata*. *M. attenuata* differs from *M. difficilis* in having a high cardinal area, fine, regular plications, and narrow form.

*Distribution:* Wolfcamp formation (Glass Mountains) at 75?. Hess formation (Glass Mountains) at 35, 107, 215, and 223. Hess formation (Sierra Diablo) at 476, 478, 479, and 481. Leonard formation (Glass Mountains) at 5, T9, 12, 14, 22, 24, 31, 37, 98?, T119, 120, 123 (common), 128, 226, 232, Tx, D5, D7, 10, and D17. Bone Canyon member (Delaware Mountains) at 490 and 491. Word formation (Glass Mountains) at 46, 51, 54, 55, 144, T144a, 148, Tx148, 150, 153, 170, 192, 238, 239, T239, T240, T240a, 242, 252, 264, and 265. Limestone of Cerro Caballo, near Las Delicias, Coahuila, about .25 mile south of 21 on bed 12. Leonard formation (Baylor Mountains) at 502. Victorio Peak member (Sierra Diablo) at 495.

### MEEKELLA DIFFICILIS Girty

Pl. IV, Figs. 16–17; Pl. V, Fig. 1

1908 *Meekella difficilis.* Girty, The Guadalupian Fauna, U. S. G. S. Prof. Paper 58, pp. 207–208, pl. 30, figs. 10–10g.

"Delaware Mountain formation," Comanche Canyon, Glass Mountains (sta. 3763), and mountains northwest of Marathon (sta. 3840).

1908 *Meekella mutilirata.* Girty, The Guadalupian Fauna, U. S. G. S. Prof. Paper 58, pp. 208–209, pl. 24, figs. 6–6b.

Basal black limestone, Guadalupe Point (sta. 2967).

*Description:* Shell medium-sized. Pedicle valve low and subpentagonal. Hinge equal to about 0.5 the greatest width of the shell; in the specimen from loc. 235, however, the ratio is 0.8, and in one from loc. 26s it is 0.7. Cardinal area inclined backward, well-defined, generally low and slightly incurved or, as in two specimens from loc. 128, nearly flat. Deltidium broad, poorly defined, occupying more than a third the width of the cardinal area. Delthyrial supporting plates subparallel, median septum absent. Brachial valve transverse, rounded, but contracting near the hinge-line, its convexity moderate, greatest near the umbo.

Plications angular and somewhat nodose, 9–14 in number, subequal, becoming less distinct laterally, extending in most specimens nearly to the beaks; they are separated by broad, less sharp sulci. Lirae rounded and subequal, generally numbering 4 in the space of 1 mm., or more in some specimens, curving up toward the crests of the plications, and in some specimens not paralleling the plications but diverging laterally and crossing them.

Dimensions in mm. of several specimens:

| | 1 (120) | 2 (26s) | 3 (235) | 4 (128) | 5 (D1) |
|---|---|---|---|---|---|
| Length of brachial v. | 26 | 25 | 25 | 16 | 17 |
| Width | 31 | 30 | 35 | 15 | 20 |
| Thickness | 19 | — | — | 13 | 15 |
| Width of cardinal ar. | 20 | 24 | 30 | 11 | 11 |
| Height of cardinal ar. | 9 | — | 9 | 9 | 6 |

*Observations:* Specimens of this species, as noted, show considerable variation. They are, as a whole, characterized by a longitudinally concave cardinal area, fine lirae, sharp, medium-sized, nodose plications originating near the beaks, and transverse outline. *Meekella pyramidalis* of the Kaibab differs in its larger size, and in the pyramidal shape of the pedicle valve due to the high cardinal area. *M. striatocostata* has less angular plications, which originate much farther from the beaks. In *M. skenoides* there is a much higher, flatter cardinal area and a conical pedicle valve, and its plications are generally more rounded

and do not reach as near the beak. *M. attenuata* is proportionally narrower, and has much finer plications and a higher cardinal area.

The specimens in the present collection do not differ in any important particular from Girty's description of *Meekella difficilis*, except that the peculiar internal characters noted in the original description were not seen here. It is very likely that *M. mutilirata* of Girty is identical with this species. The differences noted between them were the fact that the former has lower and less angular plications and finer lirae. From Girty's figures of *M. mutilirata*, however, it may be seen that the surface was largely exfoliated, so that the plications would appear to be rounded. In the collections at hand it appears that the strength of the lirae is very variable, and not a specific character.

*Distribution:* Leonard formation (Glass Mountains) at 4, 13, 15, T15a, 20, 22, 24, 26s, 27, 38, 104, T119, 120, 123, 124, 127, 128, 226, 228, T233, 235, Tb, D1, D3, D12, and D15. Leonard formation (Baylor Mountains) at 502. Bone Canyon member (Sierra Diablo) at 499. Leonard formation, transition beds (Shafter) at Th. Near Las Delicias, limestone of Cerro Caballo, bed 11, section 1; and bed 12, section 2. The species does not occur abundantly at any of these localities, most of the collections being of less than 4 specimens.

### MEEKELLA GLOBOSA King, n. sp.

Pl. V, Figs. 10–12; Pl. XXXII, Fig. 8

*Description:* Shell very large, globose, symmetrical. Pedicle valve strongly convex, with a greatly extended, conical posterior part. The cardinal area makes near the hinge an angle of about 120° with the plane of the shell margin, but curves upward toward the beak; cardinal area well-defined, evenly concave longitudinally. Deltidium prominent, equal in width to about 0.4 that of the area, with a sharply raised narrow median part. Delthyrial supporting plates sub-parallel; median septum absent.

Brachial valve subhemispherical, widest in front of mid-length, joining the pedicle valve along the anterior margin with a regular curve. Convexity great, deepest point a little back of the center. Cardinal process very long, supported by strong sub-parallel crural plates. The anterior face of the process is deeply concave and bordered on each side by the extensions of the crural plates. On the posterior face the hinge sockets are very deep, and lie near the hinge. The greater part of the process consists of the two median apophyses, which are greatly prolonged.

Surface marked by 14–21 sharp equal-sized plications, which extend to the beaks. Posterior part marked by equal sized lirae, about 5 of which occupy the space of 1 mm.; they disappear on the plications about 15 mm. in front of the beaks, but persist over the unplicated lateral margins of the shell. The anterior parts of the plications are marked by closely arranged chevron-shaped growth lines.

Dimensions in mm. of the type specimen:

| | |
|---|---|
| Length of brachial valve | 45 |
| Width | 45 |
| Thickness | 46 |
| Width of cardinal area | 25 |
| Probable height of cardinal area | 24 |

*Observations:* There is some variation among specimens of this species. The type specimen has exceptionally large plications and seems to be more strongly convex than the other less complete examples. The species is distinguishable, even if only small or incomplete specimens are found, by its narrow, sharp, regular plications extending to the beaks, and the absence of lirae except in the umbonal regions. Girty's type of *M. difficilis* has a somewhat similar aspect, but differs in having much larger plications in proportion to the size of the shell. *M. attenuata*, the only other similar type, differs from this in being smaller, incompletely plicated and completely lirated, and in having less angular plications.

*Distribution:* Leonard formation (Glass Mountains) at 128 (abundant), T128, and 151 (which may be the same as 128). The species appears to be restricted to a single locality: the fossiliferous limestone on the cuesta northeast of Split Tank on the Old Word Ranch.

### MEEKELLA GRANDIS King, n. sp.

Pl. VI, Figs. 5–7

*Description:* Shell large, transversely elliptical, the hinge-line equaling a little more than half the greatest width of the shell. Pedicle valve flat in profile, twisted in the posterior part. Cardinal area flat and high, lying at about 120° to the plane of the margin. Delthyrial supporting plates subparallel; median septum absent. Brachial valve of low convexity, the greatest curvature being in the umbonal region. A shell from locality 105 which is doubtfully referred to this species has a broad trilobate cardinal process, the median part of which consists of a high raised ridge, not very clearly bilobed as in most other Meekellas; the sides of the process consist of strong plates which, near the beak, bear dental-sockets. These lateral plates extend antero-laterally into short but strong crural plates. In a specimen from Tq the cardinal process appears to have a much broader posterior face, but this may be due to wearing off of the surface.

Plications low and generally broad, about 12–18 in number, bifurcating toward the front margin, separated by broad, round-bottomed sulci. The umbonal regions are unplicated. The badly worn specimen from 105 appears quite smooth over most of its surface, the first plications appearing 40 mm. in front of the beak; near its anterior margin the plications number only about 6, and are low and broadly rounded. Lirae medium-sized and rounded, about 4 occurring in the space of 1 mm., converging toward the crests of the plications near the anterior margin of the shell.

Dimensions in mm. of several specimens:

| | 1(174) | 2(Tq) | 3(211) | 4(105) |
|---|---|---|---|---|
| Length | 55 | 48 | 49 | 52 |
| Width | 68 | 60 | 62 | 56 |
| Height of cardinal area | 24 | — | — | — |
| Length of hinge-line | 38 | — | — | — |

*Observations:* The important characters of the shells referred to this species are the large size, low convexity of

both valves, unplicated umbonal region, and low plications separated by shallow, broad sulci.

*Distribution:* Wolfcamp formation (Glass Mountains) at 90?. Hess formation (Glass Mountains) at 105? and 211. Leonard formation (Glass Mountains) at 8–14?, and 174 (type). Leonard formation, transition beds (Shafter) at Tq (several specimens). Victorio Peak member (Brackett Draw) at Taa?.

### MEEKELLA IRREGULARIS TEXANA King, n. var.

Pl. VI, Figs. 1–4

*Description:* Shell medium-sized to rather large; the hinge-line equal to 0.7 the greatest width of the shell. Pedicle valve very high, distorted. Beak high and sharply pointed. Cardinal area poorly defined, about as high as, or higher than wide, deeply concave longitudinally. Deltidium about a third as wide as the area, with a narrow projecting elevation in the middle, the remaining part being flat, with longitudinal striae and some irregular transverse folds. The delthyrial supporting plates diverge from the margins of the deltidium and extend downward to the median line of the shell. Brachial valve only slightly convex. Within are strong, diverging crural plates and a weak median septum.

Surface marked by fine radial lirae which, near the shell margins, converge toward the crests of the plications. Plications, numbering about 12–15, low, narrow, rounded, and irregular, being nodose at intervals, particularly near growth crenulations, and somewhat crooked, especially on pedicle valves; in some specimens they increase toward the margins by bifurcation. Some shells are plicated nearly to the beaks, while on others they become faint and indistinguishable in the umbonal region. Irregularities in growth resulting in concentric depressions on the brachial valve and strong transverse folds on the pedicle valve give a very uneven form to the shell.

Dimensions in mm. of several specimens:

| | 1 (199-type) | 2 (95) | 3 (88s) | *M. irregularis* Sch. |
|---|---|---|---|---|
| Length of pedicle v. | 23 | — | 37+ | 33 |
| Length of brachial v. | 17 | 29 | 26+ | 26 |
| Width | 24 | 37 | 30 | 33 |
| Length of hinge | 13 | — | 23 | 28 |
| Height of cardinal a. | 17? | — | 21 | 17 |
| Width of deltidium | 5 | — | — | — |

*Observations:* These specimens are evidently nearly identical with *M. irregularis* Schellwien of the lower Permian Trogkofel formation of the Karnic Alps.* They have, however, a proportionately shorter hinge-line and wider deltidium. *M. procera* Schellwien, which occurs with *M. irregularis*, has a much greater length in proportion to width and less strongly developed incisions of growth. *M. evanescens* Schellwien is distinguished especially by weak plications, but also by the angular form and the extreme size and height of the cardinal area. *M. irregularis texana* is distinguishable from other American species of *Meekella* by its high cardinal area and small nodose plications. *M. striatocostata* has a low cardinal area and broad, regular plications originating some distance from the beak.*

*Distribution:* Wolfcamp formation, *Uddenites* zone (Glass Mountains) at 88, 88s, T88, 94, 95, 175a, 199 (type), 201, and 202. Middle and upper Wolfcamp formation (Glass Mountains) at 70?, 90?, 92, 93?, 93s, 169a, 196, and 205.

### MEEKELLA MEXICANA Girty

Pl. VII, Figs. 1–5

1909 *Meekella mexicana.* Girty, Manzano Group, U. S. G. S. Bull. 389, pp. 53–54, pl. 6, figs. 1–5.
San Andres limestone, Engle and Elephant Butte; Yeso formation, San Andres.

*Description:* Shell medium-sized, slightly unsymmetrical. Pedicle valve subtriangular, subconical, flattened or depressed, and extended backward. Length of hinge equal to about four-fifths the greatest width. Cardinal area high, nearly flat, and inclined strongly backward; its inclination varies from a position nearly normal to the margin to a position nearly parallel thereto, generally forming an angle of about 45° to the shell margin. Narrow median part of the deltidium sharply raised. Brachial valve strongly inflated, especially in the umbonal region; in many specimens somewhat extended in the median anterior part to produce a ventral deflection of the margin.

Plications low, originating between 10 and 40 mm. (but generally about 15–20 mm.) anterior to the beak of the brachial valve. In some specimens there are plications on the pedicle valve but none on the other. Lirae fine, narrow, and sharp, numbering 3–4 in the space of 1 mm.; they are introduced by intercalation, especially at varices of growth, where commonly they suddenly become smaller and more numerous. In general the lirae do not converge toward the crests of the plications, and they may even cross them at an angle.

Dimensions in mm. of several specimens, all from loc. 434:

| | 1 | 2 | 3 | 4 | 5 |
|---|---|---|---|---|---|
| Length of brachial v. | 23 | 27 | 20 | 22 | 20 |
| Width | 30 | 37 | 31 | 28 | 27 |
| Height of cardinal a. | 12 | — | 13 | 7 | 7 |
| Distance from the beak of first plications | 23+ | 15 | 16 | 18 | 20 |

*Observations:* This species is extremely variable in the strength of plications and the inclination of the cardinal area. Some shells, as numbers 1 and 5 above, show no

*Schellwien, E.: Fauna der Trogkofelschichten, Abh. der K. K. Geol. Reichsanst., bd. 16, heft 1, pp. 20–21, pl. 2, figs. 8–9.

*Yakowlew (Mém. du Com. Géol. Russ., n. s., Liv. 79, 1912, pp. 2–3) places *Meekella striatocostata, M. irregularis, M. procera, M. evanescens, M. depressa* Schellwien, and *M. uncitoides* Tschernyschew in synonymy with *M. eximia* Eichwald. Such an opinion is hardly tenable when we consider the constant differences from one another shown by these species. It may be that some of Schellwien's species are identical (a fact suggested by their general similarity and their having been collected from the same locality, so that their differences may be merely variations of a single species) and perhaps also some of Tschernyschew's, but *M. eximia* can in no case be the same as *M. striatocostata* or *M. irregularis*. True *M. eximia* is apparently closer to the former than to the latter, for it has a much lower cardinal area and stronger plications than Schellwien's species.

plications and would, were it not for the associated individuals, be mistaken for *Orthothetina*. *M. mexicana* is distinguishable from *M. striatocostata* by the flat, strongly inclined cardinal area, and the faintness of the plications. The lirae are less regular than those on the Pennsylvanian species.

*Distribution:* Middle Gym formation (Hueco Mountains) at 421? (one specimen with exceptionally high plications), 425, and 434 (extremely abundant). Upper Gym formation (Hueco Mountains) at 440? and 443. Wolfcamp formation (Sierra Diablo) at 465? (two small faintly plicated shells). Wolfcamp formation (Glass Mountains) at 178?.

### MEEKELLA HESSENSIS King, n. sp.

Pl. V, Figs. 8–9

*Description:* Shell medium-sized, wider than long. Pedicle valve low, with little convexity between the beak and the anterior margin. The cardinal area lies at a little more than right angles to the plane of the shell margin, its width equaling 0.6–0.7 that of the shell and its height twice its width, flat transversely and also longitudinally near the beak. Delthyrium only a fourth the width of the area. Delthyrial supporting plates subparallel; median septum absent. Brachial valve widest a little behind mid-length, anterior margin regularly rounded. Convexity moderate, greatest near center of the shell. Beak very low; umbonal region undefined by greater convexity, the transverse profile showing very little elevation in the median posterior part.

Surface marked by 10–15 fairly regular plications of variable coarseness, originating 5–10 mm. in front of the umbos. The specimen from loc. 122 has exceptionally fine plications. In some specimens the plications bifurcate near the margin. Lirae, numbering about 4 in the space of 1 mm., subequal, narrow; there is some tendency for them to curve upward toward the crests of the plications, and in some places near the margins of the shell the lirae do not parallel the plications but cross them, being directed more laterally. Strong varices of growth cross the surface at intervals.

Dimensions in mm. of several individuals:

| | 1 (122-type) | 2 (222) | 3 (208) |
|---|---|---|---|
| Length | 24 | 21 | 27 |
| Width | 33 | 30 | 38 |
| Thickness | 18 | 18 | — |
| Length of hinge-line | 23 | 17 | 29 |
| Height of cardinal area | 13? | 9? | 15 |

*Observations:* This species is distinguished from *M. striatocostata* by its proportionately greater width, its broad and low cardinal area, and its indistinct dorsal beak. It is similar to *M. skenoides* but has a much broader and lower cardinal area, greater proportional breadth, and, in some specimens, finer plications. It is distinguished from *M. difficilis* by its flat, broad cardinal area, lying near a right angle to the margin, and generally also by its more transverse form.

*Distribution:* Hess formation (Glass Mountains) at 17, 122 (type), 208, 209, 211, and 222. Uncommon.

### MEEKELLA SKENOIDES Girty

Pl. VII, Figs. 6–8

1908 *Meekella skenoides.* Girty, The Guadalupian Fauna, U. S. G. S. Prof. Paper 58, pp. 206–207, pl. 30, figs. 8–9.
Delaware Mountain formation, Comanche Canyon, Glass Mountains (sta. 3763); Delaware Mountain formation (?), southern Delaware Mountains (sta. 3501).

*Description:* Shell medium-sized to large. Pedicle valve high, more or less distorted, subconical, inclined backward so that the area makes slightly more than a right angle with the plane of the valve margins. Longitudinal profile flat or slightly curved. Cardinal area well-defined, high, flat transversely, slightly concave longitudinally; height about equal to or somewhat less than the width. Deltidium medium-sized, at its base equal to a third to a fourth the width of the area; the median part strongly convex. Delthyrial supporting plates long and sub-parallel; median septum absent. Brachial valve transverse, widest at about mid-length, anterior margin regularly rounded. Convexity moderate, greatest near the umbo.

Surface marked by 9–13 sub-angular, nearly equal plications, the number depending on the size of the shell. The shells are unplicated as far as 9 mm. in front of the beaks. Lirae thin, 3–4 in number in the space of 1 mm., subequal or somewhat alternating, and persistent over the whole surface. Toward the anterior margin of large specimens they curve up toward the crests of the plications, with implantation in the sulci corresponding to convergence at the crests of the plications; over the posterior surface, however, the lirae are parallel.

Dimensions in mm. of several specimens:

| | 1 (192) | 2 (192) | 3 (46) | 4 (147) | Girty's type |
|---|---|---|---|---|---|
| Length of pedicle v. | 22 | 25 | 45 | 50 | 11 |
| Length of brachial v. | 18 | 22 | — | — | 11 |
| Width | 21 | 28 | 44 | 52 | 13 |
| Thickness | 14 | 24 | — | — | — |
| Length of hinge-line | 15 | 18 | 23 | 24 | 8 |
| Height of cardinal a. | 12? | 20 | 20? | 35 | 9 |

*Observations:* Typical specimens of this species differ from Girty's type only in their larger size, generally smaller number of plications in specimens of the size of the type, and, in some instances, their lower cardinal areas and broader delthyria. This species is characterized by its medium-sized plications, high, flat, pyramidal pedicle valve, and high, slightly incurved cardinal area. It is unlike any foreign species, none of which is as strongly plicated. It somewhat resembles *M. striatocostata*, but has a much higher cardinal area, and attains a much greater size.

*Distribution:* Word formation (Glass Mountains) at 6, 44–46, 46, 132, 143, 144, T144, T144a, 147, 148, T148, Tx148, 153, 162, 192, 239, 240, T240, 243, T243, 247, 248?, 250, and Ta. Delaware Mountain formation (Sierra Diablo) at 504.

### MEEKELLA STRIATOCOSTATA (Cox)

Pl. VII, Fig. 9

1857 *Plicatula striatacostata.* Cox, Geol. Sur. Ky., vol. 3, p. 568, Pl. 8, fig. 7.

1909 *Meekella striatocostata.* Girty, The Manzano Group, U. S. G. S. Bull. 389, pp. 54–55, pl. 6, fig. 6. See for synonymy.
Abo sandstone, Mesa del Yeso and Abo Canyon.

*Observations:* In the Wolfcamp formation there are a few specimens that cannot, so far as their characters may be seen, be distinguished from the common Pennsylvanian species.

*Distribution:* Wolfcamp formation (Glass Mountains) at 87 low?, 93, 93s, and T93.

#### MEEKELLA sp.

Many fragmentary specimens of *Meekella* could not be classified. Probably most of them belong to species described above. Several, however, may be different. The specimens from the Hess at 223, for example, are pieces of rather large shells marked by very strong rounded plications, except in the umbonal regions, which are marked only by coarse lirae. In profile the pedicle valve is nearly flat. These shells approach closer to *M. pyramidalis* (Newberry) than any others in this fauna. A medium-sized specimen from the Hess at 106 has only 7 high plications, which are as broad as those of *M. grandis* but much higher, and extend nearer the beak. The specimens from the Word at 54 and 265 have very broad, high, sharp plications.

*Distribution:* Wolfcamp formation (Glass Mountains) at T91. Hess formation (Glass Mountains) at 1, 8, 105, 106, 219, and 223. Leonard formation (Glass Mountains) at 26, 39, 82, 83, 125, 126, T226, and Tba. Word formation (Glass Mountains) at 54, 55, 237, and 265. Gym formation (Diablo Plateau) at 449. Hess formation (Sierra Diablo) at Te.

### Genus GEYERELLA Schellwien

*Diagnosis:* Shell sub-conical, distorted. Hinge-line much shorter than the greatest width of the shell. Ventral cardinal area very high and narrow, not sharply set off from the rest of the surface. Delthyrial supporting plates extend to the middle of the shell or beyond, and merge into a septum to form a very elongated chamber below the deltidium. Brachial valve of low convexity. Cardinal process very long and bilobed. Surface marked by radial plications and by fine lirae.

Genotype: *Geyerella gemmellaroi* Schellwien.

*Geologic age:* Permian.

*Observations:* Whether this genus descended from *Orthotetes* to which it is internally similar, or whether the acquisition of a camerate internal character was an independent evolution from some other member of the *Orthothetinae* has been the subject of some debate. Girty* was inclined toward the former interpretation. He stated that the internal structure is precisely that of "the camerate group of *Derbya*, a fact to which Schellwien does not, in my view, give sufficient weight—indeed, I do not know that he calls attention to it at all." In this connection, however, Girty seems to have overlooked the fact that Schellwien* discusses at some length the similarity of *Geyerella "Derbya" correana* Derby which is a typical example of *Orthotetes* (the "camerate group of *Derbya*" of Waagen). Ivor Thomas* pointed out that the much more strongly developed internal plates of *Geyerella* than of *Orthotetes* was an important distinguishing character. To me it seems quite as likely that the arrangement of the internal plates in *Geyerella* is due to the joining together of the anterior part of the delthyrial supporting plates of *Meekella* as to development from *Orthotetes.*

*The Guadalupian Fauna, U. S. G. S. Prof. Paper 58, pp. 203–204.

#### GEYERELLA AMERICANA Girty

Pl. VII, Figs. 10–13

1908 *Geyerella americana.* Girty, The Guadalupian Fauna, U. S. G. S. Prof. Paper 58, p. 204, pl. 11, figs. 2–2b.
Middle Capitan, Guadalupe Point (sta. 2926).

1908? *Orthotetes distortus campanulatus.* Girty, The Guadalupian Fauna, U. S. G. S. Prof. Paper 58, p. 202, pl. 10, figs. 10–10c.
Middle Capitan, Guadalupe Point (sta. 2926).

*Description:* Shell attaining a very large size, with a high, conical distorted pedicle valve and a moderately convex brachial valve. In large shells the valves may meet at the anterior margin in a right angle. Pedicle valve deformed by strong concentric undulations. Hinge-line equal in length to about half the width of the shell. Cardinal area very high, and generally twisted to one side. Axis of the deltidium elevated into a narrow ridge; the rest of this plate is flat and not distinctly set off from the rest of the cardinal area. The delthyrial supporting plates converge to form the median septum at about a third the distance between the deltidium and the floor of the valve. These internal plates are very long, extending from the beak nearly to the anterior margin. Cardinal process long and massive.

Surface marked by 18–26 strong but rounded and very irregular plications, which originate 10–20 mm. in front of the beak and generally bifurcate near the anterior margin. The sides of the shell near the cardinal area are unplicated. Lirae fine and rounded, separated by broader flat striae.

Dimensions in mm. of three specimens:

| | 1(17) | 2(35) | 3(T15) |
|---|---|---|---|
| Length of pedicle valve | 35 | 55 | 47+ |
| Length of brachial valve | 28 | 40 | 30+ |
| Width | 39 | 38 | 44 |
| Length of hinge | 13 | 11 | — |
| Height of cardinal area | 17 | 40 | 40+ |

*Observations:* Girty's specimen, though from a much higher horizon, agrees well with those from the Hess insofar as can be told from such an incomplete type specimen. *Orthotetes distortus campanulatus* is much like *G. americana,* but has plications only at a distance of nearly 20 mm.

*Fauna der Trogkofelschichten, Abh. der K. K. Geol Reichnanst, bd. 16, heft 1, p. 25. In this paper he also gives the name *Geyerella distorta* n. sp. to the same form which he had previously (N. Jahrbuch f. Minéralogie, etc., 1900, hd. 1, p. 13) named *G. alpina.* It would appear that the latter name has priority and should be used.

*British Carboniferous Orthotetinae, Mem. Geol. Surv. Gt. Brit., vol. 1, pt. 2, p. 90–91.

anterior to the beak. Girty suggested that this might warrant the assignment of this variety to *Geyerella*, but himself believed them to be accidental features. The fact that *O. distortus campanulatus* and *G. americana* were collected from the same locality is in favor of their identity. The two south European species of *Geyerella* are not nearly as high and distorted as *G. americana*.

*Distribution:* Hess formation (Glass Mountains) at T15, 17, 35, and 208?. Upper Wolfcamp or lower Hess formation (Glass Mountains) at 85.

### Genus DERBYA Waagen

*Diagnosis:* Shell subplano-convex to biconvex, either regular or more or less distorted in growth. Pedicle valve flat or convex, the cardinal area more or less high and sloping posteriorly from the hinge-line, divided into a primary and secondary area; delthyrium closed to the apex by the deltidium. Within the pedicle valve there is a strong median septum which extends from the inner side of the deltidium to the floor of the valve; it is formed by the fusion of the delthyrial supporting plates. Brachial valve convex, with a narrow cardinal area or no area. Surface unplicated, marked by radial lirae.

Genotype: *Derbya regularis* Waagen.

*Geologic range:* Mississippian to Permian.

### DERBYA BUCHI (d'Orbigny)

Pl. VIII, Figs. 4–6

1842 *Orthis Buchi.* d'Orbigny, Voyage dans l'Amérique méridionale, p. 49.

1842 *Productus Andii.* d'Orbigny, *idem*, p. 54, pl. 5, figs. 1–3.

1914 *Derbya Buchi.* Kozlowski, Brachiopodes du Carb. supér. de Bolivie, Ann. de Pal., vol. 9, pp. 54–59, pl. 8, figs. 1–6.

*Description:* Shell variable in form, large, slightly biconvex, wider than long, with the greatest width at about mid-length, somewhat unsymmetrical. Cardinal angles obtuse and rounded; front margin regularly rounded.

Pedicle valve slightly convex in the umbonal region, and sloping evenly toward the front; beak erect and pointed, not elevated above the cardinal area. Cardinal area 4–6 times as wide as high, inclined 120°–130° from the plane of the valve margin, striated parallel to the base, and with a median triangular part which is likewise vertically striated. The deltidium occupies about a fifth of the area; it is strongly convex and is divided by a longitudinal furrow. Internally there is a strong median septum.

Brachial valve moderately and unevenly convex, with a slight flattening at the sides in the posterior part, in some specimens bearing a slight mesial depression. Hinge-line bordered by a narrow, linear area. Cardinal process broad and thick, projecting at right angles to the plane of the margin, divided by a broad, deep notch into two parts, each of which is grooved at the end. Crural plates strongly diverging.

Surface marked by subequal broad, rounded, crenulated, radial lirae, with intervening flat striae of either greater or less width, about 14 to 20 of each occurring in the space of 10 mm., increasing by intercalation. Growth undulations present but not prominent.

Dimensions in mm. of several specimens:

| | 1 (93—ped. v) | 2 (93—brach. v.) | 3 (441) | 4 (424) |
|---|---|---|---|---|
| Length | 43 | 47 | 36 | 31 |
| Width | 54 | 64 | 50 | 45 |
| Height of card. area | 11 | — | 6 | 6 |

*Observations:* Fragments of certain specimens have been questionably assigned to this species because of their similar ornamentation—strong, regular, closely spaced lirae. *D. cymbula* is distinguished from this species by its smaller width in proportion to length, greater convexity, higher beak, higher cardinal area, and coarser liration. *D. nasuta* is more convex, and has a much higher beak and larger cardinal process. *D. crassa* is smaller, flatter, narrower in proportion to length, and less irregular. *D. crenulata* is much narrower in proportion to length, smaller, and has a higher cardinal area. *D. buchi* is very similar to *D. regularis* Waagen of the Salt Range section, but, as Kozlowski has pointed out, the Indian species is more regular and has the cardinal process enlarged at its end.

*Distribution:* Wolfcamp formation (Glass Mountains) at 91?, 93, 93s, 197?, and 198?. Middle Gym formation (Hueco Mountains) at 424. Upper Gym formation (Hueco Mountains) at 441.

### DERBYA? CRENULATA Girty

1908? *Derbya? crenulata.* Girty, The Guadalupian Fauna, U. S. G. S. Prof. Paper 58, pp. 183–184, pl. 26, figs. 5–5d.

"Delaware Mountain formation," Diablo Mountains (sta. 3764). Delaware Mountain formation, southern Delaware Mountains (stas. 2969 and 3500).

*Description:* Shell medium-sized, transversely ovoid. Pedicle valve rather low. Surface marked near the margins by concentric undulations, otherwise nearly flat, except for the depressed ears. Hinge-line slightly shorter than the greatest width of the shell (in Girty's type the hinge is very long), the greatest width being near mid-length. Cardinal area lying at about 135° to the plane of the margin. Beak pointed and twisted to one side. Brachial valve moderately convex. Ventral median septum large. Surface marked by coarse, strongly elevated, rounded lirae, separated by broad, flat grooves of about the same width. The lirae number 7 to 8 in the space of 5 mm., and are crossed by crenulations, which in the striae appear as fine transverse wrinkles. New lirae are introduced by intercalation, giving the liration an alternating character.

Dimensions in mm. of two specimens:

| | 1 (loc. 248) | Girty's type |
|---|---|---|
| Width | 39 | 30 |
| Length of brachial valve | 27 | 19.5 |
| Width of cardinal area | 35 | 30 |
| Height of cardinal area | — | 5 |

*Observations:* The specimens here referred to this species differ from *D. crassa*, a closely related form, in having slightly stronger, crenulated lirae, and a higher cardinal area. Several differences from Girty's type make absolute identification doubtful. *D.? crenulata* Girty has greatly produced cardinal extremities and apparently more strongly

crenulated lirae. The high cardinal area and coarse lirae are, however, important points of agreement.

*Distribution:* Word formation (Glass Mountains) at 54?, 138?, T144, Tx148, 150, 153?, 241?, 246?, 248 (the best specimen), Ta, and Tx. It is uncommon, no more than one specimen having been collected at most of these localities.

**DERBYA CYMBULA Hall and Clarke?**

Pl. VIII, Fig. 3

1892? *Derbya cymbula.* Hall and Clarke, Pal. New York, vol. 8, pt. 1, p. 348, pl. 11B, figs. 2–3.

*Observations:* This identification is based on a single brachial valve, which agrees in every particular that can be seen with typical *D. cymbula* from the Cisco. It differs from *Orthothetina* sp. in having a coarser ornamentation, from *D. buchi* in its even convexity, less proportionate width, and finer liration, and from *D. crenulata* in its much finer liration. The shell is penetrated by many holes, about 2 mm. long by 0.75 mm. wide, probably made by some boring organism.

*Distribution:* Wolfcamp formation (Glass Mountains) at 169a.

**DERBYA ELEVATA King, n. sp.**

Pl. VIII, Fig. 7

1908 *Derbya* sp. b. Girty, The Guadalupian Fauna, U. S. Geol. Sur. Prof. Paper 58, pp. 185–186, pl. 10. fig. 12.

Top of Capitan, Capitan Peak (sta. 2966). Delaware Mountain formation, southern Delaware Mountains (sta. 2969).

*Description:* This species is represented by two imperfect examples. The first is from a boulder which apparently came from the upper Word. Its shape is conical; cardinal area sharply defined, extending at right angles to the plane of the valve margin, flat in so far as observed. Deltidium fairly wide and slightly convex. Median septum very high and strong. Near the anterior wall it divides into two plates, each diverging laterally to unite with the floor of the valve. Just beneath the deltidium it appears to split and to join the ridges bordering the deltidium on the inside. Surface marked by fine subequal lirae, 12 of which occupy the space of 5 mm.; also by low undulations of growth.

The other example is very doubtfully referred to this species, because of the poor preservation of its surface. The pedicle valve is very high and conical. Cardinal area flat, extending at right angles to the plane of the shell margin, 0.6 of its width occupied by the deltidium. Hinge-line equal to 0.7 the greatest width of the shell. The median septum extends from the middle of the deltidium probably to the anterior wall of the valve; as the shell is not preserved it cannot be told if it actually joins the floor of the valve, or if it is split in the lower part as is the Word specimen. Brachial valve convex, particularly at the beak, showing two strong diverging crural plates but no median septum. Surface marked by fine radial lirae, on this specimen very poorly preserved.

Dimensions in mm. of three specimens:

| | 1 (loc. 265) | 2 (El Tordillo) | 3 (Girty's *D.* sp. b) |
|---|---|---|---|
| Length of pedicle valve | 36+ | 45+ | — |
| Length of brachial valve | 43? | 40 | 10 |
| Width | 56 | 49 | 16 |
| Width of cardinal area | 30 | 35 | 16 |
| Width of deltidium | 15 | 21 | 4.5 |
| Height of cardinal area | 15+ | 38+ | 12 |

*Observations:* The specimen from loc. 265 resembles Girty's typical specimen of *D.* sp. b. in every detail except that it is much larger. The other is more doubtful, but the internal characters and the nature of the ornamentation can hardly be those of anything but a *Derbya.* The two specimens agree in having a high cardinal area, but the Glass Mountains one probably did not have as high an area as that from Mexico.

*Distribution:* Word formation (Glass Mountains) at 265. Limestone of Cerro Caballo (Las Delicias) at section 2, bed 12, between El Tordillo and Cerro Caballo.

**DERBYA NASUTA Girty**

Pl. VIII, Figs. 1–2

1908 *Derbya nasuta.* Girty, The Guadalupian Fauna, U. S. Geol. Surv. Prof. Paper 58, pp. 182–183, pl. 26, figs. 6–6c.

"Delaware Mountain formation," Diablo Mountains (sta. 2764).

*Description:* Shell large, wider than long, growth irregular. Pedicle valve rather high but flat. Cardinal area 2 to 2.6 times as wide as high, lying at nearly right angles to the margin, flat transversely and somewhat concave longitudinally. In one distorted specimen from D3 the cardinal area is bent about 60° by a transverse angulation due to more rapid growth of the pedicle valve on one side; the part of the area in front of this geniculation lies nearly parallel to the plane of the margin on that side of the beak, while the other side is normal. Deltidium about 0.25 as wide at its base as the length of the hinge-line, divided into three parts, of which the median one is strongly convex but with a longitudinal groove, and the lateral ones nearly on a level with the rest of the area. Hinge-line equal in length to 0.7 to 0.9 the width of the shell. Brachial valve evenly convex.

Interior of the pedicle valve with a long, high median septum, at the apex meeting the deltidium, along the inside of which a low ridge continues from the septum to the hinge margin. Delthyrium bordered on the inside by low ridges which converge toward the septum at the apex. Muscle scars deeply impressed. Cardinal process of the brachial valve very large and massive, with strongly diverging apophyses, which receive between them the septum of the pedicle valve, curving upward to become nearly parallel with the plane of the shell margin.

Surface marked by rather thin and high radial lirae separated by striae of nearly equal size; they increase by intercalation and to a slight extent are alternating. The lirae number 10 to 14 in the space of 5 mm., varying on

different specimens. Interruptions in growth are marked by strong concentric undulations.

Dimensions in mm. of several specimens:

| | 1 (loc. 39) | 2 (loc. D3) | 3 (loc. D3) |
|---|---|---|---|
| Length of pedicle valve | 50 | 51 | 23 |
| Width | 80 | 57 | 24 |
| Height of cardinal area | 44 | 47 | 22 |
| Length of hinge-line | 17 | 16 | 11 |
| Width of deltidium | — | 10 | 5 |

*Observations:* This species is easily recognizable by its large size, the fineness of its ornamentation, the large undulations of growth, and the dimensions of its cardinal area. Of the foreign Derbyas, *D. grandis* Waagen is nearest to this species, resembling it closely in size, ornamentation, irregularity of growth, and shape and size of the cardinal process but differing from *D. nasuta* in having the hinge about as long as the greatest width of the shell and a narrower deltidium.

*Distribution:* Hess formation (Glass Mountains) at 4?, 8?, and 98?. Leonard formation (Glass Mountains) at 38?, 39, T41?, 119?, 123?, 124, 128, 174, T226?, and D3. Word formation (Glass Mountains) at 46, 47, 55?, 144, 145, 192, and T239. Leonard formation, transition beds (Shafter) at Th. Hess formation (Sierra Diablo) at 476. Bone Canyon member (Sierra Diablo) at 485. Limestone of Pichagüia (Las Delicias) at Cerro Pichaguïa?. The species is not common at any of these localities.

## Family CHONETIDAE

### Genus CHONETES Fischer von Waldheim

*Diagnosis:* Shell generally small, concavo-convex, subsemicircular in outline, wider than long, the greatest width being along the hinge-line. Pedicle valve convex, generally flattened on the ears, with a narrow cardinal area; delthyrium closed to the apex by the deltidium; lateral margins of the area sharply defined, bearing a single row of hollow, vertical or divergent spines which increase in length toward the cardinal extremities. Internally the muscle scars are generally faint, the surface outside the scars is strongly papillose, the papillae appearing in internal molds as pits. Brachial valve concave, with a cardinal area generally narrower than that of the other valve, and without marginal spines. Internally the shell is papillose just as the pedicle valve. External surface of both valves marked by fine radial costae, by plications, or smooth.

Genotype: *Orthis striatella* Dalman.

*Geologic range:* Upper Ordovician to Permian.

#### CHONETES BIPLICATUS King, n. sp.

Pl. IX, Figs. 12–13

*Description:* Shell medium-sized, wider than long, the cardinal angles being right angles. Pedicle valve rather strongly convex; beak prominent and greatly incurved; cardinal area low, bordered by numerous spines. A narrow, rather shallow median sinus, arising near the beak, lies between two plications, which are bordered laterally by deep sinuses lying parallel to each other rather than radial to the beak. On each side of these sinuses the shell is slightly convex, flattening out toward the hinge. The anterior margin is notched backward at the middle because of the sinus.

Outer surface marked by 4.5 to 5 lirae in the space of 1 mm. They parallel the two median plications, and therefore in the median part of the shell do not increase greatly in number from the beak to the front, but in the sulci bordering these plications they spread out and new lirae are introduced. Dimensions in mm. of one specimen from loc. 476: *length* 10, *width* 15, *convexity* 6.

*Observations:* This shell is distinguishable from other species of *Chonetes* by its narrow, shallow sinus and two high lateral plications lying parallel to each other and separated from the rest of the shell surface by deep sulci. The specimen from loc. 193 is probably immature, though it does not show the biplicate character at a length of 7 mm.

*Distribution:* Upper Wolfcamp formation (Glass Mountains) at 193 and 195. Hess formation (Sierra Diablo) at 476 and 485.

#### CHONETES DELICIASENSIS King, n. sp.

Pl. IX, Figs. 17–20

*Description:* Shell medium-sized, plano-convex, wider than long, length of the hinge less than, or rarely slightly greater than the greatest width of the shell. Pedicle valve strongly convex, without a sinus. Cardinal area relatively narrow, lying generally at about 45° to the valve margin. Spines not well preserved, but probably numbering about 8 on each side of the beak, their bases inclined backward toward the beak at an angle of about 45°, then turning at right angles upon reaching the edge of the area. Muscular areas small and circular. Brachial valve quite flat or very slightly concave near the beak; its cardinal area slightly smaller than that of the other valve and sloping back so as to lie at 120° to 180° to the ventral area. Cardinal process quadrilobate. Surface marked by lirae, of which 3 to 4 occupy the space of 1 mm., increasing in number by bifurcation. The striae are defined by rows of coarse punctae, as in *C. granulifer*.

Dimensions in mm. of 3 specimens:

| | | | |
|---|---|---|---|
| Length | 10 | 14 | 16 |
| Width | 18 | 21 | 22 |

*Observations:* The only other American species having this plano-convex form is *C. platynotus* White, from New Mexico (probably from the Magdalena formation). That shell, however, is not coarsely punctate, a character which allies *C. deliciasensis* more closely with *C. granulifer;* moreover, the New Mexico species is much smaller. The only foreign species at all similar is *C. moelleri* Tschernyschew, but this is too imperfectly known for comparison, the brachial valve not having been found.

*Distribution:* Limestone of Cerro Caballo (Las Delicias) at section 2, beds 12 and 18; section 1, bed 11 (a ventral interior).

#### CHONETES GRANULIFER Owen

Pl. IX, Fig. 14

1855 *Chonetes granulifera.* Owen, Geol. Rept. Minn., Iowa, and Wis., p. 583, pl. 5, fig. 12.

1915 *Chonetes granulifer.* Girty, Fauna of the Wewoka Formation, U. S. Geol. Sur. Bull. 544, pp. 59–62, pl. 7, figs. 12–13b. (See for synonymy.)

*Description:* Shell small or medium-sized, semicircular, with its greatest breadth at the hinge-line, or slightly in front of it. Pedicle valve moderately convex, with a broad, shallow mesial depression (absent in a specimen from loc. T88); beak small, slightly arched and projecting. Cardinal area rather narrow, lying nearly parallel to or at a low angle to the plane of the valves. The cardinal margins bear 4 to 5 spines on each side of the beak. Brachial valve concave, following nearly the curve of the pedicle valve. Cardinal area well developed, being nearly as large as that of the other valve. Surface ornamented by fine lirae.

Dimensions in mm. of two specimens:

| | 1 (loc. 88) | 2 (loc. 88) |
|---|---|---|
| Width | 19 | 13 |
| Length | 11 | 8.5 |

*Observations:* This shell is distinguishable from *C. verneuilianus* by the absence of a strong mesial sinus. The large variety *C. granulifer meekianus,* which is characteristic of the Cisco, is evidently absent from the Permian of the Glass Mountains, where the specimens are more like the typical *C. granulifer.*

*Distribution:* Wolfcamp formation, *Uddenites* zone at 88, 88s, T88, 95, and 195. Wolfcamp formation (Glass Mountains) at 89 and 196.

### CHONETES HESSENSIS King, n. sp.

Pl. IX, Figs. 21–22

*Description:* Shell small, wider than long, subquadrate, concavo-convex, with only a faint trace of a sinus near the margin. Ventral beak high; ears somewhat flattened, but not very sharply defined. Ventral cardinal area slightly larger than the dorsal one, its margin marked by 5 spines on each side of the beak. Cardinal process quadrilobate, the two lateral lobes being much larger than the two inner ones, which are merely narrow ridges lying side by side.

Surface marked by coarse lirae, increasing by bifurcation, about 3 being present in the space of 1 mm. Striae are as broad as or broader than the lirae, marked by irregularly arranged punctae. Concentric growth lines present on some shells.

Length of one shell from loc. 107, 7 mm., width 9 mm.

*Observations:* This species is easily distinguishable by the coarseness of its lirae, which are not, however, so coarse as to place it in the group of the grandicostae, which evidently is not represented in this fauna. The nearest related species is *C. molengraaffi* Broili. The only difference is in the somewhat larger size and coarser liration of the Asiatic specimens. *C. strophomenoides* also is very similar, but has a lower ventral beak and fewer spines along the margin of the area relative to the size of the shell.

*Distribution:* Upper Hess formation (Glass Mountains) at 107.

### CHONETES HILLANUS Girty

Pl. IX, Fig. 6

1859 *Chonetes Flemingi.* Shumard (non Norwood and Pratten), Trans. Acad. Sci. St. Louis, vol. 1, p. 390 (date of volume, 1860) (?).

White (Permian) limestone: Guadalupe Mountains.

1908 *Chonetes hillanus.* Girty, The Guadalupian Fauna, U. S. Geol. Sur. Prof. Paper 58, p. 228, pl. 11, figs. 8–10.

Middle Capitan, Capitan Peak (sta. 2926); dark, limestone, Pine Spring (sta. 2930).

*Description:* Pedicle valve medium-sized, very broad, subtriangular, strongly convex. Sinus distinct but narrow. Hinge-line long, ears sharply pointed, each defined by broad, obscure sulcus, which is more distinct toward the beak than toward the lateral margin. Cardinal area rather high, bordered by about 9 spines on each side of the beak. Median septum extending more than half the length of the shell, bordered on each side by elongated narrow adductor muscles and large, laterally extended diductors, which are strongly defined by encircling grooves. The rest of the internal surface is marked by large granules regularly ranged in radial lines. Length, 10 mm.; width, 22 mm.

*Observations:* This species is easily distinguishable by its convexity, its wide hinge-line, and its low, narrow sinus. *C. subliratus* is larger, less alate, less strongly convex, and has a broad sinus. The surface is probably also quite different.

*Distribution:* Word formation (Glass Mountains) at T144? and 248. Very rare.

### CHONETES MOLENGRAAFFI Broili?

Pl. IX, Fig. 26

1916? *Chonetes Molengraaffi.* Broili, Paläontologie von Timor, Lief 7, no. 12, p. 142, pl. 4, figs. 6–7.

Permian: Bitauni, path between Soefa and Manoemea.

1925? *Chonetes Molengraafii.* Fredericks, Rec. Geol. Com. of the Russian Far East, no. 40, pp. 5–6; pl. 2, figs. 67–68.

Permian: Cape Kalouzin, Ussuriland.

1927? *Chonetes? Molengraaffi.* F. R. C. Reed, Palaeontologia Indica, n. s., vol. 10, mem. 1, pp. 130–131, pl. 14, fig. 16.

Permian: Yellowish mudstone at Ta-li shao, Yunnan.

*Description:* Shell medium-sized, wider than long, subquadrate, cardinal angles rectangular. Pedicle valve strongly convex, especially in the posterior part. Median part depressed. Surface marked by coarse bifurcating lirae, about 2 to 3 of which occur in the space of 1 mm. Striae about as broad as the lirae, marked by rows of pores. Cardinal margin marked by about 7 spines on each side of the beak. Length, 10 mm.; width, 14 mm.

*Observations:* This species differs from *C. hessensis* in its larger size, stronger convexity, and coarser liration. So far as the characters may be seen they agree perfectly with those figured and described for *C. molengraaffi,* but there are no perfect specimens in the collection from Texas, so that absolute identification cannot be made.

*Distribution:* Leonard formation (Finlay Mountains) at 512.

### CHONETES OSTIOLATUS Girty

Pl. IX, Fig. 24

1910 *Chonetes ostiolatus.* Girty, Fauna of the Phosphate Beds of the Park City Formation, U. S. Geol. Sur. Bull. 436, pp. 25–26, pl. 1, figs. 12–14.
Phosphate beds of the Park City formation; Thomas Fork, Wyoming; Montpelier, Idaho; Preuss Range, Idaho.

1927 *Chonetes ostiolatus.* Girty, in Geol., etc. of Part of Southeastern Idaho, U. S. Geol. Sur. Prof. Paper 152, p. 80, pl. 28, figs. 23–25.
Phosphoria formation: southeastern Idaho.

*Description:* Shell large, semicircular, widest at the hinge-line, the cardinal angle being about 80°. Pedicle valve slightly convex, with poorly defined low ears and a low median depression. Beak small and inconspicuous. There is a very shallow median depression. Surface smooth except for fine concentric striae or wrinkles, which are strong on the ears and less distinct over the rest of the surface. The shell is pierced by radially elongate, rather large pores, spaced about 2 mm. apart in the posterior part of the shell but more scattered in the anterior part. Through the outer shell layer fine lirae may be seen. The inside of the pedicle valve is finely papillose. Width 31 mm., length 18 mm.

*Observations:* But one specimen of this species is present in the collections here studied. Its characters fall easily within the limits of *C. ostiolatus* as described by Girty, though it is somewhat more extended at the hinge than in Girty's figured specimen, and in the presence of a median depression it is closer to the variety *impressus.* The distinction of the variety may be of doubtful value, as the distinguishing characters cited by Girty are rather variable. This species is very close to *C. capitolinus* Toula, but figures of that species do not show pores in the shell.

*Distribution:* Word? formation (Shafter) at Tn (no. 16005).

### CHONETES PERMIANUS Shumard

Pl. IX, Figs. 1–4

1859 *Chonetes Permiana.* Shumard, Trans. Acad. Sci. St. Louis, vol. 1, p. 390 (date of volume, 1860).
Permian: conglomerate at mouth of Delaware Creek.

1908 *Chonetes permianus.* Girty, The Guadalupian Fauna, U. S. Geol. Sur. Prof. Paper 58, pp. 226–227, pl. 20, figs. 1–3a; pl. 29, figs. 1–2.
Dark limestone, Pine Spring (sta. 2930) and east of Guadalupe Point (sta. 3762b). Delaware Mountain formation, southern Delaware Mountains (stas. 2936 and 2969).

1908 *Chonetes* sp. Girty, *idem,* pp. 229–230.
Delaware Mountain formation (stas. 2903 and 2931); basal black limestone, Guadalupe Point (sta. 2920).

*Description:* Shell small to medium-sized, subsemicircular; hinge-line varying from slightly shorter to slightly greater than the greatest width of the shell; the cardinal angle being generally a right angle. Pedicle valve moderately convex, without a mesial sinus or with a faint depression; beak somewhat higher than the adjoining parts of the shell. Ears slightly deprest, but not well-defined. Cardinal margin with 6 spines on each side of the beak. Hinge-teeth small but distinct, not supported by dental plates. The median septum extends about half the length of the valve. Diductor muscle scars long and oval, lying beside the median septum. The rest of the interior surface is covered by coarse granulations.

Brachial valve flat in the posterior part but concave near the anterior margin. Cardinal process quadrilobate, being divided in the middle by a deep, narrow groove, and on each side by broader furrows. Crural plates short, diverging at about 45° from the hinge, giving rise to two linear brachial ridges, diverging at 60° from the hinge. Muscular area divided by a short, sharp median septum. Each antero-lateral part of the interior is raised into a curved ridge. Internal surface marked by large papillae, directed toward the lateral margins.

Surface smooth except for fine striae of growth. The inner shell layer is finely striated and punctate as in *C. subliratus.*

Dimensions in mm. of several specimens:

| | 1 (loc. 6) | 2 (loc. 26s) | 3 (loc. 8–14) | 4 (loc. 31) | 5 (loc. 128) |
|---|---|---|---|---|---|
| Width | 14 | 20 | 20 | 12 | 13 |
| Length | 8 | 12 | 12 | 7 | 9 |

*Observations:* According to Girty, *C. permianus* Shumard has 3 rather than 6 spines on each side of the beak. Girty distinguished the type having a greater number of spines as *Chonetes* sp., but this feature is not here regarded as important enough to warrant distinguishing that type. Girty supposed that Shumard in his original description implied that there were 5 or 6 spines on the entire cardinal margin, but the fact that an odd number (5) is mentioned would rather suggest that Shumard meant 5 or 6 on each side. If so, Shumard's description fits best the specimens with the greater number of spines.

*C. geinitzianus* Waagen of the Pennsylvanian is broader than this species and has a well-marked sinus. *C. ostiolatus* Girty differs in its large size, low convexity, and the presence of pores piercing the shell. *C. ambiensis* Waagen, of the lower Productus limestone, appears to differ from *C. permianus* in having a higher ventral cardinal area, resulting in a greater angle between the cardinal margins on opposite sides of the beak. *C. obtusa* Schellwien is very similar but has a slight sinus. *C.* cf. *geinitzianus* Reed, from the Permian of Yun-nan, appears to be close to *C. permianus,* for it evidently has no mesial sinus in the pedicle valve. It appears, however, to be marked by stronger concentric growth lines. The description of *C. laevis* Davidson from Kashmir shows no point of difference from this species, but until it is better known any comparison would be unsatisfactory. *C. alatus* Stuckenberg has a much longer hinge. *C. rotundata* Toula has large punctures like those of *C. ostiolatus. C. amazonicus* Derby is of this type but has a nearly flat pedicle valve.

*Distribution:* Hess formation (Glass Mountains) at 207?. Leonard formation (Glass Mountains) at 8–14, D13, 15, T15, 26s, 31, 128, and 301. Word formation

(Glass Mountains) at 6, T144, 148, and T251. Upper Wolfcamp formation (Sierra Diablo) at 456?. Bone Canyon member (Sierra Diablo) at 483 and 484. Bone Canyon member (Delaware Mountains) at 490. Victorio Peak member (Sierra Diablo) at 495. Leonard formation (Baylor Mountains) at 495. Not abundant at any locality.

**CHONETES QUADRATUS King, n. sp.***

Pl. IX, Figs. 7–11

*Description:* Shell small, subquadrate, slightly wider than long; cardinal angles approximately 90° in most specimens, but in some a little less. Sinus shallow and rounded, but narrow. Visceral slopes very steep, defining sharply the low ears. Spines probably about 6 in number on each side of the beak. Surface smooth, with tubules piercing it as in *C. subliratus;* inner shell layer finely striated.

Dimensions in mm. of several specimens:

| | 1 (loc.145-type) | 2 (loc.246) | 3 (loc.247) |
|---|---|---|---|
| Length | 12 | 10 | 9 |
| Width | 14 | 12 | 12 |

There is much variation shown by some shells assigned to this species. One from loc. 246 has no fold or sinus. One from loc. 256 has a narrow and deep sinus.

*Observations:* This species probably is identical with a group of shells from the Kaibab limestone, probably the same as the ones referred to by Girty as *C. hillanus*, in Kaibab fossil lists. The Kaibab type is characterized, like the Glass Mountain species, by having quadrate ears and a narrow, shallow sinus. *C. quadratus* is closely related to *C. subliratus*. It differs in having a smaller size, quadrate outline, sharply defined ears, and a shallower ventral sinus. *C. hillanus* has, among other differences, much more extended cardinal extremities.

*Distribution:* Upper Word formation (Glass Mountains) at 145, T148, 192, 246, 247, and 256. Kaibab limestone of the Grand Canyon region.

**CHONETES SPINOLIRATUS King, n. sp.***

Pl. IX, Figs. 15–16

*Description:* Shell very small, wider than long, ears quadrate. Pedicle valve strongly convex, beak prominent and incurved. Median sinus narrow and shallow. Ears small and not well-defined. Cardinal area fairly large, but smaller than that of the brachial valve. Cardinal margin with 3 spines on each side of the beak. Brachial valve concave, but with a very faint median ridge. Cardinal process quadrified, with two median lobes almost united. Surface of both valves ornamented with moderately strong lirae, numbering 5 to 6 in the space of 1 mm.; each lira divides once between the beak and the margin. The lirae of both valves appear to be hollow and open out at regular intervals in forward-projecting spine bases about 1 mm. or less apart. Dimensions in mm. of the type specimen: length 5, width 6, convexity 2.

*Observations:* The small size, narrow median sinus, strong lirae, and the possession of spines on the lirae serve to distinguish this species from all the others in this fauna. The specimen from loc. 92 has somewhat stronger lirae than the type, and spines cannot be seen on the lirae. There is, however, no other apparent difference, and the absence of spines is probably due to difference in the preservation.

*Distribution:* Wolfcamp formation (Glass Mountains) at 91a (type), 92, and 175a.

**CHONETES SPINOLIRATUS DIABLOENSIS King, n. var.**

Pl. IX, Fig. 23

*Description:* This variety differs from typical *C. spinoliratus* in having much stronger lirae, their number being 4 or less in the space of 1 mm. The spines arising from them are larger and fewer. It is also somewhat more strongly convex. Length 4.5 mm.; width, 6.5 mm.; convexity, 2 mm.

*Distribution:* Hess formation (Sierra Diablo) at 476 and 479. Rare.

**CHONETES SUBLIRATUS Girty**

Pl. IX, Fig. 25; Pl. X, Figs. 1–7

1908 *Chonetes subliratus.* Girty, The Guadalupian Fauna, U. S. Geol. Sur. Prof. Paper 58, pp. 228–229, pl. 20, figs. 4–7.

Dark limestone, Pine Spring (sta. 2930); Delaware Mountain formation, Guadalupe Point (sta. 2919); ?Basal Capitan, hill southwest of Guadalupe Point (sta. 2906).

*Description:* Shell of medium to large size, wider than long, alate, strongly concavo-convex. Sinus of the pedicle valve strong, the shell rising in two humps on each side of the sinus and descending abruptly toward the wings, which are depressed and convex. Cardinal area broad, lying nearly in the plane of the valve margin. Spines about 7 in number on each side of the beak, their bases inclined backward toward the beak at an angle of about 45°, then turning at right angles upon reaching the edge of the area so that the spines extend back away from the beak. Delthyrium (as shown in a specimen from loc. 46) bordered by strong hinge-teeth, which are not supported by septa. Median septum low at the posterior end, but extending into a high, irregular, pustulose ridge which reaches three-fourths the length of the valve. Muscle scars irregularly striated. Lateral interior part of the shell marked by coarse granules arranged in distinct radial rows.

Brachial valve corresponding closely with the pedicle valve in curvature. Dorsal area about half as high as the ventral. Cardinal process quadrilobate on the posterior face, uniting forward with crural plates. Muscle scars separated by a median ridge, enlarging forward, and by two lateral ridges lying at a low angle to the septum, and each ending in a low protuberance. Anterior part of the inside of the valve raised into a marginal ridge, which is undeveloped along the median line. Interior surface of both valves strongly papillose.

*This species may be identical with *C. quadratus* Girty, Jour. Wash. Acad. Sci., Vol. 19, 1929, p. 707.

*Probably a synonym of *C. victoriansis* Girty, Jour. Wash. Acad. Sci. Vol. 19, 1929, p. 410.

Where perfectly preserved, the surface is essentially smooth, being marked only by very fine growth lines and by regularly spaced round holes that run obliquely out through the shell and may represent the bases of hollow spines. Where the outer shell layer has been removed by weathering, as is the case in most of the specimens from the Leonard and all of those studied by Girty, the underlying layer with radial striae is exposed; these striae are defined by rows of punctae, about 6 of which are present in the space of 1 mm. Only about a fourth of the specimens from loc. 123 show the outer smooth layer, while practically all those from the Word do. In some shells the outer layer has been partly worn through or partly destroyed during silicification, so that the striae may be seen through it.

Dimensions in mm. of several specimens:

| | 1 (loc. 250) | 2 (loc. 123) | 3 (loc. 123) | 4 (loc. 144) | 5 (loc. 253) |
|---|---|---|---|---|---|
| Length | 15 | 10 | 15 | 33 | 22 |
| Width | 26 | 14 | 26 | 20 | 14 |

*Observations:* The only apparent differences between these specimens and *C. hillanus* Girty, from the Capitan, are that these are more strongly convex, more subquadrate, and probably have a stronger sinus. If the surface of *C. hillanus* as described were perfectly preserved, then this shell differs also in having a smooth outer layer pierced by pores; as indicated above, however, it is doubtful if Girty had perfect specimens on which to base his species.

*C. subliratus* is related to *C. ostiolatus* but there is a well developed sinus, while there is none in *C. ostiolatus*. *C. subliratus* may be confused with large specimens of *C. verneuilianus* if the outer layer has been destroyed, so that the punctate and striate interior is exposed, but the earlier species has a deeper and narrower sinus, and its punctae are not arranged in lines along the striae as in *C. subliratus*. It is also a much smaller shell than the Permian species.

The Salt Range species *C. morahensis*, *C. avicula*, *C. trapezoidalis*, and *C. bipartita* are much like *C. subliratus*, but do not have at all the same configuration of the sinus as the American species. They do, however, show traces of a similar perforation of the smooth shell by pores. *C. obtusa* Schellwien, from the Auernig formation, differs from *C. subliratus* in being less convex and having a much weaker sinus.

*Distribution:* Hess? formation (Glass Mountains) at 207. Leonard formation (Glass Mountains) at 3, T9, 12, T15a, 16, 20, 31, T119, 129, 123 (extremely abundant), 124, 227, 231?, D7, and D1. Leonard? formation (Glass Mountains) at 78, 82, T82a, and 83. Word formation (Glass Mountains) at 44, 46, 54?, 139, T140, T141, 144, 145, T148, Tx148, 238, T238, 239, T239, 241, 243, 250, 253, 265, and Tx. Vidrio member (Glass Mountains) at 49. Lower Hess formation (Sierra Diablo) at 471. Hess formation (Sierra Diablo) at 476 and 479. Bone Canyon member (Sierra Diablo) at 484. Victorio Peak member ((Sierra Diablo) at 499. Leonard formation (Finlay Mountains) at 512 and 513.

**CHONETES VERNEUILIANUS Norwood and Pratten?***

Pl. IX, Fig. 5

1854 *Chonetes verneuilianus.* Norwood and Pratten, Jour. Acad. Nat. Sci. Phila., vol. 3, p. 26, pl. 2, figs. 6–6c.

*Description:* Shell small, wider than long, widest at the hinge-line. Pedicle valve convex, with a deep rounded median sinus, originating near the beak and widening slightly toward the anterior margin; ears more or less angular, flattened; beak rather prominent and incurved. Cardinal areas moderately developed, largest on the pedicle valve. Cardinal margin with 4 or 5 oblique spines on each side of the beak. Brachial valve rather deeply concave. Surface of both valves ornamented by fine lirae.

Dimensions in mm. of three specimens:

| | 1 (loc. 94s) | 2 (loc. 75) | 3 (loc. 94) |
|---|---|---|---|
| Length | 9 | 10 | 7 |
| Width | 14 | 16 | 10 |

The specimen from loc. 94 is exceptional in having a shorter hinge-line than the greatest width of the shell.

*Observations:* This species is distinguishable by its convex pedicle valve and its deep median sinus. The specimens from loc. 94s* were identified by Schuchert as *Chonetes subliratus*. That species is very similar to *C. verneuilianus*, but is in general less strongly bilobate, larger, and has an outer smooth layer perforated by pores.

A bilobate Chonetes that may belong to this species was collected from section 1, bed 11, near Las Delicias, Coahuila. It has a narrower and deeper sinus than the Wolfcamp specimens assigned to this species.

*Distribution:* Wolfcamp formation, *Uddenites* zone (Glass Mountains) T88, 94, and 94s. Wolfcamp formation (Glass Mountains) at 75, 87?, and 195.

**Genus ISOGRAMMA Meek and Worthen**

*Diagnosis:* Shells short, transversely elongate, hinge-line straight, generally making the greatest width of the shell. Pedicle valve slightly convex, with traces of a broad, obscure median sinus; brachial valve flat. Shell very thin and fragile, with a coarsely prismatic cellular structure. Surface covered with numerous regular and continuous concentric rounded ridges which are separated by furrows of equal width. In the pedicle valve there is no cardinal area, teeth exceedingly small. A pair of divergent, elevated ridges originate within the apex of the valve and extend for a third or even half the length of the shell, inclosing a thickened area or platform which terminates abruptly in a transverse anterior margin. This platform bears the adductor and diductor muscles, and probably rests on the bottom of the valve and is not vaulted. In the brachial valve there is a prominent cardinal process, from the base of which diverge two lateral ridges or socket-walls, lying just within the hinge-line; behind them are linear depressions or dental sockets.

*See *C. consanguineus* Girty, Jour. Wash. Acad. Sci., Vol. 19, 1929, p. 409.

*See Am. Jour. Sci., vol. 14, 1927, p. 392.

There is a low median ridge extending from the base of the cardinal process into the pallial region.

Genotype: *Chonetes? millepunctata* Meek and Worthen.

*Geologic range:* Pennsylvanian and lower Permian.

**ISOGRAMMA MILLEPUNCTATA (Meek and Worthen)**

Pl. X, Fig. 8

1870 *Chonetes? millepunctatus.* Meek and Worthen, Proc. Acad. Nat. Sci. Phil., p. 35.

1873 *Isogramma millepunctata.* Meek and Worthen, Geol. Surv. Ill., vol. 5, p. 568.

*Distribution:* Wolfcamp formation, *Uddenites* zone (Glass Mountains) at 203. Upper Wolfcamp formation (Sierra Diablo) at 456.

## Family PRODUCTIDAE

In the present paper an attempt is made to classify the species of this family according to the work of Thomas, Chao, Fredericks, and Muir-Wood.* As there is a great deal of conflict in the ideas of these different writers, it has been necessary to correlate their work. The classification of Chao is accepted in the main, and the following artificial key to the genera of Productidae is taken from his second work, but with some additions and changes. Genera recognized in the species described in the present paper are starred.

I Cardinal area present.

A Ventral median septum absent.

1 Shell small; costae discontinuous, nodose; cardinal teeth present. . . . *Productella* Hall.

2 Shell large, broad; striate, cardinal teeth absent. . . . *Kansuella* Chao.

3 Shell semireticulate. . . . *Sinuatella* Muir-Wood.

B Well-developed ventral median septum present.

1 Shell pustulose, cardinal area low, ventral beak incurved. . . . *Tschernyschewia* Stoyano.

2 Shell pustulose, cardinal area very high, ventral beak erect. . . . *Scacchinella* Gemmellaro.*

II Cardinal area absent.

A Visceral region of the interior of the brachial valve not defined by marginal elevations.

1 Surface pustulose.

a Surface with well-marked wrinkles bearing one or more rows of pustules. Brachial ridges diverge from the median ridge of the brachial valve some distance anterior to the cardinal process. . . . *Pustula* Thomas.*

b Surface with well-marked wrinkles bearing a row of pustules. Brachial ridges diverge from the base of the cardinal process, each pair of ridges having the shape of a ladle. . . . *Overtonia* Thomas.*

c Surface with elongate pustules in early stages, replaced in later stages by concentric bands bearing pustules of two sizes (as in *P. nebraskensis*). . . . *Juresania* Fredericks

d Surface with quincunxially arranged pustules. . . . *Waagenoconcha* Chao.*

2 Surface pustulo-plicate.

a Surface marked by tubercles in the early stages, giving rise to plications in later stages. . . . *Avonia* Thomas.*

b Surface marked by elongate pustules arranged in lines to form costae. . . . *Buxtonia* Thomas.*

3 Surface plicate and generally semireticulate. . . . *Productus* Sowerby.*

4 Surface lirate.

a Hinge-line wide, visceral cavity deep.

(1) Anterior margin evenly convex. . . . *Linoproductus* Chao.*

(2) Anterior margin of the pedicle valve elongated into a narrow proboscis. . . . *Proboscidella* Oehlert.

b Hinge-line short, visceral cavity narrow. . . . *Striatifera* Chao.*

5 Surface smooth.

a Wrinkled in early stages. . . . *Plicatifera* Chao.

b Unwrinkled. . . . *Horridonia* Chao.

B Visceral region of the interior of the brachial valve defined by marginal elevations.

1 Inner surface of the brachial valve outgrown at its geniculation to form a raised platform. . . . *Diaphragmus* Girty.

2 Inner surface of the visceral area of the brachial valve surrounded by a marginal ridge. . . . *Marginifera* Waagen.

**Genus PRODUCTUS Sowerby**

*Diagnosis:* Shells varying in size from small to very large, deeply concavo-convex in form, brachial valve geniculate, the anterior part of the shell being produced in a

*Thomas, Ivor: The British Carboniferous Producti. I.—Genera *Pustula* and *Overtonia*. Mem. Geol. Sur. of Great Britain, Palaeontology, vol. 1, pt. 4, pp. 197–366. 1914.

Chao, Y. T.: Productidae of China. Part I: Producti. Pal. Sinica, ser. B, vol. 5, fasc. 2. 1927.

Chao, Y. T.: Productidae of China. Part II: Chonetinae, Productinae and Richthofeninae. Pal. Sinica, ser. B, vol. 5, fasc. 3. 1928.

Fredericks, Georges: Communication for the Classification of the Genus Productus. Bull. du Com. Géol. (Russia), t. 46, no. 7, pp. 773–792. 1928.

Muir-Wood, Helen M.: The British Carboniferous Producti. II.—*Productus (sensu stricto); semireticulatus* and *longispinus* groups. Mem. Geol. Sur. of Great Britain, Palaeontology, vol. 3, pt. 1, pp. 1–217. 1928.

long, narrow trail. Hinge-line straight, equaling the greatest width of the shell; ears generally extended. Pedicle valve without cardinal area or hinge-teeth, the two valves being held together at the cardinal margin by the strongly incurved beak of the pedicle valve. Cardinal area, dental sockets, and socket plates absent or very rudimentary, the cardinal process large and strong, extending far beyond the hinge-line. Visceral region of the brachial valve not defined by marginal ridges. Surface of both valves marked by radiating costae, and, in typical forms of the genus, by strong concentric wrinkles in the visceral region. The *P. guadalupensis* group is unwrinkled. Hollow spines are present over the shell surface, generally most conspicuously developed on the ears.

Genotype: Here regarded as *Anomites semireticulatus* Martin.

*Geologic range:* Mississippian to Permian.

*Observations:* While it is known that the true genotype of *Productus, Anomites productus* Martin, is actually a *Diaphragmus,* it is preferred to use the name here in the old sense, as if *A. semireticulatus* were the type.*

**PRODUCTUS ARCTICUS Whitfield**

Pl. X, Figs. 16–17

1878 *Productus semireticulatus.* Etheridge, Quart. Jour. Geol. Soc., vol. 34, pp. 629–630.
Fielden Isthmus, Ellesmere Island.

1908 *Productus semireticulatus arctica.* Whitfield, Bull. Amer. Mus. Nat. His., vol. 24, pp. 54–55, pl. 1, no. 2, pl. 2, figs. 8–10.
Cape Sheridan, Ellesmere Island.

1916 *Productus inflatus.* Tschernyschew and Stepanow, Rept. of Second Norw. Arctic Exp. in "Fram," vol. 34, pp. 30–32, pl. 5.
Store Björnekap, Ellesmere Island; Svarte Veg, Heiberg Island.

?1917 *Productus inflatus.* Grönwall, Marine Carboniferous of Northeast Greenland, Medd. om. Grønland, vol. 43, pp. 578–579, pl. 29, figs. 17–19.
Cape Jungerson and Eskimo Naze, Northeast Greenland.

*Description:* Shell rather large, with the hinge-line equaling the greatest width of the shell. Pedicle valve strongly convex longitudinally and abruptly geniculate at about two-fifths the length from the beak to the anterior margin, the curvature in front of and behind the geniculation being relatively low, though the beak is strongly incurved. Visceral region narrow and pointed, the umbonal angle being about 75°; lateral slopes of this region steep, setting off the ears sharply. Median sinus narrow but distinct, originating near the beak and maintaining about the same size from there to the front margin. Brachial valve nearly flat in the posterior half, then sharply geniculated. Interior of the pedicle valve strongly pustulose, especially in the anterior and lateral parts. Diductor muscle scars small but deeply impressed, and coarsely and irregularly striate. The adductor scars lie on the ridge between the diductors, and are divided by a groove.

Surface marked by medium-sized, even ribs, numbering 7 to 9 in the space of 10 mm., which remain of practically constant number in the median part but divide by bifurcation on the lateral slopes. Posterior to the geniculation the surface is crossed by low wrinkles, which are fine and strong on the wings. The part of the shell anterior to the reticulate region bears unevenly distributed spine bases, which are very small for so large a shell. Two rows of spines lie along the lateral slopes of the visceral region.

Dimensions in mm. of several specimens:

| | 1 (loc. 247) | 2 (loc. 246) |
|---|---|---|
| Length | 36 | 36 |
| Width | 35 | 37 |
| Thickness | 20 | 21 |

*Observations:* This species is distinguished by the geniculation and the low convexity in front of and behind it, by the incurved beak, by the pointed visceral region, by the narrow median sinus, by the low wrinkling of the posterior part, and by the small spines distributed over its surface. The Texas specimens agree perfectly with the figures and brief description of Whitfield. Tschernyschew and Stepanow figure as *P. inflatus* specimens that are quite like *P. arcticus,* but seemingly differ in several respects from the *"P. inflatus"* of the Uralian.

*Distribution:* Upper Word formation (Glass Mountains) at 143, 152, 246 (common), and 247 (common).

**PRODUCTUS CAPITANENSIS Girty**

Pl. X, Fig. 15

1858 *Productus semireticulatus.* Shumard, Trans. Acad. Sci. St. Louis, vol. 1, p. 292 (date of volume, 1860).
White (Permian) limestone: Guadalupe Mountains.

1859 *Productus semireticulatus* var. *antiquatus.* Shumard, *idem,* p. 389.
White (Permian) limestone: Guadalupe Mountains.

1908 *Productus semireticulatus* var. *capitanensis.* Girty, The Guadalupian Fauna, U. S. G. S. Prof. Paper 58, pp. 254–255, pl. 12, figs. 1–3b; pl. 20, figs. 8–8a(?).
Top of Capitan, Capitan Peak (sta. 3762); middle Capitan, Capitan Peak (sta. 2926); basal Capitan, hill southwest of Guadalupe Point (sta. 2906); (?) dark limestone, Pine Spring (sta. 2930).

*Description:* Shell large. Pedicle valve strongly convex, beak low. Sinus strong, narrow, extending nearly to the beak. Surface marked by fine costae, about 5 in number in 5 mm., and in the posterior part by concentric wrinkles, which produce nodes where they cross the ribs. Width, 37 mm.; length, 27 mm. plus.

*Observations:* This species is distinguished by its fine costae and strong narrow sinus. The specimens from the upper dark limestone of the Delaware Mountain formation identified by Girty as *P. capitanensis* are more coarsely costate, in which respect they are more similar to *P. arcticus,* which occurs at the equivalent horizon in the Glass Mountains.

*Editor's footnote: Miss Muir-Wood has recently proposed the name *Dictyclostus* for the "semireticulatus" group. Ann. and Mag. Nat. Hist., Vol. V, 1930, p. 103.

*Distribution:* Gilliam member (Glass Mountains) at 179 (one incomplete specimen).

**PRODUCTUS COMANCHEANUS Girty**

Pl. XI, Fig. 1

1908 *Productus guadalupensis* var. *comancheanus.* Girty, The Guadalupian Fauna, U. S. G. S. Prof. Paper 58, pp. 261–262, pl. 31, figs. 5–5b.
Delaware Mountain formation, "Comanche Canyon," Glass Mountains.

*Description:* Shell medium-sized, very wide in proportion to length. Pedicle valve strongly convex. Beak rather prominent. Ears long and depressed. Sinus deep and narrow, giving the shell a bilobate form. Surface marked by coarse ribs, numbering 4 or less in the space of 5 mm. No concentric wrinkling is visible. Length, 13 mm.; width, 32 mm. Length of Girty's type, 16 mm., width 38 mm.

*Observations:* In both Girty's collection and my own there is but a single specimen of this species. Girty considered it to be very close to *P. guadalupensis,* but it appears to me to be quite distinct, being much shorter in proportion to width, and having larger ribs which probably are not at all crossed by concentric undulations.

*Distribution:* Word formation (Glass Mountains) at 144.

**PRODUCTUS DARTONI King, n. sp.**

Pl. X, Figs. 12–13

*Description:* Shell small, transversely subquadrate, with the greatest width at the hinge. Pedicle valve strongly convex, geniculated at about two-fifths the length from beak to front. Umbonal region highly inflated; beak incurved over the cardinal margin. Ears very sharply defined, the lateral slopes of the visceral area being very steep. Sinus broad and shallow. Brachial valve flattened in the visceral area, then abruptly geniculated.

Surface of the pedicle valve marked by low, rounded costae, numbering about 6 in the space of 5 mm., separated by much narrower striae. The visceral area is crossed by low concentric wrinkles (hardly distinguishable in the specimen from loc. 460). Small spines are unevenly distributed over the surface, and an oblique row of prominent spines lies along the inner side of each wing.

Dimensions in mm. of several specimens:

| | 1 (loc. 460) | 2 (loc. 399) | 3 (loc. 385) |
|---|---|---|---|
| Length | 19 | 22 | 19 |
| Width | 23 | 26 | 25 |
| Convexity | 13 | 12 | 11 |

*Observations:* This species is distinguished from other similar ones in the fauna by its low, fine even ribs, well-marked sinus, geniculated profile, and prominent ventral umbo. The specimens from the Victorio Peak member have a short snout-like emargination of the anterior margin, a fact suggesting relation to *"Proboscidella" genuina* Kutorga, which is externally somewhat similar. However, as Fredericks has pointed out, the presence of this anterior median extension is probably not of great significance.

*Distribution:* Lower Gym formation (Hueco Mountains) at 377b, 385, and 399. Upper Gym formation (Hueco Mountains) at 440. Upper Wolfcamp formation (Sierra Diablo) at 459 and 460 (type). Hess formation (Sierra Diablo) at 479.

**PRODUCTUS DARTONI SULLIVANENSIS King, n. var.**

Pl. X, Fig. 14

*Description:* This variety differs from typical *P. dartoni* in having a low, small beak instead of an inflated umbonal region, and somewhat coarser costae. Dimensions of a specimen from loc. 46: length 22 mm., width 25 mm.

*Distribution:* Word formation (Glass Mountains) at 46, 132, and T239. Word formation?, thin-bedded zone (Shafter) at Tj. Victorio Peak member (Sierra Diablo) at 493.

**PRODUCTUS GUADALUPENSIS Girty**

Pl. X, Figs. 9–11

1908 *Productus guadalupensis.* Girty, The Guadalupian Fauna, U. S. G. S. Prof. Paper 58, p. 261, pl. 22, figs. 1–2.
Delaware Mountain formation, Guadalupe Point (stas. 2919 and 2931).

*Description:* Shell medium-sized, inflated, wider than long. Pedicle valve so enrolled that its profile is semicircular. Beak low, bluntly pointed, not extending beyond the hinge-line; umbonal region depressed, showing in profile a very low convexity. Ears low, convex, greatly extended. Median sinus strong, narrow, originating near the beak. On one shell from loc. 247 the sinus is broad and deep, and the shell as a result is strongly bilobed. Brachial valve deeply concave, with a low median fold. Surface marked by radial lirae, numbering 6 to 8 in the space of 5 mm., and by a few widely scattered large spines in the marginal region. There are no concentric ribs in the umbonal region, but a few faint undulations of growth are present on some specimens in the mid-part of the shell.

Dimensions in mm. of several specimens:

| | 1 (loc. 247) | 2 (loc. 247) | 3 (loc. 247) | 4 (loc. 246) |
|---|---|---|---|---|
| Length | 19 | 22 | 18 | 21 |
| Width | 36+ | 44+ | 27 | 40 |

*Observations:* This species is distinguishable from related forms by its rather fine costae, absence of wrinkles, few but large spines, prominent sinus, low umbonal region, and extended ears. The specimens from the Glass Mountains have coarser ornamentation and a flatter ventral umbonal region than Girty's types. The differences give the type an aspect close to that of *P. schucherti,* but the latter species has very significant differences—pronounced wrinkling of the posterior part, large spines on the ears, and generally larger costae, and, moreover, it occurs at a lower stratigraphic level. *P. guadalupensis* differs from *P. multistriatus* in having smaller size, coarser costae, and stronger enrollment. It resembles *P. geniculatus* Girty and *P. mammatus* Keyserling in many respects, but is larger and has a deeper sinus and longer ears.

*Distribution:* Word formation (Glass Mountains) at 148, T148, Tx148, 155, 246 (common), 247 (abundant), and Tc.

**PRODUCTUS HESSENSIS King, n. sp.**

Pl. XI, Figs. 3–6

*Description:* Shell rather large, transversely subquadrate. Pedicle valve slightly convex in the posterior third, the beak overarching the hinge-line but little; somewhat posterior to mid-length there is generally more or less of an increase in curvature, and the curvature decreases gradually to the front margin. Ears broad, rather flat, and not sharply defined. Median region deprest to form a deep, narrow sinus. Brachial valve strongly geniculate, being nearly flat in the visceral region and having a long trail; along the median line there is a low elevation corresponding to the ventral sinus. Visceral regions of both valves marked by fine concentric wrinkles. Radial costae even and very fine, numbering 8 to 11 in 10 mm. Spines rather fine and on most specimens rather widely spaced, but some well-preserved specimens (loc. 510 and Tzz, aa) show rather thickly spaced spines. Spines on the ears not large or abundant.

Dimensions in mm. of several specimens:

| | 1 (loc. 106) | 2 (loc. Tzz, aa) | 3 (loc. Tzz, aa) |
|---|---|---|---|
| Length | 34 | 42 | 36 |
| Width | 45 | 46 | 34 |
| Convexity | 16 | 25 | 19 |

*Observations:* This species differs from the common Pennsylvanian types designated as *P. semireticulatus* in having a low beak, more or less geniculation, narrow median sinus, and fine costation. *P. capitanensis* is somewhat similar, but has a narrower median sinus and a more strongly geniculate profile. The specimens of *P. hessensis* from northern Trans-Pecos Texas shows less geniculation and less distinct median sinus than those from the Glass Mountains.

*Distribution:* Upper Hess formation (Glass Mountains) at 106 (type), 107, T107, and 222. Lower Gym formation (Wylie Mountains) at 507 and Tzz, aa. Lower Gym formation (Carrizo Mountains) at 510. Lower Gym formation (Hueco Mountains) at 405 and 408.

**PRODUCTUS HUECOENSIS King, n. sp.**

Pl. XI, Figs. 7–8

*Description:* Shell large, very strongly convex, transversely subquadrate. Pedicle valve evenly convex (though the specimen from loc. 382 is geniculated at about mid-length), the longitudinal curvature decreasing gradually forward. Median region flattened or depressed to form a broad, shallow sinus. Ears well-defined, broad, and depressed. Brachial valve concave in its posterior two-thirds, then very strongly geniculate. Surface ornamented with radial ribs to the number of about 10 in the space of 10 mm., of rather even constant size except immediately anterior to a bifurcation. The visceral region is crossed by low or indistinct wrinkles, which are stronger on the lateral slopes and the ears. The part of the shell anterior to the wrinkled region has many relatively small spines rather evenly distributed. A row of large spines lies in an oblique row between each ear and the umbonal slope. On the type specimen the spines either lie at the point of bifurcation of a costa, or give rise to a single large costa in place of several smaller ones, on the anterior slope.

There is considerable variation in the specimens referred to this species. That from loc. 382 has a strong geniculation, broad visceral region, broader, deeper sinus, and coarser costae than the type. The specimens from loc. 193 are small and more inflated in the visceral region.

Dimensions in mm. of several specimens:

| | 1 (loc. | 2 (loc. 380 —crushed) | 3 (loc. 380 —type) | 4 (loc. 193) |
|---|---|---|---|---|
| Length | 38 | 45? | 40 | 33 |
| Width | 53 | 70 | 58 | 50 |
| Convexity | 29 | — | 23 | 19 |

*Observations:* This form is distinguished from *P. hessensis* and *P. capitanensis* by being broader, with an indistinct sinus, and by the presence of a row of spines along the visceral shoulders. From *P. wolfcampensis* it differs in being much broader.

*Distribution:* Lower Gym formation (Hueco Mountains) at 380 (type), 382?, and 385. Upper Wolfcamp formation (Glass Mountains) at 193.

**PRODUCTUS INCA d'Orbigny**

Pl. XI, Fig. 2; Pl. XIII, Figs. 4–5

1842 *Productus inca.* d'Orbigny, Voy. dans l'Amér. mérid., pt. 4, vol. 3, p. 52, pl. 4, figs. 1–3.

1909 *Productus leei.* Girty, Manzano Group of Rio Grande Valley, U. S. Geol. Sur. Bull. 389, pp. 56–57, pl. 7, fig. 1.
San Andres formation; Engle, Caballos Mountains, Nogal Creek, and San Andres.

1914 *Productus semireticulatus* var. *Inca.* Kozlowski, Ann. de Pal., vol. 9, pp. 30–33, text fig. 5, pl. 3, figs. 1–7; pl. 4, fig. 18. See for synonymy.

1914 *Productus semireticulatus.* Douglas, Geological sections through the Andes, Quart. Jour. Geol. Soc., vol. 70, pp. 34–35, pl. 8, fig. 6.

*Description:* Shell medium-sized, subquadrate (or subtriangular if, as is commonly the case, only the visceral part is preserved), generally somewhat wider than long, with right cardinal angles. Pedicle valve strongly convex, geniculate. Beak low, projecting slightly over the cardinal line. Sides of the visceral area rounded but fairly steep, sharply set off from the ears, which are small, low, and flattened. About 10 mm. anterior to the beak there originates a narrow, shallow sinus. Brachial valve slightly concave in the visceral region, the margins of which are sharply geniculate, with a long trail anterior to the geniculation. In the median part there is an elevation corresponding to the sinus of the other valve. Brachial ridges similar to those of *P. ivesi.*

Surface of the pedicle valve marked by rather strong radial costae, crossed in the visceral part by concentric wrinkles of about the same size as the costae. Costae fairly regular and rounded, separated by rounded striae; about 6 to 7 occur in the space of 10 mm. Each costa bears, in the anterior part of the shell, several spine bases, which are unevenly distributed over the surface. A row of small spine bases lies along the cardinal margin on each side of the beak. Surface of the brachial valve similarly ornamented.

Dimensions in mm. of several specimens:

| | 1 (loc. 421) | 2 (loc. 424) | 3 (loc. 424) | 4 (loc. 414) | 5 (*P. leei*) |
|---|---|---|---|---|---|
| Length | 28 | 35 | 28 | 32 | 24 |
| Width | 32 | 44 | 32 | 39 | 31 |
| Convexity | 13 | 19 | 16 | 17 | 13 |

*Observations:* This species was imperfectly known previous to the work of Kozlowski; the name *P. leei* was therefore applied to it by Girty in 1909. There can be little doubt that the Texas specimens are conspecific both with those from New Mexico and those from Bolivia. *P. inca* is distinguished by its low beak, coarse costae, strong wrinkles, and rather abundant spines. It somewhat resembles *P. magnicostatus,* but has a less incurved beak, smaller and more regular costae, and smaller spines. It is more coarsely costate and has a much less incurved beak than *P. semireticulatus hermosanus.*

*Distribution:* Middle Gym formation (Hueco Mountains) at 414, 415, 417, 418, 419, 421?, 424, 428, 439, and 434. Upper Gym formation (Hueco Mountains) at 436, 439, 449a, 444, and 449. Upper Gym formation (Wylie Mountains) at 508?. Chupadera formation (Salt Basin) at 420 (common).

### PRODUCTUS IVESI Newberry

Pl. XII, Figs. 1–6

1861 *Productus ivesi.* Newberry, Rept. upon Col. River of the West Exp. by Lieut. Ives, p. 122, pl. 2, figs. 1–8.

Middle Carboniferous (Kaibab) limestone: banks of the Colorado near mouth of Diamond River.

1877 *Productus semireticulatus* (pars). White, U. S. Geol. Surv. W. 100th Mer., vol. 4, p. 111, pl. 8, figs. 1a–b (not 1c). (The legend of the plate refers to the figures as *P. semireticulatus* var. *ivesii.)*

Carboniferous: Near Bear Spring, Camp Wingate, and near Santa Fe, New Mexico; Camp Cottonwood, Old Mormon road, Lincoln County, and top of Grass Mountain, Ely range, 35 miles north of Pioche, Nevada; head of Patridge Creek; near Bill William's Mountain; Tenney's Ranch; Kaibab Plateau; head of Dry Fork and Kanab Canyon ("Aubrey" limestone), Arizona; crest of Hurricane Hill, near Toquerville, and Meadow Creek, south of Fillmore, Utah. (Some of these localities, such as that near Santa Fe, New Mexico, are probably Pennsylvanian and are not *P. ivesi.)*

1909 *Productus ivesi.* Girty, Manzano Group, U. S. G. S. Bull. 389, p. 65.

San Andres limestone, Nogal Creek, Mesa del Yeso, and San Andreas; Yeso formation, Mesa del Yeso.*

*Description:* Shell large, attaining a great length in proportion to width when mature, strongly convex, with extended, well-defined, strongly arched ears. Pedicle valve regularly convex, the curvature becoming gradually less toward the front. After a curvature of about 90°, or about 15 mm. in front of the beak, a slight flattening appears in the median part of the valve, and develops into a shallow but rather narrow median sinus. At the same stage the sides begin to slope more abruptly, and the low, broad ears become sharply defined from the steep lateral slopes. The entire curvature of the pedicle valve is about 180°. Brachial valve gently concave, the deepest concavity being near the beak, in front of which the surface flattens out.

Surface marked by strong radial costae, which are rather small in the umbonal region but attain a constant size over the main part of the shell, a few additional ones being introduced there by intercalation. On the pedicle valve the posterior part is crossed by concentric wrinkles, which are asymmetrical, the steepest sides being toward the beak; they are slightly more prominent and more widely spaced than the costae, and give rise to nodes where they cross the costae. Some of the nodes are the bases of large hollow spines, which are most abundant near the hinge, along which there are many large, irregularly placed spines. The ears are strongly wrinkled at right angles to the costae, and from these wrinkles there arise large laterally projecting spines. Over the median and anterior parts of the shell spines are less numerous, but still relatively abundant, occurring at intervals of 4 or 5 mm. on each costa. Their bases are hollow, and about 0.75 mm. across. The costae are somewhat variable in size, the intercalated ones being generally smaller for some distance anterior to their point of origin. In some shells the sinus pinches together forward, and the costae run into each other anteriorly along the sinus. The distance between centers of adjoining ribs is 1.5 to 2 mm., the costae and sulci being of equal dimensions. On the brachial valve the posterior part is wrinkled, but over a larger proportion of the surface than on the other valve; where radial ribs and concentric wrinkles intersect there are pits, equivalent in position to the spines of the pedicle valve. Surface marked also by fine, irregular, closely crowded undulating growth lines.

Shell of the pedicle valve very thick in large individuals; one shell is 12 mm. thick in the median part, and 7 mm. at the side, but most shells are considerably thinner. The interior of the pedicle valve bears large, deeply impressed muscle scars. The diductor scars are deeply incised, coarsely and irregularly striated, largest at about a third the length of the shell, and tapering toward the posterior end. The adductor muscle scars lie on the raised platform between the diductors, are multilobate, and are separated by a low ridge.

The interior of the brachial valve has a large trifid cardinal process, which extends beyond the hinge-line; the three branches are of about equal size, and the median one is divided by a small groove. From the process two blunt lateral ridges extend, one on each side, along the hinge-line. In the middle the process is joined by a broad median ridge, which extends forward between the muscle scars as a narrow, slightly elevated septum. The brachial ridges arise from the septum, extending first backward at angles of about 30° to the septum, then curve evenly around until parallel with the septum, and still further until they are directed backward, this end being high and straight and lying nearly parallel to the septum. Within each brachial loop there is a short elevated ridge connected with the brachial ridges, lying at an angle of 45° to the

*The *"Productus* (undet. sp.) compare *P. Ivesii"* of Meek (1877), also named by him *Productus longus* is a *Buxtonia* and occurs in the Pennsylvanian.

septum and directed toward the beak. A faint ridge connects the end of the loop again with the median septum. Inner median surface of the valve marked by short forward-projecting spines shingled over one another; the surface between the spines is strongly punctate. In front of the spinose region the surface is longitudinally roughly striated.

Dimensions in mm. of several shells:

| | 1 (loc. 123) | 2 (loc. 107) | 3 (loc. 174) | 4 (loc. 174) |
|---|---|---|---|---|
| Length | 50 | 50 | 70 | 51 |
| Width | 69 | 55 | 60 | 49 |
| Convexity | 27 | 28 | 40 | 28 |

Number 3 gives the typical dimensions of an old elongate shell. Number 4 is an unusually narrow specimen, and numbers 1 and 2 are relatively short.

If the brachial valve is somewhat caved in the valves pull apart and there is a strong appearance of a cardinal area.

*Observations:* This common Kaibab-San Andres species is very abundant in the Leonard and in the upper Hess of the Glass Mountains, and their equivalents in Sierra Diablo. Four badly preserved shells or parts of shells that apparently belong to *P. ivesi* were collected from beds of supposed Word age in the northeastern Glass Mountains and Sierra Madera. At localities 256 and 257 they are associated with fossils that are probably Word. The uncertainty of differentiation of Leonard and Word in this region makes the Word age of these fossils, however, doubtful, especially since *P. ivesi* has not been found in the Word in its normal phase in the western Glass Mountains. *P. ivesi* was referred to by Böse as "very large Producti of the group of *Productus sino-indicus* Frech."*

From *P. semireticulatus* and its varieties, *P. ivesi* differs in being much larger, narrower in proportion to length, and in having coarser costae and more numerous spines. Typical examples of the two cannot be confused. It differs from *P. inca* mainly in its much larger size and far more abundant spines. *P. uralicus* Tschernyschew differs in having fewer spines and probably also in some details of general form. *P. spiralis* Waagen has much larger costae and fewer spines (although, as Diener has pointed out, the specimens figured by Waagen have abnormally large costae). *P. sino-indicus* Frech is not well enough figured for satisfactory comparison. It appears to be narrower than *P. ivesi,* but is similar in other respects. No other foreign species appears to be closely similar.

*Distribution:* Hess formation (Glass Mountains) at 4, 7, 17, 102, 105, 106, 107, T108, 113, 128, 206, 211, 222, and 223. Leonard formation (Glass Mountains) at 9, T9, 15, T15, 21, 22, 24?, 31, 37, 38, 39, T41, 83?, 104, 119, T119, 120, 121, 123 (common and well-preserved), 124, 126, T128, 174 (extremely abundant), T233, T240, 301, D1, D4, D11, and Tb. Word formation? (Glass Mountains) at 256, 257, and 262. Hess formation (Sierra Diablo) at 141, 470, 471, 472, 476, 477, 479, and 481. Bone Canyon member (Sierra Diablo) at 484 (one of these specimens has exceptionally coarse costae and large spines), and 489. Victorio Peak member (Sierra Diablo) at 493, 495, 499, 500, and 503. Victorio Peak member (Brackett Draw) at Taa. Leonard formation (Baylor Mountains) at 503. Leonard formation (Shafter) at Tk, Tm?, 516, and 518. Leonard formation (Finlay Mountains) at 511 and 513. Gym formation (Wylie Mountains) at 508?. Limestone of Puertacitas at Puertacitas (Las Delicias) (a more finely costate individual than typical *P. ivesi).*

*U. of Texas Bull. 1762, p. 24.

**PRODUCTUS LEONARDENSIS King, n. sp.**

Pl. XIV, Figs. 4–9

*Description:* Shell medium-sized, subquadrate, with the hinge-line equal to or only slightly less than its greatest width. In most shells the hinge-line is straight, but in some the parts on opposite sides of the beak lie at an angle to each other of about 140° rather than 180°.

Pedicle valve strongly geniculated, being relatively flat in profile in the posterior and anterior halves but with the front part lying at a right angle to the visceral region, the only curvature being near the belt of geniculation. Region near the hinge nearly flat, with a low but abrupt elevation at the beak. Visceral angle relatively very narrow. Beak pointed, very low, ending essentially within the same line as the hinge margin. Ears broad and flattened, set off from the triangular, raised marginal part of the shell by well-defined sinuses. Median sinus narrow and subangular, extending from nearly the point of the beak, giving the shell a bilobed appearance. It becomes broader toward the front, but is deepest just behind the geniculation.

Brachial valve slightly concave near the beak but nearly flat near the cardinal margin, marked by a low fold corresponding to the sinus of the other valve. The brachial valve is geniculated toward the front, but less strongly than the pedicle valve.

Surface of the flattened posterior region marked by radial lirae and concentric wrinkles of the same strength, which are not asymmetric. The wrinkles converge near the hinge and become stronger; some of them give rise, about 1 to 2 mm. in front of the hinge margin, to spines which are arranged in two parallel rows on each side of the beak and are directed backward over the hinge-line. The concentric wrinkling extends as far forward as the beginning of the region of geniculation. The radial lirae number 7 to 10 in the space of 5 mm., increasing by bifurcation, the new lirae coming in in the direction of the lateral margins, while the lirae near the sinus are, except near the beak, parallel and simple.

Dimensions in mm. of several specimens:

| | 1 (loc. 120) | 2 (loc. 82) | 3 (loc. 120) | 4 (loc. 120) |
|---|---|---|---|---|
| Width | 24 | 28 | 27 | 28 |
| Length | 20 | 38 | 28 | 24 |

The differences in proportion of length to width are due largely to the length of the posterior geniculated part of the shell, for the width does not increase much after the reticulate posterior part of the shell has been formed. Mature specimens show evidence of lateral growth simultaneous with the forward growth of the geniculated front margin. In these the hinge-line bends forward at about 45° at the same stage of growth at which geniculation

starts in the central part of the shell. These laterally extended parts of the wings turn sharply upward, forming a flange of the upturned brachial valve along the lateral margins, which may be several mm. wide. The flange lies at about a right angle to the rest of the brachial valve; toward the front it merges into the anterior extension of the shell, which is not turned back.

*Observations:* This species resembles the species of *Linoproductus* in the relative fineness of its radial ornamentation and in the presence of a double row of spines along the hinge, but it is clearly a true Productus. Several Russian species, especially *P. gruenewaldti* Krotow and *P. moelleri* Stuckenberg, are very similar to this species in ornamentation, but the configuration of the shells is quite different. The strong geniculation of *P. leonardensis*, the depth of its sinus, the presence of a flange along the lateral margins, and the fine sculpture distinguish it easily from all other described species of similar general nature.

*Distribution:* Hess formation (Glass Mountains) at 8. Leonard formation (Glass Mountains) at 4, 7, 9, 10, 11, 12, 13, 15, 21, 31, 37, 82, 83, 102, 104, 119, 120 (types), T120, 123, 231, 232, and Tb. Word formation (Glass Mountains) at 48 (single specimen). Bone Canyon member (Delaware Mountains) at 491. It is a very abundant and widespread Leonard fossil.

**PRODUCTUS GRATIOSUS OCCIDENTALIS Schellwien**

Pl. XIV, Figs. 1–3

1892 *Productus gratiosus* Waagen var. *occidentalis*. Schellwien, Die Fauna des karnischen Fusulinenkalks, Palaeontographica 39, pp. 24–25, pl. 3, figs. 6–9; pl. 8, fig. 25.

1927 *Productus gratiosus* var. *occidentalis*. Chao, Productidae of China, Pal. Sinica, vol. 5, fasc. 2, pp. 44–47, pl. 4, figs. 6–10.

*Description:* Shell small, transversely subquadrate, with the hinge-line equaling the greatest width. Pedicle valve flattened posteriorly, then broadly geniculated beyond the reticulated region, in front of which the curvature is slight and even. Beak pointed, only slightly incurved over the hinge. Ears flattened, set off from the rest of the shell by a concavity. Median sinus originating near the beak, narrow and moderately deep, of constant size throughout its length. Brachial valve slightly and regularly concave in the visceral region and sharply geniculated toward the front. Median fold low and distinct.

Surface of both valves marked by many high, rounded, relatively coarse costae, separated by deep sulci, which are much narrower than the costae. The costae in the median sinus of the pedicle valve are smaller than those on each side, and they converge toward the center, where they unite or become small and disappear. Posterior region marked by many fine regular concentric ridges, giving the surface a delicate reticulation. They are somewhat smaller than the costae. There is a flange-like lateral ridge on each visceral slope, which originates as a faint elevation slightly behind the geniculation, and extends at first oblique to the radial costae. After passing over the geniculation it joins several costae and develops into a large thickened ridge, bearing a row of several spines. There are many forward-directed spines that arise from the reticulated surface, and each costa bears several spines between the anterior end of the reticulate region and the front margin.

Dimensions in mm. of several specimens:

| | 1 (loc. 93) | 2 (loc. 93) | 3 (loc. 93) | 4 (loc. Sd) |
|---|---|---|---|---|
| Length | 14 | 16 | 17 | 20 |
| Width | 20 | 22 | 22 | 26 |

*Observations:* These specimens agree perfectly in all details with Chao's description and figures, detailed comparison with Schellwien's rather inadequate figures being difficult. This species is distinguished from all others in the Glass Mountains fauna by the strong reticulation of the posterior part, the large, rounded costae separated by very narrow sulci, and the large number of spines on the costae. The variety is distinguished from typical *P. gratiosus* Waagen by the presence of the thickened spine-bearing rib on each side of the visceral area.

*Distribution:* Gaptank formation (Marathon basin) at Sd (bed 3 of the Gap Tank section). Wolfcamp formation (Glass Mountains) at T91 and 93.

**PRODUCTUS MULTISTRIATUS Meek**

Pl. XIII, Figs. 1–3

1860 *Productus multistriata*. Meek, Proc. Acad. Nat. Sci. Philadelphia, vol. 12, p. 309.
Upper Carboniferous?: East side of Long Valley, lat. 39°57′; long. 115°10′.

1876 *Productus multistriatus*. Meek, Simpson's Rept. Expl. across the Great Basin of Utah, p. 350, pl. 1, figs. 8a–b.
Yellow ls. (Upper Carboniferous?): Long Valley.

1876 *Productus multistriatus*. White, Powell's Rep. Geol. Uinta Mts., p. 90.
Lower Aubrey group: Confluence of Grand and Green rivers, Utah.

1877 *Productus multistriatus*. Meek, King's U. S. Geol. Expl. 40th Parallel, vol. 4, pp. 76–78, pl. 8, figs. 3–3e.
Carboniferous ls.: Near center of the eastern margin of Nevada; Ruby Group and Mahogany Peak, Eagan Range.

1920 *Productus multistriatus*. Girty, U. S. G. S. Prof. Paper 111, pl. 56, figs. 7–7a.
Permian: Utah.

*Description:* Shell rather large, wider than long, with extended ears, which are broad and not sharply defined from the rest of the shell. The greater width than length stated for the present specimens may not actually be the case for the species if the anterial trail were completely preserved. Profile of the pedicle valve of very low convexity in the posterior part, then sharply geniculated. Beak low and bluntly pointed. Median sinus originating near the beak, but only slightly developed posterior to the geniculation, in front of which it is deep and narrow. Surface marked by fine radial lirae, 8 to 12 of which occur in the space of 5 mm. Only very faint concentric undulations are present. Spines are arranged in a row along the

hinge, and are inclined strongly outward; the other spines are scattered sparsely and irregularly.

Dimensions in mm. of several specimens:

| | 1 (loc. 144) | 2 (loc. 144) | 3 (loc. 6) | Las Delicias | Meek's type |
|---|---|---|---|---|---|
| Length | 24 | 23 | 18+ | 20 | 25 |
| Width | 40 | 42 | 35 | 21 | 44 |

*Observations:* This species is characterized by the flat visceral region of its pedicle valve, the strong geniculation at mid-length, and the fine liration. The specimens from Las Delicias are much smaller than those from the Glass Mountains. The forms identified by Tschernyschew as *P. multistriatus* var. have much coarser costae than the American form, and no prominent geniculation. *P. weyprechti* Toula is probably the most similar foreign species. It has coarser ornamentation and is less strongly arcuate.

*Distribution:* Word formation (Glass Mountains) at 6, 144, and T240. Limestone of Cerro Caballo (Las Delicias, section 1, bed 14, west of the Noria de Malascachas.

### PRODUCTUS INDICUS Waagen

Pl. XIII, Figs. 6–9

1882 *Productus indicus.* Waagen, Productus-limestone Fossils, Pal. Indica, ser. 13, vol. 1, pt. 4, pp. 687–690, pl. 70, pl. 71, fig. 1. (See for further synonymy.)

"The commonest species of the Salt-range. It is restricted to the middle and upper divisions of the Productus-limestone."

1915 *Productus indicus.* Diener, Anthracol. Faunae of Kashmir, Kanaur, and Spiti, Pal. Indica, n. s., vol. 5, mem. 2, pp. 66–67, pl. 6, fig. 15.

Zewan beds, Kashmir.

*Description:* Pedicle valve strongly convex, showing in profile a regular curve; beak greatly incurved over the hinge-line. Wings very large, arcuate, and sharply set off from the rest of the shell. Hinge-line straight, wide, making the greatest width of the shell. Median sinus deep but fairly broad, originating near the beak and maintaining a nearly uniform width and depth from a place about 20 mm. in front of the beak to the anterior margin. Brachial valve concave in the visceral region and strongly geniculated at the outer border of that region.

Shell marked by very large, very unequal costae, some of which bifurcate and some unite with an adjacent costa. The strength of the costae varies from one specimen to another. In the sinus some costae may converge and disappear. Rather small, subequal wrinkles cover the visceral region and extend more or less upon the wings. A double row of heavy spines is borne on the posterior slope of the ears and the rest of the shell is marked by much smaller spines borne at intervals of 5 or 6 mm. along the costae.

Dimensions in mm. of several specimens:

| | 1 (loc. 46) | 2 (loc. 46) | 3 (loc. T251) |
|---|---|---|---|
| Length | 37 | 44 | 30 |
| Width | 52 | 74 | 48 |

*Observations:* The Texas specimens show no characters that are not included within the range of variations of the specimens figured by Waagen. *P. indicus* is distinguishable from *P. ivesi* by its coarser costation, smaller size, and less prominent reticulation, and from *P. occidentalis* by its larger size, much greater breadth, and larger ears.

*Distribution:* Leonard formation (Glass Mountains) at Tb. Word formation (Glass Mountains) at 46 (abundant), 58, 141, and T251. Upper Hess formation (Sierra Diablo) at 480?. Leonard formation (Shafter) at 518. Word? formation—thin-bedded zone (Shafter) at Tk.

### PRODUCTUS OCCIDENTALIS Newberry

Pl. XIV, Figs. 10–14

1858 *Productus costatus.* Marcou, Geol. of N. A., p. 46, pl. 5, fig. 5 (?).

Carboniferous: Cedar Creek, Mogollon Mountains, tributary of Gila River, sources of Colorado Chiquito.

1861 *Productus occidentalis.* Newberry, Rept. upon Col. R. of the West Exp. by Lieut. Ives, p. 122, pl. 2, figs. 9–10.

Cherty limestone near top of Carboniferous (Kaibab): banks of Cascade River near junction of Great and Little Colorado.

1908 *Productus occidentalis.* Girty, The Guadalupian Fauna, U. S. G. S. Prof. Paper 58, pp. 262–263, pl. 12, figs. 4–4c.

Middle Capitan, Capitan Peak (sta. 2926); dark limestone, Pine Spring (sta. 2930).

*Description:* Shell medium-sized, relatively very narrow, with the hinge-line forming the greatest width of the shell. Pedicle valve strongly inflated in the umbonal region, its curvature decreasing toward the anterior margin. The median sinus originates about 1 cm. in front of the beak and becomes broad and fairly deep. Ears broad, separated from the rest of the shell by deep sinuses. Costae uneven, large, from 2 to 3 mm. across and 20 to 26 in number. In some specimens the costae bordering the sinus are particularly large, while those in the sinus are much smaller and, in shells becoming laterally compressed toward the front, uniting toward the anterior margin. Other shells, however, are quite regular and the costae in the sinus are only slightly smaller than those of the rest of the shell. On the anterior part of the shell a few sparsely distributed large spines arise from the costae. Posterior and alar region marked by a few very faint concentric wrinkles, which give a somewhat nodose appearance. Some of the nodes give rise to spines, one of which, arising from an ear, is more than 2 cm. long.

Brachial valve gently concave, the deepest concavity being near the beak.

Pedicle valve with deeply incised, irregularly striated diductor muscle scars; adductor muscle scars multilobate, marking the raised platform between the diductors. Interior of the brachial valve having a large cardinal process, from the anterior end of which extends a low median ridge, and on each side a lateral ridge which extends along the hinge-line. Muscle scars small, oval, and elevated, bordered at the front by the brachial ridges, which arise from the median septum and extend laterally half way to the shell margin, curving continually forward, then reversing to produce a hook-like form, the ends pointing

back toward the point of origin but not quite meeting the septum. The rest of the interior surface of the valve marked by fine granules.

Dimensions in mm. of several specimens:

| | 1 (loc. 128) | 2 (loc. 123) | 3 (loc. 174) | Kaibab, Grand Canyon |
|---|---|---|---|---|
| Length | 36 | 27 | 40 | 35 |
| Width | 25 | 26 | 37 | 32 |
| Convexity | 20 | 14 | 26 | 18 |

Some of the Grand Canyon specimens are relatively slightly broader than the typical ratios given above.

*Observations:* This shell is easily distinguishable by its coarse costae, narrow outline, and prominent sinus, and by the limited area of reticulation and the sparseness of spines. The specimens described by Girty from the Delaware Mountain and Capitan, as well as those from the Word of the Glass Mountains, cannot be distinguished from the typical Kaibab form and its variations. However, it is very rare in the Word, and may be said to be common only in the Leonard and its equivalent.

*P. occidentalis* is distinguishable from *P. costatus* and *P. magnicostatus* by its stronger costation. The closest similarity among foreign species of *Productus* is shown by *P. grandicostatus* Chao, a form so much like this that it shows no marked difference. Since the Chinese species is based on but one specimen, it is rather difficult to make satisfactory comparison. There are several large-ribbed species of *Productus* in the Salt Range fauna, but all are much wider in proportion to length.

*Distribution:* Leonard formation (Glass Mountains) at 7, T119, T120, 123, 128, 174 (abundant), 232, and D3. Word formation (Glass Mountains) at 144, T148, Tx148, and 241. Bone Canyon member (Delaware Mountains) at 491. Victorio Peak member (Brackett Draw) at Taa. Gym formation (Diablo Plateau) at 446.

### PRODUCTUS SCHUCHERTI King, n. sp.

Pl. XV, Figs. 1–5

1908 *Productus* sp. c. Girty, The Guadalupian Fauna, U. S. G. S. Prof. Paper 58, pp. 256–257.
Delaware Mountain formation, "Comanche Canyon," Glass Mountains.

*Description:* Shell medium-sized, strongly inflated, wider than long. Pedicle valve strongly arcuate longitudinally, somewhat geniculate. Beak small and pointed, strongly incurved. Ears well-defined, strongly convex, and very much extended. Sinus deep and rather narrow. Brachial valve deeply concave, with a low median fold; beak concave and pointed; cardinal area entirely wanting. Surface marked by low, rounded costae, numbering about 5 in the space of 5 mm.; they may be obsolescent near the anterior margin. Posterior part of most specimens not crossed by well marked concentric wrinkles, but having an obscurely nodose appearance; distinct wrinkles are, however, present on the ears of both valves. The ears bear two rows of rather small spines, set well back from the cardinal margin and inclined strongly outward toward the cardinal extremities. The rest of the shell bears somewhat coarser, sparsely scattered spines.

Dimensions in mm. of several specimens:

| | 1 (loc. 120) | 2 (loc. 120) | 3 (loc. 120) | 4 (loc. 128) | 5 (loc. 124) |
|---|---|---|---|---|---|
| Length | 22 | 22+ | 26 | 27 | 22 |
| Width | 44 | 32 | 40+ | 34 | 40 |

*Observations:* This species is distinguishable by its low, medium-sized ribs, faintness of the wrinkling of the posterior part, prominent sinus, and extended ears. It belongs to the group of *P. boliviensis* d'Orbigny, and can hardly be distinguished from the specimens identified by Tschernyschew as *P. boliviensis* and *P. gruenewaldti*, except that it has a deeper sinus. *P. guadalupensis* Girty is somewhat similar, but is more strongly enrolled and has no concentric wrinkles and much finer costae. Girty's figures of *P. guadalupensis*, however, show great similarity to *P. schucherti*, (figures 1 and 1a, plate 22), having costae almost as coarse as those of the latter species.

*Distribution:* Hess? formation (Glass Mountains) at T205. Leonard formation (Glass Mountains) at 9, 12, 13, 31, 38, T82, 119, T119, 120 (abundant; types), T120, 123, 124, 128, 174, 232, Tb, Tba, D1, D4, D5, D8. Word? formation (Glass Mountains) at T240 and 265. Word? formation thin-bedded zone (Shafter) at Tq. Limestone of Cerro Caballo (Las Delicias) section 1, bed 14. Bone Canyon member (Sierra Diablo) at 484. Victorio Peak member (Sierra Diablo) at 494. Leonard formation (Baylor Mountains) at 503.

### PRODUCTUS SEMIRETICULATUS HERMOSANUS Girty?

Pl. XIV, Figs. 15–17

1903 *Productus semireticulatus* var. *hermosanus*. Girty, U. S. G. S. Prof. Paper 16, pp. 358–359, pl. 2, figs. 1–4.

?1915 *Productus inflatus* var. *coloradoensis*. Girty, U. S. G. S. Bull. 544, pp. 64–65, pl. 8, figs. 1–2.

?1921 *Productus inflatus coloradoensis*. Plummer and Moore, U. of Tex. Bull. 2132, pl. 13, fig. 24. Not *P. semireticulatus hermosanus* Plummer and Moore, pl. 20, fig. 23; pl. 25, figs. 1–2.

*Description:* Shell medium-sized to large (up to 60 mm. across), marked by a very shallow median sinus. The costae average 6 to 8 in 10 mm. Concentric wrinkles form strong nodes in the posterior region. Spines are rather sparsely distributed, being generally not closer than 10 mm. from each other on the non-reticulate part of the shell.

*Observations:* The shells referred to this variety are mostly very fragmentary. It may be seen, however, that they agree with *P. semireticulatus hermosanus* in the strength of their costae and arrangement of spines. Because of the imperfect nature of the Wolfcamp material, as well as the great confusion which has attended the identification of this variety by other authors, exact identification is not possible. Girty, in his original description, mentioned as points of difference from *P. semireticulatus* the small size, more inflated and incurved form, fewer and larger spines, and more extended and enrolled ears. Since their original description, the identification of *P. hermosanus* and of *P. inflatus (coloradoensis)* appear to have been reversed. Girty originally pointed out emphatically that the only important distinguishing character between them was the finer lirae of *P. inflatus (coloradoensis)*. In

the report on the Wewoka fauna, however, he figured and described a coarsely costate form under that name. The specimens identified by Plummer and Moore as *P. semireticulatus* var. cf. *hermosanus* (U. of Tex. Bull. 2132, pl. 25, figs. 1–2) agree almost exactly with Girty's original figures of *P. inflatus (coloradoensis)*, while those identified by them as *P. inflatus coloradoensis* are very similar to the original figures of *P. hermosanus*.

*Distribution:* Wolfcamp formation, *Uddenites* zone (Glass Mountains) at 88, 88s, T88, 94, 94s, 107, 175, and 201. Middle and Upper Wolfcamp formation (Glass Mountains) at 87 low, 87 high, 89, 91, T91, 92, 93, T93, and 195.

**PRODUCTUS SEMISTRIATUS Meek**

Pl. XV, Figs. 6–11

1860 *Productus semistriatus.* Meek, Proc. Acad. Nat. Sci. Phila., vol. 12, p. 309.
"Coal Measures": Timpanogos Canyon, lat. 40°22′, long. 111°38′.

1876 *Productus semistriatus.* Meek, Simpson's Rept. Expl. across the Great Basin of Utah, p. 347, pl. 1, figs. 7a–b.
"Coal Measures": Timpanogos Canyon.

1877 *Productus semistriatus.* Meek, Rept. of the Geol. Expl. of the 40th Parallel, pp. 74–75, pl. 7, figs. 8–8a.
Black bituminous limestone: southeast of Great Salt Lake, lat. 40°22′n, long. 111°38′w.

1920 *Productus semistriatus.* Girty, U. S. G. S. Prof. Paper 111, pl. 53 and p. 651.
Upper Mississippian: Utah.

*Description:* Shell medium-sized, transversely subquadrate. Pedicle valve highly inflated, somewhat geniculated at about a third of the length and gently and evenly convex in front of and behind the geniculation. Beak small, projecting somewhat beyond the hinge. Ears short, not well-defined, the umbonal region being relatively low; the hinge-line equals the greatest width of the shell. Median sinus broad and indistinct; in some specimens it is quite absent, there being only a flattening in the median region. Brachial valve evenly concave.

Visceral region of the pedicle valve covered by moderately distinct low, rounded costae, about 6 of which occur in the space of 5 mm.; they become obsolete over the anterior part of the shell. There is a single row of rather slender spines along the hinge-line, which apparently were directed outward away from the beak. Scattered spines are rather abundant over the body of the shell, being small on the visceral region and increasing in size toward the front margin. On the anterior slope, where the costae are obsolescent, the shell may bear several low broad, ribs, each of which arises in front of a stout spine base. The outer shell layer in this smooth anterior part is marked by many and closely spaced fine punctae. Where the surface is broken away, it is seen that posterior to the smooth region the sulci between the costae are marked by rows of large pores, just as the striae of *Chonetes granulifer* are marked by rows of punctae. The rest of the inner surface is even more finely punctate than the outer smooth surface.

Dimensions in mm. of several specimens:

| | 1 (loc. 92) | 2 (loc. 193) | 3 (SW of Hueco T.) |
|---|---|---|---|
| Length | 22 | 23 | 22 |
| Width | 33 | 38 | 30 |
| Convexity | 13 | 16 | 14 |

A specimen from loc. 93 which probably belongs to this species shows on the inside of the brachial valve a low papillose arcuate ridge, suggesting *Marginifera* structure, but not as distinct as in *Marginifera*. The specimen from loc. 90 is not geniculate like the typical ones, and has fine costae, and the ones from 94 and 88s have extended ears, which are marked on each side of the beak by a row of laterally directed spines.

*Observations:* This species is distinguishable by its geniculate profile, very shallow sinus, faintness of the reticulation of the posterior part, the merging of the costate posterior part into the smooth anterior part of the surface, the few large spines in the anterior smooth region, and the punctate shell. In 1920 Girty figured *P. semistriatus* on a plate of Upper Mississippian fossils from Utah. In 1927*, however, he does not list it as a Brazer fossil, but cites *Prod.* aff. *semistriatus* from the Pennylvanian Wells formation. As no further discussion is given, it cannot be told whether the first listing was later found to be incorrect, or how closely similar the Wells form is to the *Productus semistriatus* of Texas.

*Productus semistriatus* is very similar to *P. chandlessi* Derby, the only apparent difference being that *P. semistriatus* has more or less of a median sinus, while the Brazilian species has none at all. *P. schucherti* differs in being much broader in proportion to length, in not having a smooth anterior region, and in its more extended ears.

*Distribution:* Wolfcamp formation (Glass Mountains) at 88s, T88, 90, 92, 93?, 93s, 94, and 196?. Lower Gym group (Hueco Mountains) at 376, 381, 382, 384, 388, 390, and Baker collection 4 miles southwest of Hueco Tanks. Upper Gym group (Hueco Mountains) at 438 and 440a. Upper Wolfcamp formation (Sierra Diablo) at 456 and 460.

**PRODUCTUS WOLFCAMPENSIS King, n. sp.**

Pl. XVI, Figs. 1–5

*Description:* Shell medium-sized, subquadrate, longer than wide. Pedicle valve highly inflated, the longitudinal profile showing a progressive decrease in convexity from beak to anterior margin, but without distinct geniculation. Beak small and pointed. The median region is slightly convex in transverse profile, or somewhat flattened, there being no distinct sinus. Ears small but well defined. Brachial valve evenly concave.

Surface marked by rather fine costae, which toward the anterior part become very uneven in size, some of them tending even to coalesce. The posterior one-third or one-fourth of the surface is crossed by low and rather fine concentric wrinkles. Small spines are rather abundantly distributed over the anterior two-thirds of the surface and on the ears. Very small spines seem to be abundant over

*U. S. G. Sur. Prof. Paper 152, p. 73.

the visceral region, but are shown only on very well-preserved specimens.

Dimensions in mm. of several specimens:

| | 1 (loc. Tmm) | 2 (loc. 460) | 3 (loc. 198) |
|---|---|---|---|
| Length | 34 | 41 | 30 |
| Width | 38 | 44 | 31 |
| Convexity | 22 | 23 | 15 |

*Observations:* This species is distinguished by the greater length than width, the absence of a distinct sinus on the pedicle valve, regular convexity, and fine costae, which are uneven on the anterior part of the shell. The most nearly related species is *P. huecoensis*, which is, however, larger, much broader in proportion to length, and bears a row of spines along the visceral slope. The two species occur together rather abundantly at loc. 456.

*Distribution:* Upper Wolfcamp formation (Glass Mountains) at 2?, 83s?, 195?, 196, and 198. Upper Wolfcamp formation (Sierra Diablo) at 456 (common) and 460. Lower Gym formation (Hueco Mountains) at Tmm (type) and 383?.

#### PRODUCTUS sp.

Several specimens of *Productus* from the Hess and Leonard formations cannot be placed in any of the above species, but are too dissimilar among themselves to be described collectively. Some of them are similar to *P. inca*, but are smaller and more strongly enrolled.

*Distribution:* Hess formation (Glass Mountains ) at Tt. Leonard formation (Glass Mountains) at 9, T9, 11a, 13?, 26, 37, 129, and 174. Bone Canyon member (Delaware Mountains) at 491. Leonard formation (Shafter), Cieneguita beds at Tg (2279), transition beds at Th (15978), and 518. Leonard formation (Finlay Mountains) at 512.

### Genus LINOPRODUCTUS Chao

*Diagnosis:* Shell productoid; pedicle valve strongly convex, brachial valve flattened to slightly concave in the visceral part, strongly geniculated toward the front, resulting in the formation of an anteriorly produced long trail. Surface marked by numerous fine radiating lirae and indistinct concentric wrinkles. A double row of spines is invariably present along the margin of the hinge-line.

Genotype: *Productus cora* d'Orbigny.

*Geologic range:* Mississippian to Permian.

*Observations:* In 1928, the year following the publication of Chao's name of this genus, Fredericks introduced the name *Cora* for approximately the same group. Fredericks, however, excluded the *P. cancrini* group, because of its abundant thin spines, placing it in his genus *Cancrinella;* on the other hand he included the typical form of *Striatifera*, believing that the concavity of the brachial valve was not a generic character.

#### LINOPRODUCTUS CORA (d'Orbigny)

Pl. XVI, Figs. 6–7

1842 *Productus cora.* D'Orbigny, Voyage dans l'Amérique Méridional, t. 3, pt. 4, p. 55, pl. 5, figs. 8–9.

1909 *Productus cora.* Girty, The Manzano Group of the Rio Grande Valley, U. S. Geol. Sur. Bull. 389, pp. 58–61.

Abo sandstone: Mesa del Yeso, Abo Canyon, and Sandia Mountains.

1915 *Productus cora.* Girty, Fauna of the Wewoka Formation, U. S. Geol. Sur. Bull. 544, pp. 68–72, pl. 8, figs. 4–6. (See for American synonymy.)

1916 *Productus cora.* Broili, Permischen Brachiopoden von Timor, Pal. von Timor, lief. 7, 13, pp. 19–22. (See for Eurasian synonymy. The inclusion in the synonymy of *P. lineatus* Waagen may be questioned.)

1927 *Linoproductus cora.* Chao, Productidae of China, pt. 1, Pal. Sinica, ser. B, vol. 5, fas. 2, pp. 132–134, pl. 13, figs. 17–18, pl. 14, figs. 1–4.

*Observations:* There has been some controversy as to the limits of this species, centering particularly about the distinction of *L. lineatus* (Waagen) from *L. cora.* Waagen, Tschernyschew, and Chao have insisted on separating the two types, the distinction being that Waagen's species has a median sinus and a rectangular rather than a triangular form. Kozlowski has shown* that though the type specimen is non-sinuate, specimens from the type locality are either sinuate or non-sinuate, and he therefore concludes that Waagen's distinction is not significant. The quadrate form of *L. lineatus*, however, remains as a distinguishing feature. Chao believes a still more minute division is possible, but this hardly seems practicable, in view of the great variation shown by the species. In the Gaptank and Wolfcamp formations sinuate and non-sinuate forms occur together, and it seems useless to attempt a distinction. None of the sinuate varieties, however, has the rectangular form of *L. lineatus. L. cora* is here, therefore, regarded as a shell of triangular outline, except for the ears, and with or without a shallow sinus.

The Glass Mountains specimens of *L. cora* are identical in every detail with the types from Bolivia as figured by Tschernyschew and Kozlowski, except for the single specimen from loc. 90, which has extremely fine lirae, 10 of them occurring in the space of 5 mm.

In this connection it is well to mention *Productus delawarii* Marcou*, which I think should be placed in synonymy with *L. cora.* The specimen figured appears to be a brachial valve, although Marcou seems to have regarded it as a pedicle valve. The beak is no more pointed, and the hinge no shorter than in most brachial valves of *L. cora,* but these characters were said to distinguish the species. The locality from which it was collected is "a siliceous limestone at the western foot of Delaware Mount on the bank of an affluent of Topofki Creek, Texas. This fossil is very common but never well preserved . . . In company with *P. delawarii* a true *Orthis* is found . . ." This locality is probably somewhere in northern Trans-Pecos Texas.

*Distribution:* Wolfcamp formation, *Uddenites* zone (Glass Mountains) at 88s, 93, 93s, T93, 175, 175a, 201, and 203. Wolfcamp formation (Glass Mountains) at 2?, 75, 87?, 90, T90, 92, 195a, 196, and 198. Upper Wolfcamp formation (Sierra Diablo) at 456, 459, and 460. Lower Hess formation (Sierra Diablo) at 471. Lower Gym forma-

*Ann. de Paléontologie, t. 9 (1914), pp. 50–51.

*Marcou: Geol. of North America, p. 45, pl. 5, fig. 3. 1858.

tion (Hueco Mountains) at 383, 389, and 401. Middle Gym formation (Hueco Mountains) at 414, 416, 425, and 434.

**LINOPRODUCTUS CORA ANGUSTUS King, n. var.**

Pl. XVI, Figs. 8–9

*Description:* Shell medium-sized to large, with a roughly rectangular outline; hinge-line somewhat less than the greatest width of the shell. Posterior part of the pedicle valve very strongly inflated; the anterior half of low convexity in large shells. Beak pointed, but not much incurved over the hinge; the sides of the visceral cavity diverge from it at angles of 85° to 90°. Ears small and flattened, sharply demarked. Median sinus absent, though some specimens show a faint median flattening.

Surface marked by fine lirae, about 9 to 11 of which occupy the space of 5 mm.; they are rounded and separated by striae of varying width, some of which are broader than the lirae. Spines very few in number and inconspicuous. Concentric folds are present on the ears and on the posterior part of the brachial valve, but not on the convex part of the pedicle valve. There is a row of small spines along the hinge-margin.

Dimensions in mm. of several specimens:

| | 1 (loc. 105) | 2 (loc. 105) | 3 (loc. 15) | 4 (loc. 219) | 5 (loc. 222) |
|---|---|---|---|---|---|
| Width | 40 | 30 | 35 | 21 | 24 |
| Length | 43 | 34 | 49 | 26 | 22 |

Specimens assigned to this variety agree in being elongate, with subparallel margins, but otherwise are rather variable. The typical form is that from locs. 105, 107, and 222. The large specimen from loc. 15 is similar in size to that from 105, but is much narrower in proportion to length, and is less strongly convex.

*Observations:* This variety differs from typical *L. cora* in its narrowness and rectangular outline. It is similar to *L. lineatus*, but has no sinus. *L. tenuistriatus* (Verneuil) is smaller, and has a regular curvature instead of a flattening in the anterior part.

*Distribution:* Hess formation (Glass Mountains) at 1, 98?, 105 (type), 107, T107, 113?, 120, and 222. Hess or Leonard formations (Glass Mountains) at 15 and Tba. Leonard formation (Baylor Mountains) at 503. Leonard formation (Finlay dome) at 512 and 513.

**LINOPRODUCTUS GIRTYI King, n. sp.**

Pl. XVII, Figs. 3–5

*Description:* Shell very small, transversely elliptical. Pedicle valve gently convex. Hinge-line shorter than the greatest width of the shell, the cardinal angles being obtuse. Ears broad and flattened, ill-defined. Median sinus absent, although the specimen from loc. 23–24 shows near the anterior margin a very slight insinuation. Surface marked by strong, rounded radial lirae, numbering about 8 in the space of 5 mm., and increasing in number by intercalation. The lirae tend to expand and contract irregularly, giving the shell a rough and somewhat crenulated appearance. They are separated by flat striae of equal width. Concentric growth lines, spaced at irregular intervals, cross the lirae.

Dimensions in mm. of three specimens:

| | 1 (loc. 23–24) | 2 (loc. 13) | 3 (loc. Ax) |
|---|---|---|---|
| Length | 9 | 8 | 11 |
| Width | 14 | 14 | 14 |

*Observations:* *L. girtyi* differs from other species of *Linoproductus* of this fauna by its low convexity, small flat ears, short hinge-line, rounded, crenulated lirae with broad, flat striae, and well-marked concentric growth lines.

*Distribution:* Leonard formation (Glass Mountains) at 13, 16?, 23–24, 37, and Ax (type). Word formation (Glass Mountains) at 54 and T240.

**LINOPRODUCTUS NASUTUS King, n. sp.**

Pl. XVII, Figs. 8–9

*Description:* Shell rather small, elongate oval in outline, the hinge-line forming the greatest width of the shell. Pedicle valve strongly inflated, beak enrolled and projecting slightly beyond the hinge-line, ascending at first steeply, and then decreasing in curvature gradually to about mid-length, beyond which the curvature is slight. Transverse curvature very strong, the lateral slopes being steep and the wings flattened. There is a slight undefined median sinus in the visceral area, which dies out to give way to a median fold on the anterior slope. In mature specimens this fold appears as a spout-like projection of the anterior margin. Ears marked by about 5 strong concentric wrinkles, which become obsolete on the edge of the visceral convexity. Brachial valve slightly concave; marked near the hinge by concentric wrinkles like those of the wings of the other valve.

Surface marked by fine radiating lirae, of which from 6 to 12 occupy the space of 5 mm. In places they are small and crowded closely together. Spines rare over most of the surface, but concentrated into a row on the wings.

Dimensions in mm. of several specimens:

| | 1 (loc. 246) | 2 (loc. 247) | 3 (loc. 247) | 4 (loc. 248) |
|---|---|---|---|---|
| Length | 28 | 25 | 25 | 21 |
| Width | 20 | 28 | 23+ | 21+ |
| Convexity | 10 | 10 | 10? | 10? |

*Observations:* This species is characterized by the long hinge, posterior transverse median concavity and anterior convexity of the pedicle valve, sparseness of spines, and slight development of concentric wrinkling. Its closest American relative is *L. insinuatus* (Girty), which is much larger and broader, and lacks the sinus in the posterior region; moreover, *L. insinuatus* has a deep notch in its front margin. It is very close to *L. simensis* (Tschernyschew), but is easily distinguishable by its long hinge, more sharply demarked wings, and more strongly enrolled beak. *L. tenuistriatus* (Verneuil) and *L. schrenki* (Stuckenberg) also belong to the same group, but the former lacks a median fold on the pedicle valve, and the latter is much broader than *L. nasutus*.

*Distribution:* Leonard formation (Glass Mountains) at T119. Word formation (Glass Mountains) at 138, T144, 145, T148, 240, 241, 246 (common; types), 247 (common), and 248.

### LINOPRODUCTUS WAAGENIANUS (Girty)

Pl. XVII, Figs. 10–15

1899 *Productus Cora.* Diener (ex parte), Anthracolithic Fossils of Kashmir, Pal. Ind., ser. 15, vol. 1, pt. 2, pp. 22–23, pl. 1, figs. 12a–c.

1899 *Strophomena analoga.* Diener, *idem*, p. 51, pl. 2, fig. 17.

1908 *Productus waagenianus.* Girty, The Guadalupian Fauna, U. S. Geol. Sur. Prof. Paper 58, pp. 253–254, pl. 12, figs. 6–7a.
Middle Capitan, Capitan Peak (sta. 2926).

1908 *Productus waagenianus* var. Girty, *ibid.*, p. 254.
Delaware Mountain formation, Guadalupe Point (stas. 2903 and 2919).

1910 *Productus eucharis.* Girty, Fauna of Park City Formation, U. S. Geol. Sur. Bull. 436, pp. 28–29, pl. 2, figs. 3–4.
Phosphate beds, Montpelier, Idaho.

1915 *Productus waagenianus.* Diener, Anthracolithic Faunae of Kashmir, Kanaur and Spiti, Pal. Ind., new. ser., vol. 5, mem. 2, pp. 71–73, pl. 6, figs. 18–19; pl. 7, fig. 6.
Zewan beds, Kashmir.

1927 *Marginifera? eucharis.* Girty, in Geol., etc. of Part of Southeastern Idaho, U. S. Geol. Sur. Prof. Paper 152, p. 80, pl. 28, figs. 23–25.
Phosphoria formation, southeastern Idaho.

*Description:* Shell small to medium-sized; very wide, with an extended hinge. Ears pointed, in some shells projecting so far that the rest of the shell is wrapped around them. Pedicle valve strongly convex and much enrolled. Beak inflated, pointed, strongly incurved, slightly projecting. Sinus absent, though some individuals show a slight median flattening. Brachial valve concave, with a narrow geniculated part around the front and sides.

Surface marked by radiating lirae, of which from 7 to 14 occupy the space of 5 mm.; they are low and rounded, and separated by striae as wide as or wider than themselves, and they increase in number by intercalation. On some shells they are irregular in arrangement, probably as a result of interruption of normal growth. The lirae give rise to a few scattered spines; both lirae and striae are crossed by extremely fine concentric lines, about 10 of which are present in the space of 1 mm. The hinge is set with a row of large spines on each side of the beak. On the pedicle valve concentric wrinkles are present on the lateral margins, especially near the ears. Visceral area of the brachial valve marked as far as the geniculation by strong, sharp, regular, rather distantly separated wrinkles, of which the anterior slope is so much steeper than the posterior that in some specimens they appear lamellose. It was this character of the brachial valve that led Diener to identify a specimen from Kashmir as *Leptaena analoga.*

Dimensions in mm. of several specimens:

| | 1 (loc. 46) | 2 (loc. 120) | 3 (loc. 120) | Type of *L. waag.* | Kashmir (Diener) |
|---|---|---|---|---|---|
| Length | 12 | 13 | 11 | 14 | 18.5 |
| Width | 34 | 19+ | 32 | 20+ | 32 |

*Observations:* The great variability of the size of the lirae in this species shows that Girty's distinction of the coarsely lirate form as a variety is not justified. Although Girty failed to compare *L. eucharis*, from the Phosphoria, with *L. waagenianus*, Diener has indicated their apparent identity. There is not a single important detail in which the descriptions and figures of the two differ. In his second figuring of these species, Girty says that *P. eucharis* may be identical with *P. aagardi*, "which probably is a *Marginifera.*" The reason for assigning *L. aagardi* to *Marginifera* is not stated, and it seems to me to be hardly conceivable that it is not a *Linoproductus. L. aagardi* (Toula) appears to differ from *L. waagenianus* in being larger, having a shorter hinge-line, and bearing a median depression in the pedicle valve. According to Tschernyschew* the only distinction between "*P. waageni*" and *L. aagardi* is the finer lirae of the type specimens of the former, a character which he regards as unimportant, since many of the specimens of *L. aagardi* from the Urals and Spitsbergen have just as fine lirae as typical *L. waagenianus.* As here defined, *L. waagenianus* would agree perfectly with *L. aagardi* in the nature of its liration. The great length of the hinge of the Texas specimens is, however, an important difference.

*Distribution:* Hess formation (Glass Mountains) at 8, 17a, T205, and Tt. Leonard formation (Glass Mountains) at 12, 25, 37, 39?, 83, 120, T120, 123, 125, 232, D1, D11, D12b, and Tb. Word formation (Glass Mountains) at 45?, 46, 54, 144?, Tx148, 232, 240, T240, and 241. Bone Canyon member (Delaware Mountains) at 490. Delaware Mountain formation (Sierra Diablo) at 504. Leonard formation (Shafter) at 518. Limestone of Cerro Caballo (Las Delicias) at section 1, beds 11 and 14.

### LINOPRODUCTUS (CANCRINELLA) PHOSPHATICUS (Girty)

Pl. XVII, Figs. 6–7

1910 *Linoproductus phosphaticus.* Girty, Fauna of the Phosphate beds of the Park City Formation, U. S. Geol. Sur. Bull. 436, pp. 29–30, pl. 2, figs. 7–9.
Phosphate beds of the Park City formation near Montpelier, Idaho.

*Description:* Most of the specimens in the present collection are so distorted that the actual configuration is not apparent. The following is taken from Girty's description: Shell "broad, slightly transverse or slightly elongate in different specimens. Hinge-line about as wide as the width below. The cardinal angles subquadrate. Ventral valve strongly convex, with small depressed ears. Beak inflated and much incurved. Dorsal valve similar to the ventral, but with lower convexity and smaller and less projecting beak. Surface marked by fine radiating lirae, 8 to 10 in 5 mm. The whole surface of both valves is crossed by moderately strong, subequal but more or less irregular transverse wrinkles, which are stronger and coarser on the ears. There are also small spines, which appear to be rather numerous and regularly distributed. The development of a spine may be accompanied by an elevation of the rib from which it springs, beginning some distance above the point of appearance. In such cases the presence of spines is a conspicuous feature, but otherwise

*Rept. 2nd Norw. Arc. Exp. in the "Fram," vol. 4, no. 34, pp. 37–38. 1916.

not, since they are of about the same size as the costae and do not produce very considerable enlargements."

Dimensions in mm. of several crushed specimens:

| | 1 (loc. 247) | 2 (loc. 247) | 3 (loc. 246) |
|---|---|---|---|
| Length | 27 | 26 | 23 |
| Width | 20 | 26 | 20 |

*Observations:* The most closely related American species are *L. villiersi* and *L. pertenuis*, which have finer lirae, wrinkles, and spines. *L. phosphaticus* is very similar to *L. cancriniformis* (Tschernyschew), a widespread Artinsk species, and may actually prove to be identical with it. The Glass Mountains specimens, in fact, though somewhat crushed, appear to agree somewhat more closely in convexity with the Russian than with the American specimens. If the name *Cancrinella*, introduced by Fredericks for the group of *L. cancrini*, is to be used, this species and *L. villiersi* would be so designated.

*Distribution:* Leonard formation (Glass Mountains) at T119. Word formation (Glass Mountains) at 46, 246, and 247. Leonard formation (Shafter) at 512.

### LINOPRODUCTUS (CANCRINELLA) VILLIERSI (d'Orbigny)

Pl. XVII, Figs. 1–2

1842 *Productus villiersi.* D'Orbigny, Voy. dans l'Amér. mérid., part 4, vol. 3, p. 53, pl. 4, figs. 12–13.

1914 *Productus villiersi.* Kozlowski, Ann. de Pal., vol. 9, pp. 42–44, text fig. 12, pl. 2, figs. 39–60. See for synonymy.

*Description:* Shell small, strongly convex, hemispherical. Hinge-line slightly shorter than the greatest width of the shell. Surface of both valves marked by fine radial lirae, which number about 15 in the space of 5 mm., by fine, more or less irregular wrinkles on both valves, and by numerous spines, which spring from elongated bases.

Dimensions in mm. of three specimens doubtfully identified:

| | 1 (loc. 3) | 2 (loc. 26s) | 3 (loc. 26s) |
|---|---|---|---|
| Length | 14 | 11 | 13 |
| Width | 12 | 12 | 15 |

The specimen from loc. 3 is very narrow in proportion to length.

*Observations:* It is very likely that this species should include also *L. pertenuis* (Meek) and *L. clarkianus* (Derby). The only difference that can be made out between *L. villiersi* and the former is that in the first the spines are nearly prostrate, while in *L. pertenuis* they are more erect.

*Distribution:* Wolfcamp formation *Uddenites* zone (Glass Mountains) at 88. Wolfcamp formation (Glass Mountains) at 75, 193?, and 195. Middle Gym formation (Hueco Mountains) at 414 and 424.

### Genus STRIATIFERA Chao

*Diagnosis:* Shell productoid, generally relatively narrow, with the hinge-line shorter than the greatest width of the shell. Pedicle valve strongly to slightly convex, brachial valve concave and following the curve of the opposite valve, forming a thin but uniform visceral cavity between them through the whole length of the shell. Surface marked by fine radiating lirae and generally also by concentric wrinkles.

Genotype: *Mytilus striatus* Fischer de Waldheim.

*Geologic range:* Mississippian to Permian.

*Observations:* As originally defined, this genus included also the *Productus giganteus* group, but Chao later separated that group as *Kansuella*, which is characterized by its thick shells of large size and transverse outline. Fredericks included *Striatifera* in his genus *Cora* (the other member of that genus being *Linoproductus).*

### STRIATIFERA PINNIFORMIS (Girty)

Pl. XVII, Figs. 16–19

1908 *Productus pinniformis.* Girty, The Guadalupian Fauna, U. S. Geol. Surv. Prof. Paper 58, p. 272, pl. 12, figs. 5–5b.

Middle Capitan, Capitan Peak (sta. 2926).

*Description:* Shell medium-sized, spatulate. Posterolateral margins more or less straight, diverging at an angle of 90° or less; greatest width at about mid-length. The lateral margins merge with the anterior in a broad curve. Beak of the pedicle valve pointed and prominent, bent down at the end. Hinge-line very short. Ears small, provided with several spine-bases near the margin. Pedicle valve longitudinally very slightly convex, except in some specimens in which either the anterior or the umbonal part has a greater convexity. Median sinus or flattening quite absent. Brachial valve concave and slightly wrinkled. Visceral cavity thin.

Surface marked by rather strong, irregular concentric wrinkles and very fine rounded radiating lirae, of which from 8 to 15 occur in the space of 2 mm. The latter are of varying size—on some specimens there are alternate large and small lirae, increasing in number by intercalation. They are larger near the anterior margin. The lirae are crossed at right angles by closely spaced finer concentric lines, which are distinguishable only on well preserved surfaces. The concentric wrinkles curve back toward the wings, where they become closely concentrated.

Dimensions in mm. of several specimens:

| | 1 (loc. 37) | 2 (loc. 34) | 3 (loc. 21) | Ls. of C. Cab. |
|---|---|---|---|---|
| Length | 30 | 19? | 28 | 30+ |
| Width | 24 | 20 | 28 | 37 |

The specimens from loc. 26s have conspicuous scattered spines on the surface.

*Observations:* The fact that this species occurs only in the Hess and Lower Leonard in the Glass Mountains, horizons much lower than that of typical *S. pinniformis*, suggests that they may be a different species, but no evidence of this could be seen in the present collection, in which most specimens of the species are rather incomplete.

The nearest relatives to this species are *S. compressa* (Waagen) and *S. mytiloides* (Waagen), which Chao regards as identical. It differs from them in being less elongate, in having the concentric wrinkles extending upon the wings, and in having fewer and less prominent spines on its wings. *S. compressa corniformis* Chao is still narrower, and has incurved sides.

*Distribution:* Hess formation (Glass Mountains) at 1 and 8. Leonard formation (Glass Mountains) at 16, 21, 26s?, 34, 37, 38, T41, 120, and Tb. Word formation (Glass Mountains at 137. Bone Canyon member (Delaware Mountains) at 491. Limestone of Cerro Caballo (Las Delicias) at section 1, beds 11 and 14. Girty's type was from the Capitan limestone.

#### Genus PUSTULA Thomas

*Diagnosis:* Shell productoid. Pedicle valve strongly convex, brachial valve flattish to slightly concave, generally upturned at the margins, so that there is a thick and uniform visceral cavity the entire length of the shell. Hinge-line less than the greatest width of the shell. Median sinus well-defined. Surface marked by conspicuous concentric bands or wrinkles, bearing one or more rows of spines.

Genotype: *Producta pustulosa* Phillips.

*Geologic range:* Mississippian to Permian.

*Observations:* Chao and Fredericks believe that the type of *P. punctata* should be distinguished as *Echinoconchus*, the distinction being that in the *P. pustulosus* group there are numerous, well marked wrinkles, each of which bears a single row of pustules on the crest, while in the *P. punctatus* group there are strong, well defined bands, each bearing several rows of small pustules. I can hardly agree to such minute splitting.

#### PUSTULA SEMIPUNCTATA (Shepard)

1915 *Pustula semipunctata.* Girty, Pennsylvanian Series in Missouri, Mo. Bur. of Geol. and Mines, vol. 13, ser. 2, pp. 349–350.

1928 *Productus punctatus* and *Pustula nebraskensis.* Schuchert, Am. Jour. Sci., ser. 5, vol. 14 (1927), p. 391.

*Observations:* A single imperfect specimen from the upper Hess formation cannot be distinguished from the common Pennsylvanian form in its visible characters. Like the American form *P. semipunctata* it is elongate and has a noticeable sinus. The lower Wolfcamp formation contains specimens unquestionably belonging to this species.

*Distribution:* Wolfcamp formation, *Uddenites* zone (Glass Mountains) at 88, 88s, and 175. Wolfcamp formation (Glass Mountains) at 93s, T93, and 169a. Hess formation (Glass Mountains) at 102.

#### Genus BUXTONIA Thomas

*Diagnosis:* Shell productoid, with a thick visceral cavity. Pedicle valve evenly convex, or rarely geniculated; brachial valve flattened or concave. Surface spinose, the spine bases being arranged to form more or less interrupted radiating costae, which may become obsolete in the gerontic stage, being replaced by concentric bands on which are numerous small spine-bases. Marginal ridges short. Median septum generally bifurcating about 5 mm. below the cardinal process, the two branches uniting with the two lobes of the process.

Genotype: *Anomites scabriculus* Martin.

*Geologic range:* Mississippian to Permian.

*Observations:* The diagnosis given above differs somewhat from those of Thomas, Muir-Wood, and Chao. The arrangement of the spine-bases as costae is here emphasized, rather than the pustulose bands in the gerontic stage, which have been observed only in the genotype. *Productus nebraskensis*, which has been regarded as either a *Pustula* or a *Buxtonia*, agrees best with *Juresania* Fredericks.

Fredericks in 1923* proposed the subgeneric name *Tschernyschewiella* for productids having a median septum or "euseptoid" in the pedicle valve. The type species was *Productus porrectus* Kutorga, which, according to Kozlowski, is synonymous with *P. peruvianus* d'Orbigny. Whether the presence of a low median septum is as important as believed by Fredericks is very doubtful, as many species have a more or less well developed median ridge. At any rate, the name *Tschernyschewiella* had been preoccupied by von Toll for a Siberian Cambrian trilobite, and its use for a brachiopod is incorrect. It seems logical to suppose that *P. porrectus* (or *P. peruvianus)* is a species of *Buxtonia* and deserves no separate designation.

#### BUXTONIA OCCIDENTALIS King, n. sp.

Pl. XVII, Figs. 20–21

1858 *Productus scabriculus* Marcou, Geol. of North America, p. 47, pl. 5, fig. 6.

1861 *Productus scabriculus.* Newberry, Ive's Rep. Colorado River of the West, p. 125.

1903 *Productus sp. a.* Girty, Carb. Formations and Faunas of Colorado, U. S. Geol. Sur. Prof. Paper 16, p. 372, pl. 5, fig. 3.
Middle and upper Hermosa formation, San Juan region.

1909 *Productus nebraskensis.* Girty, Manzano Group of Rio Grande Val., U. S. Geol. Sur. Bull. 389, pp. 62–64, pl. 7, figs. 5–6.
Abo sandstone: Abo Canyon, Mesa del Yeso, and Sandia Mountains. Yeso formation: Mesa del Yeso.

*Description:* Shell medium-sized, subtriangular in outline, hinge-line less than the greatest width, ears small and depressed. Pedicle valve strongly and regularly inflated in longitudinal profile. Beak pointed and somewhat incurved over the hinge-line. Median sinus broad and shallow, not well-marked. Surface of the pedicle valve marked by elongate spine bases, which are not connected end to end to form continuous costae, as typically in the species of this genus, but generally lie somewhat out of line with one another. The spine bases are so long and so closely set to one another that the shell appears to be costate when not carefully examined. There are several concentric undulations in the surface of some specimens, but they have no relation to the arrangement of the spines. Fine growth lirae may be distinguished in some specimens. Surface of the brachial valve marked by radially elongate pits and by faint concentric wrinkles.

Dimensions in mm. of several specimens:

| | 1 (loc. 75) | 2 (loc. 75) | 3 (loc. 323) |
|---|---|---|---|
| Length | 21 | 25 | 38 |
| Width | 27 | 30 | 30 |
| Convexity | 10 | 13 | 12 |

*Observations:* This species appears to occupy an intermediate position between the Lower Carboniferous *Buxtonia*

*Rec. Geol. Com. Russian Far East, no. 28, pp. 20–21.

*scabricula* and *Juresania juresanensis*, having, unlike the former, interrupted rather than continuous costae, and, unlike the latter, no concentric arrangement of the spines, and but a single set of spines. The difference from *J. nebraskensis* is even greater, as the common mid-continent species has very strong concentric corrugations and many irregularly arranged spines of very diverse sizes.

*Distribution:* Gaptank formation (Marathon basin) at 75 (types), 272, 275, 284, Sa*, Sd, and Sh. Strawn group (Hueco Mountains) at 323 and 329. Wolfcamp formation (Glass Mountains) at 91a, T91, 93, and 203. A specimen from the basal Magdalena of the Sandia Mountains is identical with the Texas form, as well as specimens from the Pennsylvanian of Northern Arizona. The specimens identified as *P. scabriculus* by Newberry and Marcou were from near Santa Fe and from the Sandia Mountains, and there seems to be little room for doubt that they should actually be placed in *B. occidentalis*.

#### BUXTONIA PERUVIANA (d'Orbigny)

Pl. XVIII, Figs. 1–3

1842 *Productus Peruvianus*. D'Orbigny, Voy. dans l'Amér. mérid., p. 52, pl. 4, fig. 4.

1844 *Productus porrectus*. Kutorga, Zweiter Beitrag zur palaeont. Russlands, Verh. d. Russ. Kais. Mineral. Gesellsch., ser. 1, p. 26, pl. 10, fig. 3.

1914 *Productus peruvianus*. Kozlowski, Ann. de Pal., vol. 9, pp. 38–40, text fig. 7, pl. 5, figs. 1–4. (See for synonymy.)

1916 *Productus Porrectus*. Tschernyschew and Stepanow, Rept. Sec. Norw. Arctic Exped. in the "Fram," no. 34, p. 41, pl. 5; pl. 8, fig. 5.

1924 *Productus (Tschernyschewiella) porrecta*. Fredericks, Rec. Geol. Com. Russ Far East, no 28, p. 21.

*Description:* Shell large, triangular in outline. Pedicle valve strongly convex near the beak, but with gradually decreasing curvature toward the front. Beak greatly incurved over the hinge-line. Margins of the visceral area very steep. Ears small, slightly convex. Median region depressed in a broad sinus (which, however, is absent in one specimen from loc. Tmm). Brachial valve nearly flat or only slightly convex, with a low median fold; geniculate near the front. Surface marked by costae, which are continuous but consist of rows of elongate nodes, between which the costae become very faint. The nodes on the costae are the bases of small erect spines. Concentric wrinkles cross the pedicle valve of the specimen from loc. 472.

Dimensions in mm. of several specimens:

| | 1 (loc. 472) | 2 (loc. Tmm) | 3 (loc. 377b) |
|---|---|---|---|
| Length | 50+ | 48 | 58 |
| Width | 52 | 54 | 55 |
| Convexity | 51 | 22 | 30 |

*Observations:* There is no apparent difference between these specimens and typical *B. peruviana* as described by Kozlowski.

*Identified by Schuchert (Am. Jour. Sci. ser. 5, vol. 14, 1927), as *Pustula nebraskensis*.

*Distribution:* Lower Gym formation (Hueco Mountains) at 377b, 403, and Tmm. Hess formation (Sierra Diablo) at 472.

#### BUXTONIA VICTORIOENSIS King, n. sp.

Pl. XIX, Fig. 1

*Description:* Shell large, transversely subquadrate, visceral region subtriangular. Pedicle valve strongly and evenly convex, beak incurved over the cardinal area. Margins of the visceral area rather steep. Ears broad, convex, sharply set off. Median region depressed in a broad, shallow sinus. Surface marked by continuous radial costae, numbering about 9 in the space of 10 mm. They swell at intervals into small nodes, which are spine bases. Concentric wrinkles are absent.

Dimensions in mm. of the type specimen, from loc. 498: length, 50; width, 70; convexity, 25.

*Observations:* This species is distinguished from *B. peruviana* by being much broader, the greatest width being at the hinge-line, having broad, well-defined triangular ears, relatively finer costae and smaller nodes, and stronger convexity. It is much smaller and less strongly convex than *B. longa*.

*Distribution:* Victorio Peak member (Sierra Diablo) at 498 (type). Leonard formation (Glass Mountains) at 26s. Word formation (Glass Mountains) at 243? and 255?. Rare.

### Genus WAAGENOCONCHA Chao

*Diagnosis:* Shell productoid, pedicle valve of low convexity, brachial valve nearly flat. Surface marked by quincuncially arranged tubular spines in the youthful and adult stages, which become smaller and closely packed together in old age. Wrinkles faint or absent.

Genotype: *Productus humboldti* d'Orbigny.

*Geologic range:* Uppermost Pennsylvanian and Permian.

*Observations:* In 1928, the year following the publication by Chao of his diagnosis of *Waagenoconcha*, Fredericks introduced the name *Ruthenia* for approximately the same group, the genotype being *P. irginae* Stuckenberg. Besides the *P. irginae* group, however, he included that of *P. tastubensis*, which is regarded by Chao as an *Avonia*. Other species classed as *Waagenoconcha* by Chao were placed by Fredericks in his genus *Krotovia*. The distinction by Fredericks between *Ruthenia* and *Krotovia* is that the former is ornamented with fimbriate tubes, and the latter with spines only. In each genus are placed shells having either plane or concave dorsal valve. To me the division by Chao seems the more natural, the shells in both of Fredericks' "genera" having a plane dorsal valve being placed in *Waagenoconcha*, while those having a concave dorsal valve are regarded as *Avonia* (or *Krotovia*, as defined by Chao).

#### WAAGENOCONCHA LEONARDENSIS King n. sp.

Pl. XIX, Figs. 2–4

*Description:* Shell large, subquadrate, with the hinge-line slightly less than the greatest width. Pedicle valve strongly inflated, especially near the beak, and flattening out in longitudinal profile toward the front. Lateral slopes very steep in early stages and somewhat less so later. Beak pointed, enrolled, and incurved over the hinge. Ears

flattened, demarked from the rest of the shell by the concavity bordering the visceral area. Median sinus originating at the very point of the beak or slightly anterior to it, becoming strongest in the posterior median part of the shell, and becoming shallower toward the anterior margin. Surface marked by many large, elongated pustular spines, more or less arranged along two diagonal axes intersecting at a low angle.

Dimensions in mm. of several specimens:

| | 1 (loc. 15) | 2 (loc. 120) | 3 (loc. 38) | 4 (loc. Tb) |
|---|---|---|---|---|
| Length | 35 | 45 | 33 | 43 |
| Width | 43 | 50 | 35 | 42 |
| Convexity | 21 | 20 | — | 15 |

*Observations:* This species is easily distinguishable from *W. montpelierensis* by its greater convexity, its deep median sinus, and especially its large relative size and the irregularity in the arrangement of its spines. It differs from *W.? palliata* (Kayser) in having larger spines and a deeper sinus. *W. abichi* and *W. serialis* (Waagen) are similar to it in the size and arrangement of their spines, but are much smaller and have much shallower sinuses. *W. gangeticus* (Diener) also appears to be similar, but probably has smaller spines.

*Distribution:* Hess? formation (Glass Mountains) at 38 and Tv. Leonard formation (Glass Mountains) at 15, 22, 83, 120 (type), T233, and Tb.

### WAAGENOCONCHA MONTPELIERENSIS (Girty)

Pl. XIX, Figs. 5–6

?1898 *Productus irginae.* Stuckenberg, Carte géol. de la Russie, f. 127, p. 220, pl. 2, fig. 16.

1910 *Productus montpelierensis.* Girty, Fauna of the Park City Formation, U. S. Geol. Sur. Bull. 436, p. 30, pl. 2, figs. 5–6.

?1914 *Productus humboldti* var. *irginae.* Kozlowski, Brachiopodes du Carb. super. de Bolivie, Ann. de Pal., t. 9, pp. 41–42, text fig. 10, pl. 7, figs. 1–6.

1927 *Productus montpelierensis.* Girty, in Geog., etc., of Southeastern Idaho, U. S. Geol. Sur. Prof. Paper 152, p. 80, pl. 28, figs. 12–13.
Phosphatic shale member of the Phosphoria formation, southeastern Idaho.

*Description:* Shell medium-sized to large, transversely oval in outline. Hinge-line shorter than the greatest width, cardinal angles rounded. Profile of the pedicle valve regularly convex. Visceral area triangular, with steep slopes, the umbonal angle being about 90°. Beak small, strongly incurved, rapidly expanding. Ears small and depressed. Median sinus shallow and regularly rounded, originating about 10 mm. in front of the beak and becoming less distinct toward the front, where it is hardly at all distinct. Brachial valve flat in the postero-central part, concave anteriorly and laterally, with a low undefined fold corresponding to the sinus of the other valve.

Surface of both valves ornamented with small, short, sharp, tubular spines, directed forward. On the pedicle valve these spines are arranged in intersecting diagonal lines, but in places, in irregular transverse lines bearing closely spaced spines. On the brachial valve the spines are arranged in very regular qunicunx. On the best specimen there is one faint growth varix near the anterior margin.

Internal characters are not determined.

Dimensions in mm. of specimen from loc. 146: length, 34; width, 43; convexity, 15.

*Observations: W. montpelierensis* shows no apparent difference from *P. humboldti irginae* as figured by Kozlowski, and only minor differences can be made out between it and *P. irginae* figured by Tschernyschew: a lower convexity and blunter and more strongly inflated beaks. Fredericks* entertains no doubt as to the actual identity of the Eurasian and American species. If *W. montpelierensis* is a different species from *W. irginae,* the question arises as to whether they differ according to time or according to place: are all the American forms *W. montpelierensis* and all the Eurasian forms *W. irginae,* or are the high Pennsylvanian and low Permian forms *W. irginae in Eurasia, Bolivia,* and the *Uddenites* zone of the Wolfcamp formation, and the middle Permian forms *W. montpelierensis* in the Phosphoria, the Leonard, and the Word? These questions cannot be settled now, because of the imperfect nature of the material from the Wolfcamp formation. All the specimens in the present collection will therefore be regarded as *W. montpelierensis,* although it is considered extremely likely that if that species is different from *W. irginae,* the form occurring in the *Uddenites* zone is *W. irginae. W. montpelierensis* appears to be a characteristic species of the North American Arctic Permian fauna. The *Pustula* aff. *P. humboldti* cited from the Alaskan Artinsk fauna is, as suggested by Girty*, possibly identical with *W. montpelierensis.* The specimens identified as *Pustula* aff. *P. irginae* and *Pustula* aff. *montpelierensis* by Girty* from the Kaibab in Arizona are almost certainly identical with this species. It is interesting to note that *W. irginae* is described by Tschernyschew and Stepanow as occurring abundantly in the lower Permian limestone of Store Björnekap, Ellesmere Island, in Arctic America. In Eurasia the species is restricted to Russia and Siberia, and does not occur in the faunas of the Tethyan region. From these facts it appears that *W. irginae* and *W. montpelierensis* are characteristically a northern type rather than a Tethyan one.

*W. montpelierensis* and *W. irginae* are easily distinguishable from other members of the genus by the extreme fineness and regularity of arrangement of their spines. Typical *W. humboldti,* for example, has long spines with much less regular quincuncial arrangement. Another closely related species is *W. waageni* (Rothpletz), from the East Indies, which has a deeper and narrower sinus, a more strongly enrolled beak, and concentric undulations of growth. It is certainly much closer to *W. irginae* than to *W. humboldti,* with which Broili compared it.

*Distribution:* Wolfcamp formation, *Uddenites* zone (Glass Mountains) at T88. Leonard formation (Glass Mountains) at 3, 13, T119, 120, 124, and 174. Word formation (Glass Mountains) at 146, T148, T240, and 247. Victorio Peak member (Sierra Diablo) at 496. Limestone of Cerro Caballo (Las Delicias) at section 1, bed 14.

---

*Mem. Com. Geol., n. s., liv. 193 (1915), p. 29.

*U. S. Geol. Sur. Prof. Paper 152, p. 79.

*In Darton, N. H.: Resumé of Arizona Geology, pp. 103–104.

### Genus OVERTONIA Thomas

*Diagnosis:* Shell small, productoid. Surface of the pedicle valve marked by concentric elevated bands, each of which bears a single row of relatively large spine bases. Within the brachial valve the brachial ridges, instead of originating on each side of the muscular impression in a direction practically at right angles to the median septum, extend for almost the whole of their length at an angle of approximately 45° to the septum. The anterior ridge in each pair originates at the inner side and near the anterior end of the median septum and just in front of the roughly oval muscular area. It extends forward for a distance equal to half the length of the valve, and at its anterior limit it curves around in a postero-lateral direction, and then backward in a course nearly subparallel to the original direction, ending at the margin of the muscular area at about mid-length of the latter. The two branches of each pair of ridges become slightly nearer to each other at about the middle of their course, so that each pair of brachial ridges resembles a ladle with a short, broad handle.

Genotype: *Productus Simbriatus* J. Sowerby.

*Geologic range:* Mississippian to lower Permian.

*Observations:* Only two species have been recognized as belonging to this genus: *O. fimbriata* (Sowerby), the genotype, and *O. cristato-tuberculata* (Kozlowski). It is possible that *Pustula biseriata* (Hall), of the Mississippian Spergen formation, is an *Overtonia*, but this cannot be determined, for, though Weller has figured the interior of the brachial valve (Mississippian Brachiopoda, plate 17, fig. 15), the details of its structure cannot be seen in the figure, and none of the internal characters are described.

#### OVERTONIA CRISTATO-TUBERCULATA (Kozlowski)?

Pl. XX, Figs. 1–3

1914 *Productus cristato-tuberculatus.* Kozlowski, Brachiopodes du Carb. Supér. de Bolivie, Ann. de Pal., vol. 9, pp. 45–46, pl. 2, figs. 61–62.

*Description:* Shell small, ovoid, with the hinge-line slightly less than the greatest width of the shell. Pedicle valve regularly and strongly inflated. Beak pointed and incurved. Ears small and not very distinct. Median sinus absent. Surface marked by about 10 regular, obtusely angulated concentric ridges, steepest on the anterior sides, separated by broad, shallow depressions. These ridges are crowded together on the sides of the shell. Between the anterior face of each ridge and the crest of the next succeeding one there are long parallel spine-bases, evidently bearing forward-projecting spines. A few of the spine-bases are prolonged as costae marking the anterior faces of the ridges and connecting with the spine-bases of the next succeeding ridge. Brachial valve slightly and uniformly concave and marked as the pedicle valve. Length, 12 mm.; width, 12 mm.

*Observations:* These specimens differ from those figured by Kozlowski, so far as can be seen, only in a slightly greater size, the largest of those from Bolivia being 9 or 10 mm. long. Since in the Wolfcamp specimens the internal characters of the brachial valve cannot be determined, there is some doubt as to their generic reference. If they are, as appears, identical with the Bolivian species, then their generic reference is certain, for the internal characters of the brachial valve, as worked out by Kozlowski, are like those of *O. fimbriata*, the genotype.

*Distribution:* Wolfcamp formation, *Uddenites* zone (Glass Mountains) at 88, 94, 94s, and Tba. The species is present also in a collection from the Wayland shale member of the Cisco, 1.5 miles southeast of Gunsight, Texas.

### Genus AVONIA Thomas

*Diagnosis:* Shell productoid, with concave brachial valve. Surface spinose in the first stages, developing costae later, through elongation of the spine bases. Marginal ridges short, diverging from the hinge-line. Cardinal process small. Median septum narrow. Adductor muscle-scars not dendritic.

Genotype: *Productus youngianus* Davidson.

*Geologic range:* Mississippian to Permian.

*Observations:* As here used, the name *Avonia* is applied to species with a variety of surface sculptures, some of them being marked exclusively by tubercles, and others having the tubercles give rise to a rather regular costation. This is the sense in which the name was used by Chao in part I of the "Productides of China." In part II of that work the greater part of the group was transferred to Fredericks' genus *Krotovia*. There is probably some justification in restricting the genus *Avonia* to species which, like the genotype, are spinose only in the visceral region and are evenly costate over the rest of the surface, but the differences between the typical form and such completely spinose species as *A. signata* seem to be more of degree than of kind. The species assigned to *Avonia* according to the present definition were grouped by Fredericks (1928) in *Ruthenia*, *Krotovia*, and *Thomasia*, the subdivision being on an entirely different basis than that of Chao.

#### AVONIA BOULEI (Kozlowski)

Pl. XX, Figs. 7–9

1914 *Productus boulei.* Kozlowski, Brachiopodes du Carb. Supér. de Bolivie, Ann. de Pal., vol. 9, pp, 47–48, text fig. 13, pl. 3, figs. 8–9.

1914 *Productus* aff. *spinulosus.* Douglas, Geological sections through the Andes, Quart. Jour. Geol. Soc., vol. 70, pp. 33–34.

*Description:* Shell small to medium-sized, outline subtriangular except for the ears; greatest width of the shell at the hinge-line. Pedicle valve strongly and evenly convex, beak incurved and projecting somewhat over the hinge-line. Borders of the visceral area diverge at about 90° from the beak, but become nearly parallel toward the front; they are very steep near the beak, and gradually decrease in slope forward. Ears well defined, slightly convex. Median region generally flattened, but not depressed into a sinus. Brachial valve deeply concave. Surface of the pedicle valve marked by large elongate spine bases, increasing in size from the beak toward the front. They are especially abundant on the slopes of the visceral area. Brachial valve marked by pits which are the reverse of the spine bases of the other valve.

Dimensions in mm. of several specimens:

| | 1 (loc. 193) | 2 (loc. 193) | 3 (loc. 95) | 4 (loc. 389) |
|---|---|---|---|---|
| Length | 20 | 14 | 26 | 18 |
| Width | 24 | 17 | 33 | 19 |
| Convexity | 11 | 7 | 15? | 11 |

*Observations:* The Texas specimens resemble in every way those from Bolivia, except that some of them attain a greater size. The most similar American species is *A. subhorrida*, but this is distinguishable by the larger size, stronger convexity, greater length in proportion to width, and the presence of a median sinus. It is even more similar to *A. tuberculata* (Moeller), but differs therefrom in its smaller spines and the absence of a sinus. *A. wallaciana* (Derby) is similar, the only apparent difference being that the Brazilian species has smaller spines. It is quite likely that the specimens identified by Meyer* as *Productus wallacei* are actually *A. boulei*, for they were collected in Peru from a fauna like that of Bolivia in which the typical form of *A. boulei* was found.

*Distribution:* Wolfcamp formation, *Uddenites* zone (Glass Mountains) at 95. Wolfcamp formation (Glass Mountains) at 87 low, 93, 93s, and 193. Lower Gym formation (Hueco Mountains) at 377b, 389, 390, 392, and 4 miles southwest of Hueco Tanks (Baker coll.). Hess formation (Sierra Diablo) at 472.

#### AVONIA? INCURVATA King, n. sp.

Pl. XX, Fig. 6

*Description:* Shell small, subquadrate, greatest width at the hinge-line. Pedicle valve strongly arched, the umbo being greatly elevated. Beak large and very much incurved over the hinge-line. Ears extended and fairly well defined. Median region faintly depressed into a sinus. Brachial valve concave, ears flattened. Surface marked by broad, low, rounded costae, separated by shallow furrows; they arise some distance anterior to the beak, and bear scattered large erect spines. There are uneven concentric undae, some of which, near the front margin, are especially strong. It is possible that this species is a *Marginifera*.

Dimensions in mm. of the typical specimen: length, 16; width, 22; convexity, 9.

*Observations:* This species is distinguished by its triangular outline, strongly incurved beak, large ears, and extended hinge-line.

*Distribution:* Word formation (Glass Mountains) at 46 (rare).

#### AVONIA LATIDORSATA (Girty)?

1907 *Productus latidorsatus.* Girty, the Guadalupian Fauna, U. S. Geol. Surv. Prof. Paper 58, pp. 264–265, pl. 11, figs. 11–13b.

*Description* (after Girty): "Shell rather small, thin, transverse. Ventral valve moderately convex, the anterolateral curvature being stronger than the transverse. Beak small, incurved, not projecting far beyond the hinge-line. Ears small, depressed, somewhat arched, probably slightly projecting. Sinus shallow, ill defined. Dorsal valve moderately convex, with flattened visceral region. Beak inconspicuous. Ears small, depressed, somewhat arched, and slightly projecting.

"Surface nearly smoooth. Traces of ribs can be seen on most specimens, but they are usually faint. Equally or somewhat more indistinct concentric wrinkles also are found, and rather strong growth lines. Spines are numerous and small over the posterior portion of the shell, rather scarce and larger over the front, lateral portions, and ears. The ribs in many cases, especially where at all well marked, are more or less discontinuous, and originate severally at the bases of the larger spines."

Dimensions in mm. of the Word specimen: length, 13; width, 22; convexity, 9.

*Observations:* In the present collection there are but two specimens referable to this species, and since they are incomplete, the identification is not certain. The species is distinguished from related ones in this fauna by being short and broad, having a median sinus, and bearing few spines.

*Distribution:* Word formation (Glass Mountains) at 241.

#### AVONIA MEEKANA (Girty)

Pl. XX, Figs. 4–5

1908 *Productus meekanus.* Girty, The Guadalupian Fauna, U. S. Geol. Sur. Prof. Paper 58, p. 263, pl. 30, figs. 13–13a.
Delaware Mountain formation, Guadalupe Point (sta. 2931); Delaware Mountain formation, Glass Mountains (sta. 3763).

*Description:* Shell small. Pedicle valve moderately convex. Beak small. Hinge-line shorter than the greatest width of the shell; ears small and not very sharply set off. Sinus absent, although one specimen has a median flattening of the posterior part of the shell. Surface marked by concentric wrinkles and spines. Wrinkles small and irregular, strong on the sides and ears but fainter and somewhat interrupted elsewhere. Spines medium-sized, numerous, and regularly arranged 1.5 to 2 mm. apart, projecting strongly downward and forward so that their bases are very elongate. Where the surface is worn, there appear radiating lirae, which are faint and interrupted by the spines; about 3 of them occupy the space of 1 mm.

Dimensions in mm. of one specimen: length, 15; width, 16; convexity, 8.

*Observations:* This species is very similar to the more common *A. signata*, but has much more closely arranged and smaller spines. The only foreign species to which it is closely similar is *A. pustulata* (Keyserling), which has similar ornamentation but a quite different configuration, being much wider and more convex.

*Distribution:* Word formation (Glass Mountains) at 144 and Tc (rare).

#### AVONIA SIGNATA (Girty)

Pl. XX, Figs. 16–24

1908 *Productus signatus.* Girty, The Guadalupian Fauna, U. S. Geol. Sur. Prof. Paper 58, pp. 263–264, pl. 22, figs. 4–4b.

1908 *Productus signatus* var. Girty, *idem.*, p. 264.
Delaware Mountain formation, Guadalupe Point (sta. 2931).

*Meyer, H. L. F.: Carbonfaunen aus Bolivia und Perú, N. Jahrbuch, vol. 37, p. 632. 1914.

1908 *Productus* sp. e. Girty, *idem*, p. 266, pl. 22, figs. 5–5a. Delaware Mountain formation, Guadalupe Point (sta. 2920).

*Description:* Shell medium-sized, generally slightly longer than wide, rarely somewhat wider than long, with the hinge-line narrower than the greatest width of the shell. Pedicle valve moderately arched, the curvature near the beak amounting to 90° of a circle. The convex margins of the visceral area diverge from the beak at right angles or more toward the greatest width of the shell, in front of which place the lateral margins are parallel. Median part flattened, without a sinus, bordered laterally by an abrupt up-turning to the margin. Ears small, depressed, and rounded. Brachial valve regularly concave.

Surface marked by rather regularly arranged spines mounted on narrow elongate bases, which make up a sort of coarse, irregular, and discontinuous ribbing. The abundance of these spines is extremely variable; they are from 1.5 to 6 mm. apart, but generally about 2 mm. apart and evenly spaced at that interval on any one shell. There appear to be all gradations in amount of spacing from the more common fine type to the coarse type. Where the shell surface has been removed it is seen that the spines arising from these elongate bases originate deep within the shell wall and pass up through the bases. They reach a length of at least several millimeters. Surface marked also by fine concentric wrinkles, which are stronger and more closely spaced near the beaks of some specimens, but in others are only slightly developed. Near the anterior margin of some shells the spine bases are prolonged into costae of about the same width as the spine bases, but varying in size on different individuals. The spines on the ears are much larger than those on the body of the shell.

Shell thin. Adductor muscle scars finely labyrinthiform, not situated on a raised platform, but lying at the same level on the shell interior as the diductor muscle scars, which are coarsely striated longitudinally. An interior of a brachial valve doubtfully referable to this species, from loc. 246, has a low, thin median septum. Brachial ridges obscured by pustules, which mark the entire interior.

Dimensions in mm. of several specimens:

| | 1 (loc. 247) | 2 (loc. 144) | 3 (loc. 144) | 4 (loc. 144) | 5 (loc. 253) |
|---|---|---|---|---|---|
| Length | 21 | 24 | 23 | 18 | 22 |
| Width | 20 | 22 | 22 | 21 | 21 |

*Observations:* The average type of the shells referred to this species differ from Girty's type in being generally more elongate, but as certain Glass Mountains specimens agree with the original it cannot be doubted that the two are identical. It is probable that the lirae noted by Girty on an internal mold are merely the striations of the adductor muscle scars, rather than the reflection of surface lirae, for no true lirae are present on any of the specimens here examined. The differences noted by Girty between *A. signata, A. signata* var., and *A.* sp. e appear to come within the limits of this variable species as here defined. *A. signata* is very similar to *A. opuntia* (Waagen), differing from it, according to Girty, in the more elongated character of its spine bases. This feature, however, is variable, many specimens of *A. signata* actually having nearly as short spine bases as the Indian species.

*Distribution:* Word formation (Glass Mountains) at 138, 143?, 144 (abundant), T144, 146, 148, T148, Tx148, 150, 153, 154?, 162, 180, T239, T240, 243, 246, 247, 248, T241, 253, Tc, and Tx. Word? formation (Glass Mountains) at D8 and T91.* Delaware Mountain formation (Sierra Diablo) at 504.

### AVONIA SUBHORRIDA (Meek)

Pl. XX, Figs. 10–11

1877 *Productus subhorridus.* Meek, King's U. S. Geol. Expl. 40th Par., vol. 4, pp. 75–76, pl. 7, figs. 3–3b.
Carboniferous limestone of Wasatch Mountains; Hamilton Butte, Ruby Group; Mahogany Peak, Egan Mountains, and north slope of Moleen Peak, Elko Range, west side of Long Valley, Ruby Group.

1920 *Pustula subhorrida.* Girty, in Ore Deposits of Utah, U. S. Geol. Sur. Prof. Paper 111, p. 655, pl. 56, figs. 1–2 (copied from Meek).
Permian, Utah.
Also cited by Girty (U. S. Geol. Sur. Prof. Paper 152, p. 79) as characteristic of the Phosphoria formation of Idaho, Wyoming, and Montana.

*Description:* Shell medium-sized, transversely ovate, hinge-line slightly less than the greatest width. Pedicle valve very strongly convex, especially in the umbonal region, with a shallow, broad median sinus (there is no sinus in the specimen from T239). Beak pointed and incurved, but not greatly overhanging the hinge-line. Lateral slopes of the visceral area very steep, ears small and sharply set off. Surface set with fairly evenly distributed large scattered spines, the bases of which are to some extent elongated longitudinally. Fine concentric lirae are present also. Brachial valve concave.

Dimensions in mm. of two specimens from loc. Tx148: No. 1, length 20, width 22, convexity, 10; No. 2, length 15, width 18, convexity 9.

*Observations:* The Texas specimens agree with those from the Phosphoria in every respect except that the sinus is not quite as deep in the former.

*Distribution:* Word formation (Glass Mountains) at Tx148 and T239.

### AVONIA SUBHORRIDA RUGATULA (Girty)

Pl. XX, Figs. 12–15

1908 *Productus subhorridus* var. *regatulus.* Girty, The Guadalupian Fauna, U. S. Geol. Sur. Prof. Paper 58, pp. 267–268, pl. 30, figs. 11–12c.
Delaware Mountain formation, Guadalupe Point (sta. 2931?); Glass Mountains (stas. 3763 and 3840).

*Description:* Shell small, transversely subquadrate. Pedicle valve moderately convex. Beak low. Ears small and depressed, hinge-line equal to or less than the greatest

*Loc. T91 is presumably Wolfcamp. There appears to have been an error in the labeling of this specimen.

width of the shell. Median region depressed as a very shallow sinus. Brachial valve gently concave. Surface of the pedicle valve marked by fine, irregular concentric wrinkles, some of which are stronger and of the nature of growth lines. Spines large, long, and scattered; most of them are directed slightly forward or lie perpendicular to the surface. The spine bases give rise to faint costae, which extend forward a short distance, then die out. The interior of the brachial valve lacks marginal ridges, though one specimen shows ridges that lie in the posterior part of the shell, diverging from the base of the cardinal process, as in a *Marginifera* having the anterior part of the marginal ridge undeveloped.

Dimensions in mm. of several specimens:

| | 1 (loc. 137) | 2 (loc. 50) | 3 (loc. 123) | Girty's type |
|---|---|---|---|---|
| Length | 13 | 15 | 13 | 12 |
| Width | 16 | 18 | 13 | 14 |
| Convexity | 8 | 9 | 8 | 7 |

*Observations:* This variety differs from typical *A. subhorrida* in being smaller, and in having the posterior part crossed by fine wrinkles. *A. boulei* is much larger, less strongly convex, and non-sinuate.

*Distribution:* Leonard formation (Glass Mountains) at 31 and 123. Word formation (Glass Mountains) at 44, 46, 50, 54, 137, 138, 150, 240, and 241. Upper Gym formation (Hueco Mountains) at 440a. Limestone of Cerro Cabello (Las Delicias) at section 1, bed 19.

**AVONIA WALCOTTIANA (Girty)**

Pl. XXI, Figs. 1–5

1908 *Productus walcottianus* (ex parte). Girty, The Guadalupian Fauna, U. S. Geol. Sur. Prof. Paper 58, pp. 269–270, pl. 21, figs. 27–27b (not 28–28b).
Delaware Mountain formation, Guadalupe Point (sta. 2903), and southern Delaware Mountains (sta. 2969?).

*Description:* Shell small, probably longer than wide when complete. Pedicle valve strongly convex, geniculate, with a small visceral region. Median region depressed into a rather strong sinus. Ears small and short. Posterior third of the pedicle valve spinose but not costate. Anterior to this region the spine bases are elongate, and give rise to costae, which are large near the spine base but decrease in height in front of it. Other spines tend to arise along the same line, so that discontinous costae are formed, which swell out near the spine bases and contract between them. Visceral region crossed by small but strong concentric wrinkles. One specimen from loc. 46, relatively broad and with a long hinge, has larger and fewer spines than the typical specimens.

Dimensions in mm. of several specimens:

| | 1 (loc. 240) | 2 (loc. 241) | 3 (Girty's type) |
|---|---|---|---|
| Length | 15 | — | 16? |
| Width | 16 | 16 | 20 |
| Convexity | 10 | — | 10 |

*Observations:* Girty tentatively referred his specimens illustrated on his plate 21, figs. 28–28b, to this species. Tschernyschew* expressed the view that it is distinct, and it is now embraced in the new variety *A. walcottiana costata.* The persistent nature of the costae which arise from the spine bases distinguishes *A. walcottiana* from the other species of *Avonia* here described.

*Distribution:* Word formation (Glass Mountains) at 46, 144, 239, T239, 240, T240a, 241, 248, and Tc.

**AVONIA WALCOTTIANA COSTATA King, n. sp.**

Pl. XXI, Figs. 8–13

1908 *Productus walcottianus* (pars). Girty, The Guadalupian Fauna, U. S. Geol. Sur. Prof. Paper 58, pp. 269–270, pl. 21, figs. 28–28b (not 27–27b).
Delaware Mountain formation, Guadalupe Point (sta. 2931).

*Description:* Shell small, narrow, longitudinally subquadrate. Pedicle valve strongly convex, with a very much incurved beak; greatest curvature at about a third the length from the beak to the front margin. Lateral slopes steep; umbonal angle about 90°. Ears small. Beak pointed. Umbonal disc marked by small erect spines and by sharp but fine concentric wrinkles; the long anterior slope covered by rather strong, even, rounded costae, from which small erect spines arise at frequent intervals.

Dimensions in mm. of several specimens:

| | 1 (loc. 241) | 2 (loc. 55) | 3 (Girty's fig. 28) |
|---|---|---|---|
| Length | 21 | 13 | 22 |
| Width | 17 | 13 | 18 |
| Convexity | 12 | 8 | 10 |

*Observations:* These shells agree perfectly with the specimen from Girty's sta. 2931 assigned by him to *P. walcottianus.* Typical *A. walcottiana* is more strongly geniculate than this variety, as well as broader and shorter, and marked by interrupted spine bases rather than continuous costae.

*Distribution:* Word formation (Glass Mountains) at 55?, T240 (type), 241, and Tx. Limestone of Cerro Caballo (Las Delicias) at section 1, beds 12 (SW of the Noria) and 14.

**Genus HORRIDONIA Chao**

*Diagnosis:* Shell productoid, subtriangular in outline. In the pedicle valve there is a strong median sinus. Brachial valve slightly concave. Surface smooth except for a few scattered very stout hollow spines.

Genotype: *Productus horridus* Sowerby.

*Geologic range:* Upper Carboniferous (Eurasia) and Permian.

*Observations:* In 1928, the year following the publication of Chao's name of this genus, Fredericks introduced the name *Sowerbina* for the same group, but with *Productus timanicus* Stuckenberg as the type. Chao's name clearly has priority.

**HORRIDONIA TEXANA King, n. sp.**

Pl. XXI, Fig. 25

*Description:* In the present collection there is a single specimen of this species, a pedicle valve. It is subtriangular in outline, with the hinge-line straight and

*Rept. 2nd Nor. Arctic Exp. in the "Fram," vol. 4, no. 34, p. 34. 1916.

probably at least as long as the greatest width of the shell. Beak strongly incurved and projecting far beyond the hinge-line, its end being sharply pointed. A narrow, deep median sinus is present on the pedicle valve, starting at the tip of the beak and widening gradually anteriorly; at the anterior margin it is broad and flattened. Surface smooth. Spines not observed, but quite possibly obscured by silicification. Length, 40 mm.; width, 46 mm.; convexity, 21 mm.

*Observations:* This shell appears to belong to the genus *Horridonia*, the sinus and general form being of the characteristic type of *H. horrida*, *H. timanica*, and *H. granulifera*. From them all, however, it differs in having a deeper sinus and a triangular rather than a pentagonal outline.

*Distribution:* Middle Word formation (Glass Mountains) at 255.

### Genus MARGINIFERA Waagen

*Diagnosis:* Shell small, productoid. Visceral cavity deep, brachial valve geniculate. Visceral area of the brachial valve chambered off from the anterior trail by a marginal ridge, which extends from each side of the cardinal process first laterally, then forward, and then around the anterior margin of the visceral area to meet the part of the ridge of the other side. The median anterior part of the ridge may be only imperfectly developed. In the pedicle valve a similar ridge is developed only within the wings. There are three types of surface ornamentation: those with a tuberculose ornamentation, such as *M. cristobalensis*; those with pustules and generally also with wrinkles in the early stages, changing to plicae in the later periods, such as *M. manzanica;* and, the typical form, those with semireticulate ornamentation and their phylogerontic derivatives, such as *M. lasallensis* and *M. typica*.

Genotype: *Marginifera typica* Waagen.

*Distribution:* Viséan to Permian.

*Observations:* The variety in the surface sculpture of these shells indicates that the genus is polyphyletic, but no attempt can here be made to split it along more natural lines. It is likely that the first two types of ornamentation noted above were derived from an *Avonia* ancestor, while the typical form of the genus derived its type of sculpture from a semireticulate *(Productus)* ancestor.

### MARGINIFERA CAPACI (d'Orbigny)

Pl. XXI, Figs. 14–18

1842 *Productus capaci.* D'Orbigny, Voy. dans l'Amér. mérid., t. 3, pt. 4, p. 51; pl. 3, figs. 24–26.

1861 *Productus longispinus.* Salter, Fossils from the High Andes, Quart. Jour. Geol. Soc. of London, vol. 17, p. 64, pl. 3, fig. 2.

1903 *Marginifera haydenensis.* Girty, Carboniferous Formations and Faunas of Colorado, U. S. Geol. Sur. Prof. Paper 16, p. 389, pl. 5, figs. 9–11.

1914 *Productus capaci.* Kozlowski, Brachiopodes du Carb. super. de Bolivie, Ann. de Pal., t. 9, pp. 22–26, text figs. 1–2, pl. 2, figs. 1–15; pl. 5, fig. 12.

*Description:* Shell small, hinge-line forming the greatest width of the shell, sides nearly parallel. Pedicle valve strongly convex and geniculate, the visceral region being relatively short and the anterior slope at maturity about twice as long. Ears fairly large, quadrate, and sharply defined. There is a flattening or slight depression along the median line. Surface marked by faint or obscure irregular costae, and, in the posterior part, by low concentric wrinkles. Spines rather large, numbering 8 to 10 on the anterior slope. Average length, 14 mm.; width, 16 mm.; convexity, 8 mm.

*Observations:* This species is distinguished by its relatively great length, very much enrolled ventral umbo, shallow sinus, evanescent costae, and large spines. In his description of *M. haydenensis*, Girty stated that it could be distinguished from *M. capaci* by its finer costae. The subsequent publication by Kozlowski of figures and descriptions of typical shells from Bolivia has shown that the ornamentation of the two is quite similar, and that they must be synonymous. The Texas specimens are as a whole less deeply sinuate than the ones from Colorado and Bolivia, but they appear to be quite similar in other respects. The actual horizon of the Colorado specimens described by Girty is doubtful. They were collected by the Hayden survey and were not definitely labeled with respect to locality and stratigraphic position. *M. capaci* is very similar to *M. timanica* Tschernyschew, but the latter is even more strongly geniculated and bears no sinus at all.

*Distribution:* Wolfcamp formation, *Uddenites* zone (Glass Mountains) at 88, T88, 203, and 204? Wolfcamp formation (Glass Mountains) at T82, 91, 92, 93, 93s, 196, and Tba. Lower Gym formation (Hueco Mountains) at 376, 381, 383, 384, 387, 390, and 402. Upper Gym formation (Hueco Mountains) at 439.

### MARGINIFERA CRISTOBALENSIS Girty

Pl. XXI, Figs. 19–21

1908 *Productus latidorsatus* var. Girty, The Guadalupian Fauna, U. S. Geol. Sur. Prof. Paper 58, pp. 265–266.

Basal black limestone, Guadalupe Point (sta. 2920).

1909 *Marginifera? cristobalensis.* Girty, Fauna of the Manzano Group, U. S. Geol. Sur. Bull. 389, p. 65, pl. 7, fig. 2.

San Andres formation: Fra Cristobal and San Andres?.

*Description:* Shell very small, outline longitudinally ovoid. Pedicle valve strongly convex, somewhat flattened or even faintly sinuate along the median line. Length of the hinge-line equal to or slightly less than the greatest width of the shell. Ears very small and depressed. Surface near the beak crossed by very fine indistinct concentric wrinkles. Costae not present. Many fairly large spines are distributed over the surface; some of them have elongate bases, but many do not. Brachial valve regularly concave. Marginal ridge low but distinct.

Dimensions in mm. of several specimens:

| | 1 (loc. 120) | 2 (loc. 123) | 3 (loc. 104) | 4 (Girty's type) |
|---|---|---|---|---|
| Length | 14 | 14 | 12 | 11 |
| Width | 13 | 14 | 13 | 12 |

*Observations:* The shells referred to this species have less distinct concentric wrinkling and more prominent spines than Girty's figures of his type specimen show. The original description, however, agrees almost perfectly with the Texas specimens. *M. cristobalensis* is similar in form to *M. manzanica,* but differs markedly in not being costate. It has a great similarity in appearance to *Avonia subhorrida rugatula,* but is narrower and more globose, and has a fainter sinus. The Eurasian species *M. spinosocostata* (Abich) and *M. helica* (Abich) are extremely similar to *M. cristobalensis.*

*Distribution:* Hess formation (Glass Mountains) at 8, 35, 104, and 105. Leonard formation (Glass Mountains) at 3, 7, T9, 12, 13, 16, 21, 23, 26?, 31, 37, 82, 83, T119, 120 (very abundant), 123, 124, 125, 128, T226, 228, 232?, Ax, Tb, and D1. Middle Gym formation (Hueco Mountains) at 424?. Bone Canyon member (Sierra Diablo) at 484.

### MARGINIFERA DUGOUTENSIS King, n. sp.

Pl. XXI, Figs. 6–7

*Description:* Shell small, much wider than long. Pedicle valve strongly convex, geniculated. Ears large and extended, sharply demarked from the visceral region by sinuses. Median sinus narrow but deep, originating nearly at the beak, giving the shell a bilobed appearance; its width varies, being in some specimens very narrow, but on all the shells it is sharply defined and subacute. Surface marked by low, rounded, irregular radial costae, each of which is about 1 mm. wide. Spines very sparsely distributed. The umbonal region and ears are crossed by faint concentric ridges.

Dimensions in mm. of two specimens:

| | 1 (loc. 193) | 2 (loc. 193) |
|---|---|---|
| Length | 10 | 9 |
| Width | 18 | 11 |
| Convexity | 5 | 4 |

*Observations:* This species is most nearly related to *M. wabashensis,* but differs in having a narrower and deeper sinus and longer ears.

*Distribution:* Upper Wolfcamp formation (Glass Mountains) at 193 (types). Wolfcamp formation, *Uddenites* zone (Glass Mountains) at 201?. Upper Wolfcamp formation (Sierra Diablo) at 456?.

### MARGINIFERA LASALLENSIS (Worthen)

1873 *Productus lasallensis.* Worthen, Geol. Sur. Illinois, vol. 5, p. 569, pl. 25, fig. 9.

*Observations:* This species is not abundant at any locality, and since there are no complete specimens the identification may be rather uncertain. It is similar in its ornamentation to the shells from Bolivia identified by Kozlowski as *Productus inflatus,* which have well developed marginal ridges. Some of the specimens figured by Kozlowski are very large, while most are about the same size as *M. lasallensis,* and it is to be wondered if the large specimens may not be distinct, and be true *Productus,* the smaller ones being actually *M. lasallensis.*

*Distribution:* Wolfcamp formation, *Uddenites* zone (Glass Mountains) at 88, 88s, and 95.

### MARGINIFERA MANZANICA Girty

Pl. XXI, Figs. 22–24

1909 *Marginifera? manzanica.* Girty, Fauna of the Manzano Group, U. S. Geol. Sur. Bul. 389, pp. 64–65, pl. 7, fig. 3.
San Andres formation: Fra Cristobal and San Andres?.

*Description:* Shell very small, outline transversely ovoid. Pedicle valve strongly convex, with a mesial flattening or very shallow depression in its anterior half. Hinge-line equal to or a little less than the greatest width of the shell. Ears small and depressed. Posterior part marked only by many fine wrinkles. A few millimeters in front of the beak ribs originate, first as nodes, then becoming persistent and strong and so continuing to the anterior margin; they number about 4 in the space of 5 mm., or 14 or more over the entire surface. The costae give rise to small spines, mostly near the anterior and lateral margins. Brachial valve gently concave in the visceral region and strongly curved around the margin, marked in the visceral region by fine, regular concentric wrinkles, beyond which, toward the margin, there are coarse, not very strong radial costae. At the place at which the wrinkles are succeeded by costae there is a low marginal ridge on the inside of the valve.

Dimensions in mm. of several specimens:

| | 1 (loc. 4) | 2 (loc. 222) | 3 (Girty's type) |
|---|---|---|---|
| Length | 11 | 13 | 12 |
| Width | 13 | 13 | 12 |
| Convexity | 6 | 7 | 6 |

*Observations:* These specimens agree perfectly with Girty's description and figure, except that Girty states that the hinge forms the greatest width of the shell. This species has a great resemblance to *M. ingrata* and *M. carniolicus* Schellwien from the Trogkofel formation, but is distinguished from both by the greater strength of its costae. *M. lopingensis* (Kayser) from the Permian of China has similar ornamentation but is much larger. The most similar American species is *M. whitei,* which is larger and has a greater number of plications.

*Distribution:* Hess formation (Glass Mountains) at 35, 106, and 223. Leonard formation (Glass Mountains) at 4, 9, T9, 15, 20, 21, 24, 26s, 27, 37, 38, 104, T120, T233, Tb, and Tba. Hess formation (Sierra Diablo) at 465. Bone Canyon member (Sierra Diablo) at 484. Victorio Peak member (Sierra Diablo) at 494. Leonard formation (Baylor Mountains) at 502. Leonard formation, transition beds (Shafter) at Th (15979).

### MARGINIFERA OPIMA (Girty)

Pl. XXII, Figs. 20–25

1859 *Productus Popei* (ex parte). Shumard, Trans. Acad. Sci. St. Louis, vol. 1, p. 389.
White (Permian) limestone, Guadalupe Mountains.

1908 *Productus popei* var. *opimus.* Girty, The Guadalupian Fauna, U. S. Geol. Sur. Prof. Paper 58, pp. 258–259, pl. 20, figs. 12–14a.
Dark limestone, Pine Spring (sta. 2930).

*Description:* Shell medium-sized, wider than long. Pedicle valve flat in the visceral region, but geniculated so that the anterior part of the shell lies at a right angle to it. Beak not incurved, and projecting only slightly over the hinge-line. Ears small and depressed. Sinus generally broad and shallow. Costae strong but rounded, numbering from 21 to 28, or about 7 in the space of 10 mm. The costae on the median part of the shell originate at the beak, where they are fine, and increase in size to the geniculation, in front of which their size is constant; the costae on the sides of the visceral region and on the wings originate, both fan-like, by intercalation laterally of the costae arising from the beak, and by bifurcation of the costae in the median part of the shell. On the posterior part of the shell the costae have a somewhat wavy, irregular course. On one large specimen the anterior one-third of the shell is almost smooth, the costae becoming obsolete. Spines few and large, present mainly on the lateral parts of the shell near the ears, where there is a row of three or more of them along the concavity defining the ears.

Brachial valve deeply concave. Cardinal process broad. The brachial ridges arise from a median ridge which extends forward from the base of the process, and is rather high and sharp at its termination at about mid-length of the shell. From each side of the cardinal process there arises a ridge which extends about the margin of the visceral area, at first diverging but slightly from the hinge-line, then swinging forward and finally inward toward the median line, around the front of the visceral area. The anterior part of the marginal ridge is not high and sharp like the sides and the posterior part, being only a slightly elevated highly papillose band.

Dimensions in mm. of several specimens:

| | 1 (loc. 247) | 2 (loc. 247) | 3 (loc. 247) | 4 (loc. 145) | 5 (loc. 145) |
|---|---|---|---|---|---|
| Length | 21 | 25 | 28 | 33 | 28 |
| Width | 29 | 36 | 36 | 32 | 42 |

*Observations:* This shell was originally mentioned by both Girty and Shumard as a variety of *Productus popei,* as the two occur together and have similar configuration and ornamentation. However, it seems to be so easily separable from *M. popei* that it is best to regard it as distinct. It is larger, with more costae, a broad, flat posterior face, strong geniculation, and a peculiar fan-like manner of increase of the costae on the sides of the visceral region. Some of the same characters are shown more or less in *M. popei,* but the posterior face of the latter is generally nearly smooth.

*Distribution:* Word formation (Glass Mountains) at 54, 143, 144, 145 (common), 148, T148, Tx148, 154?, 162, 239, T239, T240, 243, 246, 247 (abundant), 248, and 255. Altuda member (Glass Mountains) at 50. Delaware Mountain formation (Sierra Diablo) at 504.

### MARGINIFERA POPEI (Shumard)

Pl. XXII, Figs. 16–19

1858 *Productus Popei.* Shumard, Trans. Acad. Sci. St. Louis, vol. 1, p. 290 (date of volume, 1860). (Permian, New Mexico and Texas.)

1859 *Productus Popei* (ex parte). Shumard, Trans. Acad. Sci. St. Louis, vol. 1, p. 389, pl. 11, figs. 8–8b. White (Permian) limestone, Guadalupe Mountains.

1908 *Productus popei.* Girty, The Guadalupian Fauna, U. S. Geol. Sur. Prof. Paper 58, pp. 257–258, pl. 20, figs. 9–11b. Dark limestone, Pine Spring (stas. 2930 and 2924). Delaware Mountain formation, southern Delaware Mountains (sta. 2957).

*Description:* Shell small, subquadrate, wider than long, with the greatest width along the hinge. Pedicle valve evenly curved, its profile forming slightly more than a semicircle. Median sinus rather deep and narrow, starting near the beak, giving the shell a somewhat bilobed form. Ears small and rather sharply set off. Brachial valve concave, with a low median fold. Within, the median septum extends about two-thirds the distance to the marginal ridge, which is low but distinct. Surface of the pedicle valve marked by from 6 to 9 coarse, rounded costae on each side of the sinus, some of which bifurcate in the posterior half of the shell; the costae are strongest on the anterior two-thirds of the shell and faint or nearly obsolete on the posterior third. Spines scattered, most of them being on the sides of the visceral area.

Dimensions in mm. of several specimens:

| | 1 (loc. 247) | 2 (loc. 247) | 3 (loc. 246) | 4 (loc. 248) | 5 (loc. 144) |
|---|---|---|---|---|---|
| Length | 20 | 12 | 11 | 20 | 15 |
| Width | 25 | 18 | 13 | 27 | 19 |

*Observations:* This species is distinguishable from the other productids in the fauna by the combination of its small size, bilobate form, strong costae, and by the absence of reticulation on the posterior part. It is somewhat similar to *P. occidentalis,* but is much smaller and has a stronger sinus. It resembles *M. typica* Waagen, but has larger costae than the Indian species and no reticulation near the beak.

*Distribution:* Word formation (Glass Mountains) at 6, 46, 54, 55, 143, 144, T144, 148, T148, Tx148 (abundant), 154, 162, 192, T239, T240a, T240, 242, 246, 247, 248, 261, Ta, and Tc. Very abundant at 246 and 247. Vidrio formation (Glass Mountains) at 43. Leonard formation (Glass Mountains) at 15, 38, and 124. Hess? formation (Glass Mountains) at 98.

### MARGINIFERA POPEI MINOR King, n. var.

Pl. XXII, Figs. 1–2

*Description:* Shell small, its width averaging 16 mm., and its length 13 mm. (probably longer when complete). The pedicle valve is strongly geniculate. Costae low and indistinct, numbering 6 to 8 on each side of the sinus, which is narrow and rather deep.

*Observations:* This variety is distinguished from typical *M. popei* by its generally smaller size, narrower form, narrower sinus, and stronger geniculation. In none of these respects, however, is it very markedly different from *M. popei,* and it does not seem proper to regard it as a distinct species.

*Distribution:* Limestone of Cerro Caballo (Las Delicias) at section 1, bed 14 (west of the Noria), bed 11 (west of the Noria), bed 12, and bed 19.

**MARGINIFERA RETICULATA King, n. sp.**

Pl. XXII, Figs. 3–10

*Description:* Shell small, wider than long, subquadrate. Pedicle valve of low convexity, being depressed in the posterior part and then geniculated. A deep, narrow sinus is present in the anterior part of the shell; it originates back of the geniculation but in front of the beak. Brachial valve nearly flat posterior to the geniculation, with the beginnings of a sinus. Inside, at the line of geniculation there is a raised marginal ridge. Surface of the anterior part of the shell marked by 18 to 20 strong, sharp, irregular costae, separated by rounded sulci. Surface of the posterior part also marked by concentric folds of the same strength as the costae, forming a grid-like surface. There are also very fine concentric lines of growth. Spines are sparsely scattered over the surface, and a row of them lies along the cardinal slope.

Dimensions in mm. of several specimens:

| | 1 (loc. 26s) | 2 (loc. 26s) | 3 (loc. 3) | 4 (loc. 3) | 5 (loc. 232) |
|---|---|---|---|---|---|
| Length | 15 | 14 | 14 | 18 | 14 |
| Width | 22 | 20 | 18 | 24 | 18 |

*Observations:* This species resembles *M. whitei*, with which it is associated, but differs in having a strong sinus, a geniculated rather than a regularly rounded profile, and strong reticulation of the umbonal region. The agreement in other details, however, is so strong as to indicate their near relationship. *M. reticulata* is very closely similar to *Productus gratiosus* Waagen, but differs from it in being strongly geniculated and having coarser costae and a more strongly nodose posterior part.

*Distribution:* Hess? formation (Glass Mountains) at T107 and Tv. Leonard formation (Glass Mountains) at 3, 4, T9, T15, T15a, 16, 24, 26s, 226, 227, 228, 232, T232, T233, Ax, Tb (types), D1, and D4. Leonard formation (Shafter) at Td.

**MARGINIFERA RETICULATA ANGUSTA King, n. var.**

Pl. XXII, Figs. 11–15

*Description:* This variety differs from typical *M. reticulata* in being much narrower in proportion to length and in having only a very shallow sinus. The costae average 15 in number, and are very uneven in size.

Dimensions in mm. of several specimens:

| | 1 (loc. 26s) | 2 (loc. Tb) | 3 (loc. 4) |
|---|---|---|---|
| Length | 13 | 12 | 14 |
| Width | 17 | 16 | 17 |
| Convexity | 6 | 7 | 7 |

*Observations: M. reticulata angusta* is similar in some respects to *M. whitei*, but has much less regular costae, and generally has a shallow sinus. It is much smaller than *Productus mexicanus* of Shumard and of Girty, but otherwise quite similar.

*Distribution:* Upper Hess formation (Glass Mountains) at 207. Leonard formation (Glass Mountains) at 4, 12, 15, T15, T15a, 24, 26s (types), 38, 39, Tb, and Tba. Victorio Peak member (Sierra Diablo) at 494.

**MARGINIFERA SUBLAEVIS King, n. sp.**

Pl. XXXIII, Figs. 13–19

*Description:* Shell small, wider than long, subquadrate. Pedicle valve strongly inflated and geniculated. Ears small, the length of the hinge being equal to or less than the greatest width of the shell. Median sinus rather strong, extending from near the geniculation to the anterior margin. Brachial valve flattened near the beak, then geniculated, the anterior part bearing a median fold. The umbonal part of the shell is quite smooth, but faint radial lirae arise about at the geniculation and cover the anterior slope. The lirae are low and fine, 7 of them occupying a space of about 5 mm. Spines few but large.

Interior of the brachial valve papillose behind the geniculation. Median septum obsolescent. Adductor impressions indistinctly striated. The brachial ridges originate from the median line at the anterior margin of the adductor impressions, and extend forward and laterally nearly to the geniculation, then curve about nearly to meet the first part of the ridge. Strongly developed marginal ridges are present a little in front of the geniculation.

Dimensions in mm. of several specimens:

| | 1 (loc. 120) | 2 (loc. 174) | 3 (loc. 123) | 4 (loc. 125) |
|---|---|---|---|---|
| Length | 17 | 15 | 15 | 18 |
| Width | 23 | 24 | 20 | 21 |
| Convexity | 11 | 9 | 7 | 9 |

*Observations:* The combination of complete absence of reticulation on the posterior part of the shell with the faintness of the costae, the geniculation of the shell, and the presence of a strong median sinus distinguish this species. The most closely similar in appearance is *Avonia latidorsata* of the Delaware Mountain and Capitan. *A. latidorsata*, however, appears to have a fullness in the center of the visceral area so that the pedicle valve is subhemispheric, the faint and narrow sinus dying out both in front and behind. The sinus in *M. sublaevis*, on the other hand, is pronounced and broad, and the sides of the visceral region are inflated into lateral bumps.

*Distribution:* Hess formation (Glass Mountains) at 106. Leonard formation (Glass Mountains) at 4, T9, 12, T41, 82, 83, T119, 120 (types), T120, 123, 124, 125 (very abundant), 174, D9, D12, D13, D22, and Tb. Word formation (Glass Mountains) at T144a and 241?. Hess formation (Sierra Diablo) at 476, 478, and 479. Bone Canyon member (Sierra Diablo) at 485 and 491. Leonard formation (Baylor Mountains) at 503. Leonard formation (Finlay Dome) at 512 and 513. Leonard formation (Shafter) at 514.

**MARGINIFERA? TEXANA (Girty)**

Pl. XXIII, Figs. 5–7

1908 *Productus texanus.* Girty, The Guadalupian Fauna, U. S. Geol. Sur. Prof. Paper 58, pp. 259–260, pl. 21, figs. 25–26b.

Delaware Mountain formation, Guadalupe Point (stas. 2903 and 2919).

*Description:* Shell small. Pedicle valve strongly arched, transversely subquadrate, with a distinct but shallow sinus and abruptly descending sides. Greatest width at the hinge-line. Ears large and convex but poorly defined. Brachial valve regularly and slightly concave. Surface marked by strong costae, numbering 4 to 5 in the space of 5 mm.; they are obscure in the posterior region, which is nearly smooth. Toward the front of the pedicle valve the costae become somewhat coarser as a result of the union of some of the older costae into one. The ears are faintly wrinkled, and bear a few spine bases. Internal characters not known.

Dimensions in mm. of several specimens:

| | 1 (loc. 247) | 2 (loc. 247) | 3 (loc. 247) | 4 (Girty's type) |
|---|---|---|---|---|
| Length | 18 | 18 | 13 | 19 |
| Width | 25 | 30 | 20 | 22 |
| Convexity | 12 | 8 | 6 | 11 |

*Observations:* The only noticeable point of difference between these specimens and Girty's type is the greater length of their ears. They agree in having a very faint sinus and moderately strong costae, which bear very few spines. The costae are finer and the sinus less pronounced than in *M. popei*, with which *M.? texana* is associated. *M.? texana* is much smaller than *M. opima*. As Girty has pointed out, there is great similarity to *M. capaci* (d'Orbigny), but the costae of that species are much finer and the surface is more spinose.

*Distribution:* Word formation (Glass Mountains) at T148, 246, and 247 (types).

**MARGINIFERA WABASHENSIS (Norwood and Pratten)**

Pl. XXIII, Fig. 4

1854 *Productus wabashensis.* Norwood and Pratten, Acad. Nat. Sci. Philadelphia, Jour., (2), vol. 3, p. 13, pl. 1, figs. 6a–d. (Date of whole volume, 1855.)

1903 *Marginifera wabashensis.* Girty, Carboniferous Formations and Faunas of Colorado, U. S. Geol. Sur. Prof. Paper 16, pp. 375–379, pl. 5, figs. 8–8a (see for synonymy).

*Description* (from Girty): Shell broad, highly arched, "with strong median sinus and slightly extended ears. The beak is small and transgresses the hinge-line but little. Concentric wrinkles are almost wanting; the striae are fine and obsolescent, and the surface supports a few comparatively large scattering spines."

*Distribution:* Wolfcamp formation, *Uddenites* zone (Glass Mountains) at 88s (abundant), 94, 95, and 201. Wolfcamp formation (Glass Mountains) at 75, 89, and 195. Leonard? formation (Glass Mountains) at T9 and 20.

**MARGINIFERA? WHITEI King, n. sp.**

Pl. XXIII, Figs. 1–3

1877 *Productus mexicanus?* White (non Shumard), U. S. Geog. Sur. W. 100 Mer., Rept., vol. 4, p. 120, pl. 8, figs. 6a–c.

Carboniferous: Camp Cottonwood, old Mormon road, Lincoln County, Nevada; near Salt Lake, New Mexico.

1883 *Productus mexicanus.* Kayser, Richthofen's China, vol. 4, p. 182, pl. 28, figs. 7a–b.

Permian: Lo-Ping, China.

1902 *Productus mexicanus.* Tschernyschew, Com. Geol. (St. Petersburg), Mem., vol. 16, no. 2, p. 264, pl. 52, fig. 10.

*Schwagerina* zone of the Urals.

1909 *Productus mexicanus?* Girty, Fauna of the Manzano Group, U. S. Geol. Sur. Bull. 389, pp. 57–58, pl. 7, fig. 4 (?).

Yeso formation: Mesa del Yeso; San Andres formation: Nogal Creek and San Andres.

*Description:* Shell small, subquadrate, hinge-line equal to or greater than the greatest width of the shell. Pedicle valve subhemispherical, rather strongly convex and regularly arched, but not inflated near the beak. Some of the shells show a faint sinus, but others are nearly evenly hemispherical. Ears small but well-defined. Surface marked by 14 to 18 coarse, rounded, unequal costae, some of which are twice as large as the others. Posterior part of the shell crossed by concentric folds, which give it a nodose aspect. The entire surface is covered with fine growth lines. Spines few and large, arising from the costae. Brachial valve flattened in the visceral region, then geniculated.

Dimensions in mm. of several specimens:

| | 1 (loc. 107) | 2 (Girty—San Andres) | 3 (White) |
|---|---|---|---|
| Length | 12 | 22 | 14 |
| Width | 14 | 22 | 15 |
| Convexity | 6 | 13 | — |

*Observations:* This species is distinguishable from others in this fauna by its strong costae, general absence of a sinus, low beak, small ears, and nodose posterior part. It agrees perfectly with the specimens from Nevada and New Mexico identified by White as *Productus mexicanus,* yet it is certainly different from *P. mexicanus* Shumard, which is a large shell collected originally from the Capitan. The identity of *M. whitei* with *P. mexicanus* Girty from the Yeso and San Andres is doubtful, though they occur in a similar faunal association. The shells from New Mexico agree better with Shumard's species than do the Leonard specimens. No important difference can be seen between *M. whitei* and the specimens from China and Russia identified as *P. mexicanus* White. *M. manzanica* is also very similar, but is smaller, narrower, and has fewer plications.

*Distribution:* Hess formation (Glass Mountains) at 17a, 107, T107, 207, 208, and 222. Hess formation (Sierra Diablo) at 476, 479, and 485. Leonard formation (Glass Mountains) at T226. Leonard formation (Findlay Mountains) at 512. Upper Gym formation (Hueco Mountains) at 440?.

**MARGINIFERA? WORDENSIS King, n. sp.**

Pl. XXIII, Figs. 8–11

*Description:* Shell very small, slightly wider than long. Pedicle valve very convex, geniculated at mid-length, the posterior slope being gently convex and the anterior flattened. Ears broad and flat, so extended that the greatest

width of the shell is at the hinge-line. Beak pointed and not incurved. Median part depressed into a faint sinus, or in some specimens convex. Surface marked by about 10 strong costae, each about 1.5 mm. wide, and each bearing several small spine bases. Brachial valve slightly concave. A distinct marginal ridge is not present, there being only a papillose band about the anterior part of the visceral area.

Dimensions in mm. of several specimens:

| | 1 (loc. 162) | 2 (loc. 239) | 3 (loc. T240) |
|---|---|---|---|
| Length | 11 | 10+ | 11 |
| Width | 13 | 15 | 14 |
| Convexity | 7 | 4+ | 5 |

*Observations:* This species is distinguished by its small size, geniculated profile, and strong costae. It is similar to *M. manzanica* but has larger costae. It has stronger costae and stronger geniculation than *M.? whitei*, and is shorter in proportion to width.

*Distribution:* Word formation (Glass Mountains) at Tx148, 162, 239, T239, T240 (types), 265, and Tx.

### Genus SCACCHINELLA Gemmellaro

*Diagnosis:* Shell irregular in growth, pedicle valve conical, brachial valve nearly flat. Hinge-line shorter than the greatest width of the shell. Ventral cardinal area very high, longitudinally striated, with an indistinct deltidium. Dorsal area very low or absent. Shell surface marked with many small tubular spines on the pedicle valve and similarly arranged pustules on the brachial valve. Inner shell layer pustulose. In the specimens having very high, conical pedicle valves, the inside of the valve is filled by irregularly arranged transverse partitions. Within the pedicle valve there is a high median septum and near the apex faint flabelliform muscular impressions lie on each side of the septum. In the brachial valve the cardinal process consists of two distinct long apophyses, each base of which continues forward as a ridge, straddling the septum of the pedicle valve.

Genotype: *Scacchinella variabilis* Gemmellaro.

*Geologic range:* Permian.

*Observations:* According to Gemmellaro, *Scacchinella* is in youth attached at the apex, later by one of the sides or by the anterior part of the pedicle valve. The evidence for this conclusion is not stated.

There is some disagreement as to the taxonomic relations of this genus. It is placed by Schuchert* in the Orthotetinae, and by Brolli* in the Richthofenidae. It is more probably related to neither. Its spinose surface, indistinct deltidium, deeply bilobed cardinal process unsupported by crural plates show that it cannot be placed with the Orthotetinae. According to Likharew* the median septum is entirely differently constructed from that in the strophomenids. The broad cardinal area, absence of an enveloping layer, and the development of the median septum from the infolding of the inner side of the cardinal area rather than independent of the area as in *Richthofenia* prove that it is quite different from the Richthofenidae.

*In Zittel's Textbook of Palaeontology, second edition. 1913.

*In Zittel's Gründzuge der Paläontologie, fifth edition. 1921.

*Palaeontolog. Zeitsch., bd. 10 (1928), pp. 287–288.

The most clearly related genus to *Scacchinella* is *Tschernyschewia*, the important features of which are the spinose surface, low cardinal area, high median septum, and cardinal process consisting of two elongate apophyses, from the bases of which thickened ridges extend anterolaterally. All these features are like those of *Scacchinella*, except that the pedicle valve of the latter is much deeper, and the cardinal area much higher. A low dorsal median septum is present in *Tschernyschewia*, which has not been noted in *Scacchinella*.* The only argument against regarding *Tschernyschewia* as the ancestor of *Scacchinella* is the fact that the only known occurrences of the former are in the middle Permian (Lo-Ping) of China and the upper Permian near Djulfa, Armenia, while *Scacchinella* first occurs in the very lowest Permian. Otherwise the agreement is so nearly perfect that one cannot help but suspect that with more complete knowledge of Permian faunas *Tschernyschewia* may be found at a lower horizon. *Tschernyschewia* appears to have arisen from *Buxtonia*. *B. porrecta (=B. peruviana)* has a low ventral median septum, and for this reason was made the type of a new subgenus, *Tschernyschewiella*, by Fredericks (although the name proves to have been preoccupied). It appears likely that *Scacchinella* is a member of the subfamily Productinae, which arose from *Buxtonia* through *B. porrecta* and *Tschernyschewia*.

### SCACCHINELLA GIGANTEA Schellwien

Pl. XXIII, Figs. 20–25; Pl. XXIV, Figs. 1–5; Pl. XXV, Figs. 16–18

1898 *Scacchinella variabilis*. Schellwien, Sitzungsber, d. Akad. d. Wiss. Berlin, p. 697, no. 61; and Verhandl. d. k. k. geol. R.-A. Wien, no. 16, p. 360.

1900 *Scacchinella gigiantea*. Schellwien, Fauna der Trogkofelschichten, Abh. der k. k. géol. Reichs-Anstalt, bd. 16, heft 1, pp. 35–37, pl. 4, figs. 1–3; pl. 5, figs. 1–8.

*Description:* Shell large, the pedicle valve being a high cone, with the posterior side flattened by the large cardinal area; the brachial valve operculiform. Apex of the pedicle valve bluntly pointed, the valve expanding gradually, with irregular concentric undulations. Hinge-line somewhat shorter than the greatest width of the shell. Cardinal area nearly flat, but in some specimens from loc. 35 bent backward about 20° at a transverse angulation near the middle of the area. The area is generally twisted to one side. Deltidium very narrow, linear, and poorly defined. Surface of the cardinal area vertically striated, and marked by transverse growth striae. Brachial valve very slightly convex, with an inconspicuous beak. Surface of both valves uneven; the outer layer of shell is not preserved in the specimens of the present collection, as they were broken from limestone, but it appears to have been like that of *Buxtonia*.

In the pedicle valve the median septum is very long and high, extending from the beak to the hinge-line, where its

*The relations of *Tschernyschewia*, *Scacchinella*, and *Richthofenia* have been discussed in detail by Stoyanow: On Some Permian Brachiopoda of Armenia, Mém. du Com. Géol. (Russie), n. s., liv. 111, pp. 55–74. Likharew (Palaeontolog. Zeitsch., bd. 10, pp. 274–288) has justly criticized Stoyanow's comparison of *Scacchinella* with *Richthofenia*.

dorsal edge lies parallel to the inside of the brachial valve and about 2 mm. from it. Its height is variable, equaling as little as a fourth and as much as three-fourths the diameter of the shell, but generally equaling about half the diameter. The septum consists of two parallel lamellae, which apparently represent greatly developed and closely appressed dental lamellae. The shell tends to cleave between these lamellae. Most of the inside of the pedicle valve is filled with transverse partitions or tabulae. These plates do not extend individually across the entire diameter of the shell, but join each other at low angles to form a very coarse vesicular structure. The shell breaks so easily along these partitions that entire specimens can rarely be removed from the rock. The cardinal process was deeply bilobed, consisting of two thin and very elongate lamellae which straddle the great septum of the pedicle valve.

Dimensions in mm. of several specimens:

| | 1 (loc. 208) | 2 (loc. 208) | 3 (loc. 208) | 4 (loc. 208) | 5 (Sch.) |
|---|---|---|---|---|---|
| Height of cardinal area | 30+ | 70+ | 38 | 12 | 59 |
| Length of hinge-line | 21 | 34 | 30 | 12 | 46 |
| Width of brachial valve | 34 | — | — | 17 | 59 |
| Length of brachial valve | 33 | — | 37 | 13 | 55 |

The specimens from the Wolfcamp formation are very small (the largest one being 17 mm. wide) and it cannot therefore be told if they would be of large size and contain partitioning plates when mature. They agree perfectly, however, with small specimens of *S. gigantea* from the Hess. The specimen from loc. 71 has a width of 55 mm., but the complete length of the pedicle valve cannot be determined because only the anterior part is preserved. Traces of transverse partitions are present. It must have been a very large shell.

*Observations:* All the features shown by the Glass Mountains specimens strongly indicate their complete identity with those from the Karnic Alps. The internal characters of the brachial valve were not, however, fully determined in the specimens of the present collection. *S. gigantea* is distinguished from *S. variabilis* Gemm. by its much larger size, higher ventral median septum, the presence of partitions in the pedicle valve, and by the internal differences of the brachial valve.

*Distribution:* Wolfcamp formation, *Uddenites* zone (Glass Mountains) at 199. Wolfcamp formation (Glass Mountains) at 71. Hess formation (Glass Mountains) at 4?, 15, 17a, 35 (abundant), 106, 108, T108, 208 (abundant), 209, and 211.

### Genus AULOSTEGES Helmersen

*Diagnosis:* Shells small to very large, pedicle valve convex, brachial valve flat or concave. Surface wrinkled and very spinose. Apex of the pedicle valve more or less deformed; ventral cardinal area large and nearly flat, bearing in the middle a narrow deltidium, provided, if well preserved, with small spines. Hinge-teeth rudimentary if present at all. Dorsal cardinal area linear, cardinal process large. Brachial ridges generally not well developed.

Genotype: *Aulosteges variabilis* Helmersen (=*Orthis wangenheimi* de Verneuil).

*Geologic range:* Permian.

*Observations:* There is a great deal of doubt as to the correctness of the name *Aulosteges* for the group of shells here assigned to it. This matter has been discussed at length by Girty and Chao.* The genotype of *Strophalosia, S. excavata,* may be a member of the group commonly designated as *Aulosteges.* If so, the name *Aulosteges* is a synonym, and a new name must be found for the shells commonly known as *Strophalosia.* The distinguishing characters of the two genera as recognized by Waagen are that in *Strophalosia* the umbo is deformed by attachment, the muscle scars of the brachial valve are raised and not dendritic, and the cardinal teeth are constantly present, while in *Aulosteges* the shell is free from umbonal attachment, the muscle scars are dendritic, and the cardinal teeth are either absent or extremely rudimentary. These cannot be used to distinguish the species in the present collection, however, for in *A. beedei* the muscle scars are raised and not dendritic, though the other characters are in agreement. More reliable features defining the two groups are the following: *Strophalosia* is small, with low, imperfectly developed cardinal area; it was either cemented to some larger organism or marked by a distinct scar of attachment. *Aulosteges* is large, generally with a high, well-developed cardinal area, and without evidence of attachment. There is a great deal of difference in the internal characters of the brachial valve in the various species assigned to *Aulosteges,* and it is possible that several genera may be distinguished in it on that basis.

### AULOSTEGES BEEDEI King, n. sp.

Pl. XXV, Figs. 12–15

*Description:* Shell small to medium-sized, of low curvature and subquadrate outline. Pedicle valve of low convexity, upturned near the margin. Ventral beak very small and low; umbonal region only obscurely separated from the ears. Hinge-line about equal to the greatest width of the shell. Cardinal area high, very uneven in curvature, marked by strong longitudinal undulations and lirae, and by fine vertical lines. The area is divided by the delthyrium, which is covered in its posterior half by narrow, triangular, strongly convex deltidium. The half of the delthyrium near the hinge-line is occupied by the large cardinal process.

Brachial valve slightly concave or quite flat, with no cardinal area. Its posterior part appears to overgrow the edge of the cardinal area of the other valve. Cardinal process long and narrow, the posterior face lying at an acute angle to the plane of the shell margin, slanting toward the front. The posterior face has the form of a rounded ridge; the anterior face is deeply excavated by a groove. From each side of the base of the process extends a ridge, which expands forward and bears the adductor muscle scars; these are raised on elongate platforms, with a deep groove between. The internal characters are described from specimens from loc. Tx.

Surface of the pedicle valve marked by very strong concentric wrinkles, which extend from ear to ear, or, in the umbonal region, from one side of the beak to the other against the hinge-line. On the lateral slopes the growth

*Girty: U. S. Geol. Sur. Prof. Paper 58, pp. 274–275. 1908.
Chao: Pal. Sinica, ser. B, vol. 5, fasc. 3, pp. 69–71. 1928.

wrinkles pass into strong laminae, which converge toward the ears. Spaced at intervals of about 2 mm. are large spine-bases, from which there extend spines, lying against the shell surface and directed radially from the beak; they are 2 to 3 mm. in length. Radial lirae are absent. Brachial valve faintly wrinkled, but without spines.

Dimensions in mm. of several specimens from loc. 123:

| | (1) | (2) | (3-brachial valve only) |
|---|---|---|---|
| Length | 20 | 28 | 15 |
| Width | 19 | 23 | 18 |

*Observations:* This species is distinguished by its low, uneven convexity, elongate quadrate outline, strong concentric wrinkles, the absence of radial lirae, the abundance of prostrate, forward-directed spines, and the high, irregular cardinal area.

*Distribution:* Leonard formation (Glass Mountains) at 123 (types). Word formation (Glass Mountains) at 53 and Tx.

### AULOSTEGES GUADALUPENSIS Shumard

Pl. XXV, Figs. 8–11

1858 *Aulosteges guadalupensis.* Shumard, Trans. Acad. Sci. St. Louis, vol. 1, p. 292 (date of volume, 1860).
White (Permian) limestone: Guadalupe Mountains, New Mexico and Texas.

1859 *Strophalosia (Aulosteges) Guadalupensis.* Shumard, *idem*, p. 390, pl. 11. figs. 5a–b.
White (Permian) limestone: Guadalupe Mountains.

1890 *Strophalosia? Guadalupensis.* Beecher, Am. Jour. Sci., ser. 3, vol. 40, p. 241.

1908 *Aulosteges guadalupensis.* Girty, The Guadalupian Fauna, U. S. Geol. Prof. Paper 58, pp. 277–278, pl. 20, figs. 22–22a.
Upper dark limestone, Pine Spring (sta. 2930).

*Description:* Shell medium-sized to large, outline subtriangular. Pedicle valve moderately and regularly convex. Umbonal region flattened; beak erect and sharply pointed. Hinge-line shorter than or equaling the greatest width of the shell. Ears small and rounded. Median sinus broad, shallow, but distinct, originating some distance in front of the beak and extending to the anterior margin. Cardinal area triangular and high, sharply demarked from the rest of the shell. Deltidium not distinguishable, but the delthyrium appears as a shallow notch in the mid-part of the area. Brachial valve flattened in the visceral region, then slightly geniculated in the marginal region. Surface of the pedicle valve covered with small tapering spines that incline strongly forward, separated by pits. On some shells the spine bases are arranged more or less quincuncially, while on others they show no definite arrangement, or seem to be borne on irregular longitudinal costae. Brachial valve pitted by inward-projecting spine bases.

Dimensions in mm. of several specimens:

| | 1 (loc. 144) | 2 (loc. 58 | 3 (loc. 248) |
|---|---|---|---|
| Length | 23 | 18 | 35 |
| Width | 22 | 37 | 58 |

*Observations:* This species is easily distinguished from others in the fauna by its small, longitudinally elongate spine bases, which are thickly set over the entire surface.

*Distribution:* Lower Word formation (Glass Mountains) at 58 and 241. Middle and Upper Word formation (Glass Mountains) at 144 (abundant), T144a, 146, 148, T148, Tx148, 239?, T239, 240, T240, 243, 247, 248, 252, 265, and Tx. Word formation (Glass Mountains) at 167?. Delaware Mountain formation (Sierra Diablo) at 504.

### AULOSTEGES MAGNICOSTATUS Girty

Pl. XXV, Figs. 1–4

1908 *Aulosteges magnicostatus.* Girty, The Guadalupian Fauna, U. S. Geol. Sur. Prof. Paper 58, pp. 278–279, pl. 31, figs. 4–4b.
Delaware Mountain formation, Glass Mountains

*Description:* Shell small, wider than long, semicircular. Pedicle valve with an extended beak and a longitudinally flattened posterior part to about two-thirds the length from the beak to the front margin, where it is strongly geniculated. Cardinal area triangular, symmetrical or twisted, divided by a narrow but strongly convex deltidium. Brachial valve flattened in the visceral part and strongly curved up at the front and sides, its beak small.

Surface of both valves marked by very fine, irregular, sharp concentric wrinkles, which bear many spine bases. Toward the front and sides the wrinkles die out and are replaced by relatively coarse radial costae, which are faint at first but become strong. At the posterior end of each costa are one or more spine bases. The anterior margin of the pedicle valve is bordered by a raised band, which is smooth except for concentric striae. The costae extend under this band; just posterior to the band is a row of large spines, one on each costa. The brachial valve bears no band. Fine but sharp growth lines cover the entire surface.

Dimensions in mm. of several specimens:

| | 1 (loc. 123) | 2 (loc. 228) | 3 (loc. 4) | 4 (Girty's type) |
|---|---|---|---|---|
| Length | 11 | 10 | 12 | 9 |
| Width | 14 | 12 | 14 | 8 |
| Ht. card. area | 1.5 | — | — | 2 |

*Observations:* Some of the specimens in the present collection, particularly those from locs. 1, 104, and 128, are relatively broader than Girty's type. The presence of the marginal smooth band and the large costae of the geniculated part of the shell are the most striking characters of this species.

*Distribution:* Hess formation (Glass Mountains) at 1, 35, and 207. Leonard formation (Glass Mountains) at 4, 8?, T9, 26s, 31, 38, 25–29, T41, 104, T119, 120, 123, 128, 228, D5, and Tb. Word formation (Glass Mountains) at 237 and 245. Bone Canyon member (Sierra Diablo) at 484.

### AULOSTEGES MEDLICOTTIANUS Waagen

Pl. XXVI, Figs. 7–11; Pl. XXVII, Figs. 8–9

1884 *Aulosteges medlicottianus.* Waagen, Salt Range Fossils; Pal. Ind., ser. 13, vol. 1, Productus Limestone Fossils, Pt. 4, pp. 663–666, pl. 62.
Amb group (lower Productus limestone): Salt Range.

?1914 *Aulosteges* aff. *medlicotti.* Tschernyschew, Fauna der Oberpal. Ablager. des Darwas, Mem. Com. Geol. (Russie), n. s., liv. 104, pp. 4, 66, pl. 10, fig. 4.
Artinsk: Darwas, Bokhara.

*Description:* Shell medium-sized, subquadrate (but generally appearing to be subtriangular because the ears are broken off), width equal to or slightly less than the length; sides subparallel. Pedicle valve of low curvature over the visceral region to about mid-length of the shell, then strongly curved up to the front margin. In transverse direction the surface is moderately convex, but with more or less of a median sinus. Ears flattened and sharply set off. Beak bluntly pointed and not prominent, extending slightly over the gently concave cardinal area. The small triangular delthyrium was probably covered by a deltidium, although that structure is not preserved. The anterior margin of the shell may turn back to form a narrow flange, as wide as 5 mm. Brachial valve concave, with a faint median fold. Cardinal process large, similar to that of *A. beedei*, but, because this valve is more concave it projects farther back.

Surface of the pedicle valve marked by wavy lirae of uneven size, which number about 15 in the space of 5 mm., they are generally obscured by silicification. The entire surface is crossed by strong concentric wrinkles and fine concentric growth lines. Large hollow spines are distributed over the surface at intervals of a few millimeters. Surface of the brachial valve lirate and wrinkled, marked by dimples rather than spines. Shell fibrous and strongly punctate.

Dimensions in mm. of several specimens:

| | 1 (loc. 38) | 2 (loc. 222) | 3 (loc. 108) | 4 (loc. 124) | 5 (loc. 124) |
|---|---|---|---|---|---|
| Length | 34 | 30 | 30 | 37 | 36 |
| Width | 32 | 35 | 38 | 43 | 37 |

*Observations:* This species is distinguished from related forms in this fauna by its flattened posterior region, strong concentric wrinkles over the entire surface, fine, uneven lirae, and its large, rather abundant spine bases. It is somewhat smaller than typical *A. medlicottianus*. Waagen states that his species has fine radial wrinkles of the shell substance rather than external radial markings, and it may be that the lirae above described are of this nature. *A. medlicottianus* differs from *A. hispidus* Girty, a Phosphoria species, in being larger, having a subquadrate rather than a subtriangular form, and in having stronger concentric wrinkles and larger spines. *A. poyangensis* is somewhat similar, but has a very prominent ventral beak.

The specimen described by Girty under the name *Aulosteges medlicottianus* var. *americanus* appears to have no relation to the form now under discussion. Girty suggested that it might be a *Strophalosia* or a *Streptorhynchus*, but to me it seems even more to resemble *Rhipidomella*, the points of resemblance being the evenly lirated shell, oval outline, broad delthyrium, and the tubules piercing the lirae. The specimen described by Girty as *Aulosteges* sp. b is very probably *A. medlicottianus.*

*Distribution:* Upper Wolfcamp formation (Glass Mountains) at 70. Hess formation (Glass Mountains) at 1, 4, 107, T107, 108 (abundant), 222, and 91x?. Leonard formation (Glass Mountains) at T9, 15, 23?, 38 (abundant), T39, 120?, 121, 124, 226, T226, T233, Tb (abundant), Tba, Ax, and D3 (abundant). Word formation (Glass Mountains) at 44, 46, and 144.

### AULOSTEGES SUBCOSTATUS King, n. sp.

Pl. XXV, Figs. 5–7

*Description:* Shell small, slightly wider than long, subtriangular in outline except for the ears. Pedicle valve strongly and evenly convex, with the greatest convexity near the beak, which is pointed and extends somewhat over the hinge-line. Visceral region triangular, with steeply descending sides, depressed into a broad, shallow, undefined sinus of variable prominence. Hinge-line straight, shorter than the greatest width of the shell, bearing a low cardinal area. Brachial valve slightly concave, with a well-defined cardinal area, which is slightly smaller than that of the other valve.

Shell very thick. In the pedicle valve the diductor muscles diverge radially from the beak in the form of deeply impressed grooves, separated by a broad platform bearing the unevenly striated adductor scars. Median septum absent.

Surface of the pedicle valve marked by fine concentric growth lines and by sparsely distributed spine bases, which are elongated so as to appear as low, interrupted costae. Just in front of the cardinal margin is a row of large laterally directed spines. Surface of the brachial valve nearly smooth. Shell fibrous and finely punctate.

Dimensions in mm. of several specimens, all from loc. 123:

| | | | |
|---|---|---|---|
| Length | 18 | 17 | 20? |
| Width | 20 | 18 | 20 |
| Convexity | 9 | 9 | 12 |

*Observations:* This species is distinguished by the triangular form of its visceral region, its extended ears, its strong convexity, the presence of both valves of a low cardinal area, the absence of strong concentric wrinkles, and the few spines, whose bases are greatly elongate. It is similar in general appearance to *Strophalosia costata* Waagen, but there is no evidence of a scar of attachment as in that species.

*Distribution:* Leonard formation (Glass Mountains) at T119, 120, and 123 (types).

### AULOSTEGES TRIAGONALIS King, n. sp.

Pl. XXVII, Figs. 1–3

*Description:* Shell small, generally longer than wide, outline triangular. Pedicle valve flattened longitudinally to about two-thirds its length from beak to front, where it is strongly geniculated. Median region depressed into a deep, wide, rounded sinus, which starts a short distance behind the geniculation. Beak extended and sharply pointed, with no ears, the hinge-line extending no farther than the edge of the visceral region. Cardinal area small and triangular, most of its area being taken up by the delthyrium, which is occupied by the cardinal process of the brachial valve. Brachial valve evenly concave.

Surface of both valves ornamented by strong, irregular concentric wrinkles bearing many spines on the pedicle valve. Radial ornamentation not present. Shell fibrous.

Dimensions in mm. of several specimens:

| | 1 (loc. 8) | 2 (loc. 37) | 3 (loc. 231) | 4 (loc. 237) |
|---|---|---|---|---|
| Length | 13 | 13 | 12 | 12 |
| Width | 14 | 12 | 13 | 12 |

*Observations:* This species is easily distinguishable by its small size, very short hinge, narrow cardinal area, median sinus, and absence of radial ornamentation.

*Distribution:* Hess formation (Glass Mountains) at 8. Leonard formation (Glass Mountains) at 9, 37, 83, 231 (type), 232, 233, D4, and D7. Word formation (Glass Mountains) at 237, 239, D239, and Tx. Leonard formation (Baylor Mountains) at 503. Bone Canyon member (Guadalupe Mountains) at 221?.

### AULOSTEGES TUBERCULATUS King, n. sp.

Pl. XXVII, Figs. 4–7

*Description:* Shell subtriangular, width and length about equal. Pedicle valve low, geniculated near the front margin. Beak narrow, expanding gradually into the triangular visceral area, which is widest at the front, the umbonal angle being about 70°. The ears are nearly flat but large, so that the hinge-line about equals the greatest width of the shell. The entire pedicle valve is abundantly spinose, the ears bearing the largest spines, which are spaced 2 to 5 mm. apart and directed backward. The visceral part of the shell is marked by approximately quincuncially arranged large conical tubercles, which are the bases of spines. Fine concentric wrinkles, which are especially strong on the ears, cross the entire surface. On the marginal part of the shell the spines give rise to large subangular costae, each about 3 mm. wide. The beak of the valve may be nearly straight or considerably arched over the cardinal area, which is high and striated parallel to the hinge margin. Delthyrium narrow, the upper part covered by a small deltidium, and the lower a broad open notch.

Brachial valve nearly flat over most of its area, but with an undulating surface. Marginal part geniculated. Surface of the posterior part marked by dimples of about the same size and appearance as the tubercles of the other valve, the only difference being that they point inward rather than outward. The geniculated part is marked by costae like those of the other valve.

Dimensions in mm. of several specimens:

| | 1 (loc. 247) | 2 (loc. 247 brachial) | 3 (loc. 146) | 4 (loc. T239) |
|---|---|---|---|---|
| Length | 55 | 35 | 31 | 50 |
| Width | 60 | 40 | 34 | 42 |

*Observations:* This species is distinguished from other members of the genus by its coarse tuberculate-costate ornamentation and sprawling growth.

*Distribution:* Word formation (Glass Mountains) at 144 (abundant), 146, Tx148, 152, 239, T239 (types), T240, 241?, 243, and 247. Delaware Mountain formation (Sierra Diablo) at 504.

### AULOSTEGES WOLFCAMPENSIS King, n. sp.

Pl. XXVI, Figs. 1–6

*Description:* Shell rather large, subtriangular, expanding from a bluntly pointed posterior to a broadly rounded anterior end, the outline being interrupted by the projection of small ears. Hinge-line slightly less than the greatest width of the shell. Pedicle valve rather strongly convex, but with flattened profile both in front of and behind a region of strong curvature which lies at about mid-length or in front of mid-length. Transversely the profile is flattened over the visceral cavity and descends steeply to the lateral margins. Median part depressed into a shallow, narrow, but distinct median sinus. Ears small and flattened. Cardinal area short and low, lying in about the plane of the shell margin.

Brachial valve regularly concave, but abruptly geniculate near the margin. The interior of a brachial valve from loc. 93, probably belonging to this species, has a large cardinal process, which is narrow on its posterior side near the hinge-line, and expands toward the anterior side, where it is divided at the end into three lobes. From the base of the process extend two strong lateral ridges, which lie parallel to the hinge margin, and a low median septum, which is bordered on each side by raised lobes bearing the muscular impressions. Brachial ridges indistinct.

Surface of both valves marked by strong rounded radial lirae, numbering about 9 in the space of 5 mm., crossed by very fine, regularly arranged concentric lirae. At intervals of 4 to 5 mm. on the pedicle valve there are large forward-directed spines, arranged more or less quincuncially, practically restricted to the posterior half of the valve; on the brachial valve there are corresponding depressions. Posterior part of the pedicle valve crossed by a few faint concentric wrinkles, but the entire brachial valve strongly wrinkled.

Dimensions in mm. of several specimens:

| | 1 (loc. 93) | 2 (loc. 93s) | 3 (loc. 143) | 4 (loc. 196) |
|---|---|---|---|---|
| Length | 35? | 30 | 33 | 33 |
| Width | 40 | 36 | 40? | 38 |
| Convexity | 14 | 15 | 16 | 15 |

*Observations:* This species is distinguished from others in this fauna by the triangular outline of the visceral cavity, faint wrinkling of the pedicle valve, strong radial lirae, and regularly arranged, sparsely distributed spines. The internal characters indicate a relation to the productids, as the cardinal process is more productid than that in such a specialized form as *A. beedei.* The appearance of the muscle scars in the brachial valve is, however, similar to that of *A. beedei,* though they are not as much elevated above the general level of the valve. It may be that this species forms a connecting link between the productids and *Aulosteges.* If so, it probably arose from a *Linoproductus,* judging from the surface ornamentation. *Aulosteges* is generally regarded as having arisen from the earlier *Strophalosia* stock, and is put in the same subfamily, but it is very likely that it arose independently from the productids at the beginning of the Permian. Unfortunately, no conclusions confirmatory or otherwise of

this can be drawn from Tschernyschew's figures and descriptions of *Aulosteges dalhousii*, the first Eurasian species.*

*Aulosteges medlicottianus* is much more strongly wrinkled and spinose than *A. wolfcampensis*, and is not strongly lirate; its general form is, however, similar. *A. hispidus* Girty has a somewhat similar configuration, but its radial lirae are much fainter and the spines are most prominent near the anterior margin, rather than in the posterior half of the pedicle valve as in *A. wolfcampensis*.

*Distribution:* Wolfcamp formation, *Uddenites* zone (Glass Mountains) at 88, 88s, T88, 94, 201, and 203. Wolfcamp formation (Glass Mountains) at 71, 85, 86, 87 low?, 90, 92, 93 (types), 93s, 193, 196, 198, and Tba.

#### Genus STROPHALOSIA W. King

*Diagnosis:* Shells productoid, with a distinct scar of attachment at the ventral umbo, or attached by nearly the entire outer surface of the pedicle valve. Pedicle valve with a well defined cardinal area, the delthyrium closed by a deltidium; hinge-teeth prominent but not supported by lamellae. Brachial valve with a narrow cardinal area; cardinal process erect, bifid on its outer and trifid on its posterior face, supported laterally by socket plates and continued anteriorly into a median septum which extends half the length of the valve. Surface of the pedicle valve spinose, the spines near the beak commonly recurved and embracing some external object, and in some species all the spines assisting in the attachment of the shell. Brachial valve spinose, lamellose, or smooth externally.

Genotype: *Orthis excavata* Geinitz.

*Geologic range:* Middle Devonian to Permian.

*Observations:* As pointed out in the discussion of *Aulosteges*, the group of shells to which the name *Strophalosia* is generally applied may not be congeneric with the genotype of *Strophalosia*, in which case a new name must be found for the group.

#### STROPHALOSIA HYSTRICULA Girty

Pl. VII, Fig. 9; Pl. XXVII, Figs. 10–13

1908 *Strophalosia hystricula.* Girty, The Guadalupian Fauna, U. S. Geol. Sur. Prof. Paper 58, p. 275, pl. 30, figs. 14–14a.

Delaware Mountain formation, "Comanche Canyon," Glass Mountains.

*Description:* Shell small, subcircular. Pedicle valve moderately convex, beak somewhat elevated. Hinge-line shorter than the greatest width of the shell. Ears small and undefined. Cardinal area linear. Surface covered by long, tubular, upwardly convex spines, directed forward so as to be tangent to the surface. On the ears there are very large laterally directed spines, each about 6 mm. long. Brachial valve slightly concave. Along the hinge-line are curved spines, directed toward the beak.

Dimensions in mm. of several specimens:

| | 1 (loc. 128) | 2 (loc. 128) | 3 (Girty's type) |
|---|---|---|---|
| Length | 10 | 8 | 10 |
| Width | 13 | 9 | 11 |

*Observations:* As noted by Girty, this species is closely allied to *S. cornelliana* Derby, but is distinguished by its more distinct and elevated beak and larger and less numerous spines, fewer of which are reclining.

*Distribution:* Wolfcamp formation (Glass Mountains) at 87 low and 93s. Leonard formation (Glass Mountains) at 128 and D3. Word formation (Glass Mountains) at 144 and Tx. Hess formation (Sierra Diablo) at 479. Bone Canyon member? (Sierra Diablo) at 470.

### Family RICHTHOFENIDAE

#### Genus TEGULIFERINA Schuchert and Le Vene (=TEGULIFERA Schellwien)

*Diagnosis:* Shell irregular in growth, the brachial valve being round and flat and the pedicle valve more or less high and distorted, and to a greater or less extent enveloping the margins or the entire surface of the brachial valve. Brachial valve produced posteriorly into a tongue which is articulated in a notch near or slightly to one side of the apex of the other valve. Cardinal area completely lacking. Some species of the genus have a rim of inwardly directed spines just within the aperture of the pedicle valve. The outside of the pedicle valve bears long spines serving to attach the shell.

Genotype: *Tegulifera deformis* Schellwien.

*Geologic age:* Middle Pennsylvanian to Upper Permian.

*Observations:* In the Texas Permian the highest occurrence of *Teguliferina* is in the Wolfcamp formation, in the Lower Permian. The only species occurring in an Upper Permian fauna is *T. transkaukasia* (Stoyanow), which differs considerably in appearance from most species of the genus. The form described by Ivanow* as *T. rossica* certainly cannot belong to this genus. It has a cardinal area, and Ivanow concluded that an area was present in other species of *Teguliferina* but had not been recognized before. There can hardly be any doubt that other described species of *Teguliferina* do not bear an area, and it seems most natural to conclude that Ivanow had either a *Richthofenia* or some other related genus.

#### TEGULIFERINA BÖSEI King, n. sp.

Pl. XXVII, Figs. 14–16

*Description:* Shell medium-sized, having the form of a more or less broad asymmetric cone, rounded at the apex. Growth very irregular. Pedicle valve large, completely overgrowing the flat brachial valve, which is attached on one side of the cone and lies subparallel to the dorsal side of the cone. Cardinal area not present, the brachial valve being articulated by a three-cornered extension which lies in a notch on the inside of the other valve. Cardinal process small, projecting from the end of the posterior extension of the brachial valve. Pedicle valve strongly wrinkled and bearing many long slender spines. It is asymmetric because the apex of the cone is twisted up on

*This species is reported from the *Omphalotrochus* and the *Schwagerina* horizons of the Uralian. Only specimens from the latter are figured. If the identification from the former horizon is accepted, the geologic range of the genus must be extended down to include the highest part of the Upper Carboniferous.

*Sur la systématique et la biologie du genre *Spirifer* et de quelques brachiopodes de CII et CIII du gouvernement de Moscou. Bull. Soc. Nat. Moscou, Sec. Géol. 3, 1926, p. 111, figs. 4–6.

the dorsal side. Around the anterior margin of the pedicle valve there are several lamellar outgrowths on all but the dorsal side, and the mouth of the shell is encircled within on the ventral side by a thickened band which bears large, irregular, inward-pointing spines, that are largest in the center of the ventral side and diminish in size laterally. Surface marked by fine concentric lirae, spines, and very faint radial lirae. Dimensions in mm. of a specimen from loc. 93: width, 22 mm.; height, 15 mm.

*Observations:* The specimens from the *Uddenites* zone placed with this species are positively identified generically but not specifically, none being complete enough to show the details of the structure. *T. bösei* is very close to *T. deformis* (Schellwien), the genotype, differing only in the much less elongate mature form of the Texas specimens and the presence of a ring of spines inside the pedicle valve.

*Distribution:* Wolfcamp formation, *Uddenites* zone (Glass Mountains) at 88, T88, 95, and 203. Wolfcamp formation (Glass Mountains) at 71, 85, 87 low, 90, 93* (types), T93, and 193.

**Genus PRORICHTHOFENIA, novum**

*Diagnosis:* Similar to *Richthofenia* but without a myophore chamber lying along the inside of the cardinal area. Pedicle valve irregularly conical, generally twisted to one side. Cross-section nearly circular or transversely ovate. Brachial valve operculiform, when closed lying across the aperture of the cone. *Prorichthofenia* commonly grows in colonial habit, a number of specimens being attached to each other by spines and diverging from a common axis.

Pedicle valve composed of three shell layers. External layer thin, strongly wrinkled, set with numerous long, hollow spines, which rise from the internal shell layer and pass through the median and outer layers. Internal layer homologous with the outer shell of other brachiopods (the external one being a secondary investment). Medial layer thick and cystose, occupying the space between the other layers; it is thickest near the apex, becoming very thin above the plane of articulation of the brachial valve, and again thickening in a ring about the aperture.

Internal shell irregularly conical, bearing on its posterior face a cardinal area, extending from the apex of the cone to the place of articulation of the brachial valve, divided into three subequal parts by the sharp elevation of the median part and marked by transverse growth striae. This cardinal area appears to be the trace on the shell formed by the growth upward along the posterior side of the pedicle valve of the short hinge line of the brachial valve. On each side of the cardinal area the shell folds inward, defining it sharply. The posterior prolongation of the brachial valve fits into the groove so formed.

In the apical part of the ventral cone there is a median thickening of the shell on the inner anterior side, which may form a fairly high median septum. On each side of this ridge, in the posterior apical part, there is a deep pit. The diductor muscles were attached on each side of the median ridge, while the adductor scars lie along the axis of the ridge.

A narrow shelf or flange, formed by a fold of the inner shell layer, encircles the interior of the front and sides of the shell at the level of articulation of the brachial valve; it apparently formed a support for the brachial valve when it was closed. Above this shelf the interior of the shell is covered with small spines, especially on the posterior side. In addition, a few large spines rise from the anterior side and are directed upward so as not to interfere with the movement of the brachial valve. The rim of the shell bears large spines on the anterior side, which are rooted in the medial shell layer and extend inward and upward. Posterior part of the rim formed of several lamellae directed outward and upward.

Brachial valve with a short hinge-line, in front of which there is a slight constriction and then an expansion into a subcircular form. Upper surface finely spinose; lower surface papillose, with spines on the anterior and lateral margins directed downward. These spines would interlock with the spines on the rim of the pedicle valve when the shell was open. The tongue-like projection of the posterior side of the brachial valve extends into the groove below the cardinal area; the median elevation of the area incloses the cardinal process, which is small, rounded, and bilobed. From the base of the cardinal process, extend two small parallel ridges, inclosing the dendritic adductor impressions, which are divided by a median groove. A low median septum may be present in the anterior part.*

Genotype: *Crania permiana* Shumard.

*Geologic age:* Permian.

*Observations:* This genus differs from *Richthofenia*, with which it has previously been identified, in having quite different internal structure. In *Richthofenia* the groove below the cardinal area is separated from the rest of the body chamber by a partition lying immediately below the cardinal area. The tube so formed is called a myophore tube. It is generally divided in two by a median septum, and crossed at right angles by small partitions.* The importance of the difference between *Prorichthofenia* and *Richthofenia* was first noted by Likharew, who proposed that the forms lacking a myophore chamber be placed in a new genus.* In *Prorichthofenia* the muscles are attached to the median ridge in the apical part of the living chamber, without a special tube for their reception.

The species belonging to *Prorichthofenia* are:

*Prorichthofenia permiana* (Shumard) of the Delaware Mountain and Capitan formations of Trans-Pecos Texas.

*P. uddeni* (Böse) of the Leonard formation of Trans-Pecos Texas.

*This species was mentioned by Böse (U. of Tex. Bull. 55, p. 47) as "*Richthofenia*" which "resemble somewhat *R. Uddeni*," and specimens from this locality were identified by Schuchert (Am. Jour. Sci., ser. 5, vol. 14, p. 391) as "*Richthofenia permiana* (small)."

*See also Böse, E.: Contributions to the Knowledge of Richthofenia in West Texas, U. of Texas Bull. 55, 1916.

*Di-Stefano, G.: Le Richthofenia dei calcari con Fusulina di Palazo Adriano nella valle del fiume Sosio. Palaeontographica Italica, vol. 20, 1914, pp. 1–27, pls. 1–3.

Stoyanow, A. A.: On Some Permian Brachiopoda of Armenia, Mém. Com. Geol. (Russie), n. s., lov. 111, 1915, pp. 59–68, pl. 3.

Likharew, Boris: Über einige seltene und neue Brachiopoden aus dem Unterperm des nördlichen Kaukasus, Palaeontologische Zeitschrift, Bd. 10, 1928, pp. 258–289.

*Idem, pp. 282–283, fig. 13.

*P. likharewi* King of the Hess and Leonard formations of Trans-Pecos Texas and the zone of *Waagenoceras* near Las Delicias, Coahuila.

*P. teguliferoides* King, a primitive type transitional between *Teguliferina* and *Richthofenia*, of the Hess formation of Trans-Pecos Texas.

The following probably also belong to the genus:

*Prorichthofenia* sp., from the lower Permian of the Caucasis, figured by Likharew (text fig. 13) as Genus nov. sp. nov., and described by him on pp. 282–283.

*Prorichthofenia? rossica* (Ivanow) from the Government of Moscow, originally described as a *Tegulifera*. Because of the presence of a cardinal area (unlike true *Teguliferina)*, Likharew suggested that it might belong to the same group as *P. permiana.**

It has generally been supposed that the cardinal area of *Richthofenia*, *Prorichthofenia*, and *Gemmellaroia* is like that of other Protremata, with a median raised deltidium. That this cannot be true is indicated by several facts. The median raised third of the cardinal area is not defined from the lateral parts of the area by a different type of transverse striation, as is true in shells having a true raised deltidium. The outline of the brachial valve of *Richthofenia* is like that of *Teguliferina* and *Gemmellaroia:* there is a tongue-like extension of the posterior margin into a corresponding groove in the shell of the pedicle valve. In *Teguliferina* there is no cardinal area; the brachial valve merely fits into a depression on the inside of the pedicle valve, which is notched in the middle to inclose the cardinal process. If the brachial valve of a *Teguliferina* type had shifted its position along the posterior inner wall of the pedicle valve it would form a cardinal area on the inside of the inner layer, the extension of the cardinal process forming a median groove having the appearance of the inner side of a deltidium. This is the situation in *Richthofenia* and *Prorichthofenia.* The cardinal area is crossed by continuous growth lines indicating successive positions of the brachial valve. Similarly, in *Gemmellaroia* the growth lines on the cardinal area are continuous with the growth lines on the rest of the shell. Gemmellaro stated that in this genus there is a raised "pseudodeltidium" but that an area was lacking. The "pseudodeltidium" is actually homologous to the entire area of *Richthofenia* and *Prorichthofenia,** there being no median elevation in *Gemmellaroia* formed by extension of the cardinal process into the posterior wall during upward growth of the brachial valve (as in the "deltidium" of Richthofenia).

The presence of a myophore chamber is taken by Likharew* to be the essential character of the family Richthofenidae. In view of the agreement of the other shell characters it seems quite unnatural to separate those forms having a well developed myophore chamber in another family from those without it. Moreover, as has been shown, it is hardly to be doubted that *Teguliferina* is the ancestral form of the family, yet because of its structure, a myophore chamber could not occur in that genus.

Stoyanow* regarded as the essential character of his subfamily Richthofenidae the presence of a ventral median septum. Likharew* has clearly shown that the median septum of *Scacchinella* is of quite different nature from that occurring in the myophore chamber of *Richthofenia, Gemmellaroia,* and *Tectarea.* To me it appears that the important characters uniting the related genera *Teguliferina, Prorichthofenia, Richthofenia, Gemmellaroia,* and *Tectarea* into the family Richthofenidae are the high, deformed pedicle valve, the posterior tongue-like projection of the brachial valve, which fits into a notch in the pedicle valve, the presence in all but *Teguliferina* of a cardinal area, and the long anchoring spines. *Teguliferina, Prorichthofenia,* and *Richthofenia* have an external shell layer which envelops the entire surface, while *Gemmellaroia* and *Tectarea* are only slightly enveloped. The arrangement of the adductor muscles of the brachial valve is somewhat different in *Gemmellaroia* than in *Richthofenia* and *Prorichthofenia;* in the former they are simple and enclosed by apophyses, rather than broad and dendritic. The most highly specialized internal characters are those of *Tectarea:* in it there is a very long massive cardinal process which extends far into the myophore chamber; from the base of the process extend two diverging ridges.

It seems likely that *Prorichthofenia, Richthofenia, Gemmellaroia,* and *Tectarea* evolved from the primitive form *Teguliferina,* which in turn probably arose from some member of the subfamily Strophalosiinae.

**PRORICHTHOFENIA LIKHAREWI King, n. sp.**

Pl. XXVIII, Fig. 1; Pl. XXIX, Figs. 1–5; Pl. XXX, Fig. 1

1915 *Richthofenia permiana.* Haack, Zeit. der Deutsch. geolog. Gesellsch., pp. 491–482, pl. 38, figs. 6a–b. Pichagua, near Las Delicias.

*Description:* Shell similar in external configuration to *P. permiana,* though generally smaller. The cystose filling of the apical part of the cone is, however, much thicker; it fills the lower part of the cone, in some instances, to a depth of more than 15 mm. There is a great thickening in the median part of the anterior wall, which is produced toward the posterior side into a high, sharp septum, directed toward the cardinal area. Unlike the situation in *P. uddeni* this septum does not reach the posterior side of the shell. In some specimens the septum extends upward along the anterior side of the shell nearly to the shelf on which rests the brachial valve, but in others it occurs only in the apical part of the living chamber, the rest of the anterior wall being regularly rounded. On each side of the septum in the apical part of the shell there is a deep pit. The muscles must have been attached on each side of the septum and in the pits on each side. The inner surface of the cardinal area lies exposed to the living chamber. The other characters are as in *P. permiana.* One large piece of a colony from loc. 128 consists of about 50 specimens of this species grown together and attached by spines; the apices of the shells are directed toward a common center, as in a colonial coral.

Dimensions in mm. of several specimens:

*Idem, p. 283, note.

*See also Likharew, *idem,* p. 279.

**Idem,* p. 282.

*Mém. du Com. Géol. Russ., n. s., liv. 111, p. 74.

***Idem,* pp. 281–288.**

| | 1 (loc. 128) | 2 (loc. 128) | 3 (loc. 123) | 4 (loc. 120) |
|---|---|---|---|---|
| Length of pedicle v. | 41 | 30 | 31 | 31 |
| Length of brachial v. | 12 | 18 | 16 | 16 |
| Width of brachial v. | 21 | 17 | 22 | 11 |

*Observations:* This species appears to be ancestral to *P. permiana,* the main differences between them being the greater development of the cystose filling of the apex and the great thickening and extension into a septum of the anterior wall of the living chamber.

*Distribution:* Hess formation (Glass Mountains) at 1?, 4, 8, 17b, 35, 103, 106, 107, 117?, 211, 223, and 226. Leonard formation (Glass Mountains) at 3, 8, 11a, 11b, 7–14, 12, 13, 16, 23, 26s, 31, 34, 37, 38, 39, T41, 83, 102?, 104, T119, 120 (abundant), T120, 123 (abundant), 124 (abundant), 126, 128 (abundant; types), 228, 231, 232, Tb, Ax, D2, D3, D7, and D22. Hess formation (Sierra Diablo) at 469 and 479. Bone Canyon member (Sierra Diablo) at 491. Leonard formation (Baylor Mountains) at 502. Leonard formation, just above brecciated zone (Shafter) at Td. Limestone of Pichagüia (Las Delicias ) at Cerro Pichagüia.

### PRORICHTHOFENIA PERMIANA (Shumard)

Pl. XIX, Figs. 6–9; Pl. XXX, Figs. 8–13

1859 *Crania Permiana.* Shumard, Notice of fossils from the Permian strata of Texas and New Mexico, etc., Trans. Acad. Sci. St. Louis, vol. 1, p. 395 (date of volume, 1860).

White (Permian) limestone, Guadalupe Mountains.

1908 *Richthofenia permiana* (ex parte). Girty, The Guadalupian Fauna, U. S. Geol. Sur. Prof. Paper 58, pp. 283–286, pl. 14, figs. 27–27d; pl. 20, fig. 23; pl. 22, figs. 6–6b; pl. 31, figs. 2–3.

Middle Capitan, Guadalupe Point (sta. 2926); dark limestone, Pine Spring (sta. 2930) and Guadalupe Point (stas. 3762b and 3762c); Delaware Mountain formation, Guadalupe Point (sta. 2931); Delaware Mountain formation, southern Delaware Mountains (stas. 2964 and 2969); and Delaware Mountain formation, "Comanche Canyon," Glass Mountains (sta. 3763). (Not from stas. 2920, 2967, and 3763 part.)

1916 *Richthofenia permiana* (ex parte). Böse, Contributions to the knowledge of Richthofenia in the Permian of West Texas, U. of Tex. Bull. 55, pp. 36–42, pl. 1, figs. 1–16; pl. 2, figs. 8, 15; pl. 3, figs. 1, 4–7.

Zone of *Medlicottia* n. sp. (Word), Gilliam Canyon, mountains north of Leonard Mountain, Hess Tank, Word Ranch, Mt. Ord Range. (Not from the zone of *Waagenoceras* n. sp. (Leonard).

*Description:* Pedicle valve long, narrow, and irregularly conical; transverse section elliptical, being longer in the lateral axis; longitudinal axis generally curved, being convex toward the anterior side. Surface covered by many long hollow spines, generally projecting downward away from the shell, which serve for attachment to other Richthofenids or other shells; where the individuals grow in groups one shell may lie against two or three others and bear spines only on the free surfaces. Outer layer strongly rugose and marked by concentric growth lines. The cystose filling of the lower part of the cone is thickest at the apex and along the median part of the anterior wall in the apical region, but even there is relatively thin, being not more than a few millimeters thick even in large shells. Along the posterior side is the long, narrow cardinal area, depressed in the median part in a long concavity. Living chamber generally equal in length to about .8 the length of the valve; it is bordered above by the shelf on which rests the brachial valve. The inner side of the cardinal area is exposed to the living chamber except near the apex, where it is more or less covered by the secondary cystose filling of the apical region. Opposite the apical part of the cardinal area there is considerable later deposit on the inner anterior wall. It rises along the median line to form a low, broad ridge, bearing the elongate scars of the adductor muscles along the top and of the diductor muscles on the sides. The inside of the living chamber is marked by scattered tubes which lie against the shell wall and open out in the direction of the aperture. Above the rim on which rests the brachial valve the inner wall of the pedicle valve is thickly set with upward-projecting spines, which are larger on the anterior side, and increase in size toward the aperture, which is bordered by a ring of long, longitudinally striated spines.

Brachial valve flat and round, with a posterior projection into the notch in which lies the cardinal area. The inner side of the valve bears many small spines in the anterior and lateral parts. Cardinal process small and bilobed, projecting into the groove in the middle of the cardinal area. In front of it lie the dendritic adductor muscle scars.

Dimensions in mm. of several specimens:

| | 1 (loc. 144) | 2 (loc. 144) | 3 (loc. 144) | 4 (loc. 247) |
|---|---|---|---|---|
| Length of pedicle v. | 39 | 34 | 48 | 43 |
| Length of brachial v. | 18 | — | 22 | 19 |
| Width of brachial v. | 22 | 18 | 28 | 23 |

*Observations:* This species differs from *P. likharewi* in having a low, broad median ridge rather than a high sharp septum, and in much less development of apical cystose tissue. It is distinguishable from *P. uddeni* by its more slender form and the absence of sharp median ridge.

*Distribution:* Word formation (Glass Mountains) at 44–46?, 46, 54, 55, 135, 136, 138, 139, 142, 144 (extremely abundant), T144a, T144, 145, 146, 148, T148, Tx148, 152, 153, 160, 171, 192, 232, T233, T238, T239, D239, T240, 241, 243, T243, 245, 247, 265, Tc, and Tx. Limestone of Cerro Caballo (Las Delicias) section 1, bed 12; section 2, bed 12, between El Tordillo and Cerro Caballo.*

### PRORICHTHOFENIA TEGULIFEROIDES King n. sp.

Pl. XXX, Figs. 2–7

*Description:* Shell small. Pedicle valve having the form of a very asymmetric cone, with the apex twisted toward the posterior side. The anterior side is very long and convex outward, while the posterior side is about a third as long and concave outward. On the posterior side there

*The specific determination of these Las Delicias specimens is uncertain, as the apical parts are not preserved.

is a short cardinal area, at the top of which the flat brachial valve is attached; the posterior articulating extension of the brachial valve fits into the groove of the inside of the cardinal area. The brachial valve rests on a rim extending around the inside of the marginal part of the pedicle valve, its plane lying at about 45° to the main axis of the pedicle valve. There is no conspicuously developed apical callosity or internal septa. A row of spines lies about the aperture on the anterior side. The outside of the shell is covered by numerous round, prostrate anchoring spines. Cardinal process of the brachial valve extended back from the cardinal margin as a narrow ridge with two short lobes at the end. Dimensions of one specimen from loc. 479: length of anterior side, 18 mm.; of posterior side, 6? mm.; width of brachial valve, 15 mm.; length of brachial valve, 12 mm.

*Observations:* This species is intermediate in structure between *Teguliferina* and the more specialized forms of *Prorichthofenia.* The cardinal area is very short, a fact which indicates that the migration of the point of articulation of the brachial valve from the apex (as in *Teguliferina)* to a point high on the posterior side of the shell (as in *Prorichthofenia likharewi,* etc.) was only the beginning. Were it not for the presence of the small cardinal area, it would be placed in *Teguliferina,* for its general aspect is the same and there is no distinctly developed apical callosity or internal septa.

*Distribution:* Hess formation (Glass Mountains) at 105, 207, 208, and 222. Hess formation (Sierra Diablo) at 475, 476, 478, and 479 (abundant, types). Hess formation (Finlay Mountains) at 511.

**PRORICHTHOFENIA UDDENI (Böse)**

Pl. XXXI, Figs. 1–6

?1908 *Richthofenia permiana* (pars). Girty, The Guadalupian Fauna, U. S. Geol. Sur. Prof. Paper 58, pp. 284–285, pl. 24, figs. 10–10a; pl. 31, figs. 1–1a.
Basal black limestone, Guadalupe Point (Stas. 2920 and 2967). Delaware Mountain formation, "Comanche Canyon," Glass Mountains.

1916 *Richthofenia Uddeni.* Böse, Contributions to the knowledge of Richthofenia in the Permian of West Texas. U. of Texas Bull. 55, pp. 43–48, pl. 2, figs. 1–7, 9–14, 16; pl. 3, figs. 2–3.
Zone of *Waagenoceras* n. sp. (Leonard); ? zone of *Medlicottia* n. sp. (Word).

*Description:* Pedicle valve broadly conical; apex pointed and turned toward the posterior side. Transverse section transversely oval. Outer surface rugose and marked by strong growth lines; it gives rise to many long hollow spines, projecting generally downward or laterally for attachment to a foreign body. Median layer thickest in the apical region, especially along the middle of the anterior side; it is not, however, more than a few millimeters thick even in large shells. Directly below the aperture there are several sharp flanges marking successive growth stages; at the aperture the shell flares out in a broad thin sheet. Living chamber generally equal in length to about two-thirds the depth of the valve; it is bordered above by the shelf on which rests the brachial valve. The inner side of the cardinal area is not exposed to the living chamber, being covered by a later deposit of the mantle. In the median apical part of the living chamber there is a broad, rounded elevation, extending from the posterior side across the bottom of the chamber and along the median anterior surface, where it gradually decreases in elevation but is distinguishable as far up as the shelf. The top of the ridge gives rise on the posterior side to a sharp median septum, which connects with the later deposit covering the inside of the cardinal area, and rises halfway up the posterior side of the shell toward the place of articulation of the brachial valve. On each side of the septum, directly within the fold of the inside of the cardinal area, there is a well-defined pit; these pits and the sides of the septum were evidently the places of attachment of the muscles. Several small, upwardly directed spines are present inside the living chamber. Immediately above the shelf on which rests the brachial valve, on the anterior and lateral inner surface of the shell, there are ridges which project upward, and at the aperture give rise to thick, bifurcating spines, which are directed upward and inward toward the axis of the shell.

Brachial valve flat, its outline corresponding to the cross-section of the upper part of the living chamber.

Dimensions in mm. of several specimens:

| | 1 (loc. 128) | 2 (loc. 128) | 3 (loc. 120) | 4 (loc. 124) |
|---|---|---|---|---|
| Length of pedicle v... | 27 | 22 | 27 | 45 |
| Length of brachial v... | 25 | 17 | 18 | — |
| Width of brachial v... | 25 | 20 | 23 | 28 |

*Observations:* That this species is a primitive type is indicated by its broadly conical form, short cardinal area, and single, rather than multiple rim of spines about the aperture. These features, and likewise the presence of the median septum on the posterior side (rather than on the anterior side as in *P. likharewi)* serve to distinguish it from the other species in the present fauna.

The specimens identified by Girty as *Richthofenia permiana* from the basal black limestone at Guadalupe Point, as well as one of Girty's specimens from the Glass Mountains, appear to agree in outer configuration with *P. uddeni.* Since the internal structures were not seen in some, and not well enough preserved in others to distinguish them from *P. permiana* conclusively, they are questionably referred to *P. uddeni.* It may be noted that *P. permiana* would not be expected to occur at the horizon of the basal black limestone (Bone Canyon member).

*Distribution:* Leonard formation (Glass Mountains) at 7–14, 20, 21, 24, 91x, 120 (good silicified specimens), 121, 128 (common), and Tb. Word formation (Glass Mountains) at T251. Bone Canyon member (Sierra Diablo) at 485. Leonard formation, transition beds (Shafter) at Th (15963—figured by Böse, pl. 2, figs. 10, 11, 14); above transition beds (Shafter) at Ti (16009—figured by Böse, pl. 2, figs. 12, 16).

## Family LYTTONIIDAE

### Genus PARAKEYSERLINGINA Fredericks (=EOLYTTONIA Fredericks)

*Diagnosis:* Shell concavo-convex, hinge-margin narrow. Pedicle valve gently and irregularly convex, expanding

over the beak. Outer surface marked by uneven concentric growth lines; shell bilaminate, the external layer being fibrous, the internal one punctate. Within the pedicle valve there is a complexly lobate septal ridge corresponding to the brachial ridges which are supposed to be present in the brachial valve. There is a narrow median invagination of the ridge standing back from the front margin nearly to the beak, and 9 to 12 or more lateral lobes, which lie at a right angle to the median lobe or are directed somewhat backward. The septal ridge or fillet is distinctly looped, and does not, as in *Lyttonia* and *Oldhamina*, have the appearance of septa connected at their ends by loops; this is due to the feeble development of the invagination of the outer shell layer which in *Lyttonia* and *Oldhamina* rises between the septal ridges to form the axes of the septa. Brachial valve unknown, but supposedly like that of Lyttonia.

Genotype: *Keyserlingina darvasica* Tschernyschew.*

*Geologic age:* Lower Permian (Schwagerina zone of Darwas, Bokhara; lower Permian of Ussuriland; Wolfcamp formation).

*Observations:* This genus occupies an intermediate position between *Keyserlingina* and *Lyttonia.* From the former it differs in having a greater number of lateral lobes, which are longer, are bent toward the lateral margins, and lie closer together. From the latter it is distinguished by having fewer lateral lobes, which lie farther apart, and in having the median and lateral lobes not united in pairs to form the septa by the development of axes of the outer shell layer which project up between the lobes as in *Lyttonia.*

Fredericks has proposed two generic designations for species which appear to have the same diagnostic characters. In 1916* the name *Parakeyserlingina* was introduced for *Keyserlingina darvasica* Tschernyschew, while in 1923* he introduced the name *Eolyttonia* for *Oldhamina mira* and *O. ivanovi* Fredericks. The differences indicated between them hardly seem to be of generic value. The one occurs in the basal Permian of Bokhara and the other in beds of Artinskian age in Ussuriland (southeastern Siberia); it therefore seems likely that they lived synchronously between the time of *Keyserlingina* and that of *Lyttonia* and *Oldhamina.* Because of the little difference, if any, between the two, it seems best to recognize only the first named—*Parakeyserlingina,* although *Eolyttonia* would be a more satisfactory name.*

According to Fredericks* the development of the Lyttoniidae took place in the following manner:

The ancestor of the family is supposed to be *Marginifera.* In that genus Fredericks believes the marginal fold to be due to abnormal development of the dental plates ("apical plates") resulting from reduction of the cardinal area. Strong transverse development of the fold leads to the formation of a closed ring as in *Paramarginifera* Fredericks (genotype *M. clarkei* Tschernyschew). This ring is said to be analogous to the spondylium of brachiopods having well-developed cardinal areas. Further development of the ring leads to invagination of the median anterior part, forming two lobes: this stage is that of *Pseudokeyserlingina* Fredericks (hypothetical). With the development of a pair of lateral invaginations of the marginal ridge we have *Prokeyserlingina* Fredericks (hypothetical), and with two more pairs, *Keyserlingina* Tschernyschew. Simultaneous with these changes comes the reduction of the brachial valve until it covers only the area inclosed by the marginal ridge. *Poikilosakos* Watson is thought to be an offshoot from the main stem of the evolution of the family, for it has greatly reduced muscle scars, one of which is more strongly developed than the other, and the marginal fillet is frilled. Further development of this line results in the formation of a greater number of lateral loops of the marginal fillet, which come to lie closer together. In *Parakeyserlingina* the invaginations of the marginal ridge are very long and narrow, but are grooved along the axis, so that they do not form a septa as in *Lyttonia. Lyttonia* and *Oldhamina* are still further specialized: they have a greater number of lateral invaginations, which are very narrow and have in the axial part of each a thickening of the outer shell layer ("interlobal callosity" of Fredericks) which rises between the parts of the marginal ridge on opposite sides of the invagination (and also in the median invagination of the shell) until it becomes higher than the bordering ridges and forms a septum. As a result the inside of the pedicle valve of *Lyttonia* and *Oldhamina* have numerous lateral septa and a long median septum, which are connected at their distal ends by loops; the original structure, that of an invaginated ridge, is thus obscured.

Whatever may be the truth of the supposed evolution from *Marginifera* to *Keyserlingina,* the evolution from *Keyserlingina* to *Lyttonia* and *Oldhamina* is well substantiated in both European and American forms of this group.

---

*Described and figured in: Fauna der Oberpalaeozoischen Ablagerungen des Darwas, Mém. Com. Géol. (Russie), n. s., liv. 104, 1914, pp. 13–14, 49.

*Mém. Com. Géol., (Russie), n. s., liv. 156, p. 67.

*Rec. Geol. Com. Russian Far East, no. 28 (1924), pp. 22, 25. In this reference the name *Eolyttonia* is said to have been proposed in 1922, but the original reference is not stated.

*Fredericks regards *Lyttonia tenuis* Waagen as an *Eolyttonia,* relying evidently on Waagen's few figures of that species. This hardly seems to be justified, for without actual specimens for study it is very easy to make errors in interpreting the structure of the septal fillet. The specimen figured by Kindle (Trans. Roy. Soc. Canada, ser. 3, vol. 20, sect. 4, p. 110) from British Columbia, under the name of *Leptodus* cf. *tenuis,* has the appearance of a *Parakeyserlingina,* as each septal fillet appears to consist of two ridges separated by an axial groove, but a definite judgment on this point is not possible.

*On the subfamily *Lyttoniinae* Waagen: Mém. du Com. Géol. (Russie), n. s., liv. 156 (1916), pp. 55–79. In Russian.

Addenda to the Dissertation of G. N. Fredericks, "Paleontological Notes. 2, Some Upper Paleozoic Brachiopoda of Eurasia," presented in candidacy for the degree of Master of Mineralogy and Geognosy, pp. 1–2. 1917. In Russian.

Upper Paleozoic of the Ussuriland. I. Brachiopoda. Rec. Geol. Com. Russian Far East, no. 28 (1923), pp. 22–27. Vladivostok, 1924. In Russian.

Upper Paleozoicum of the Ussuriland. II. Permian Brachiopoda of Cape Kalouzin. Rec. Geol. Com. Russian Far East, no. 40 (1924), pp. 11–16. Vladivostok, 1925. In Russian.

Note sur l'origine de l'appareil septal de *Lyttoniinae,* Ann. Soc. paléont. Russie, vol. 3, pp. 108–110. In Russian.

New Lyttoniinae from upper palaeozoicum of the Oural, Bull. Soc. Oural. Sci. Nat., pp. 59–65. In Russian.

New Lyttoniinae from the Upper Carboniferous of Krasnoufimsk, pp. 83–89, 1927? In English.

In this connection it is necessary to consider the genus *Uralia* Likharew* *(Uralina* Schuchert and LeVene). This is based on what is supposed to be the brachial valve of a lyttoniid. This shell has an entire margin and a rounded outline; there is a median ridge and seven pairs of transverse folds. Likharew believes that in all the early forms of the Lyttoniidae the brachial valves had entire margins, and that the splitting of the shell into lobes occurred only in *Lyttonia* and *Oldhamina.* He conceives the brachial valve of the more primitive forms to be like that of *Uralina,* each fold in the brachial valve overlying a lobe of the marginal ridge in the pedicle valve. *Uralina* is thought to be a stage intermediate between *Keyserlingina* and *Lyttonia,* and to include probably *Oldhamina (=Eolyttonia) mira* Fredericks. If Likharew's supposition as to the probable structure of the brachial valve of the early lyttoniids were correct, therefore, *Uralina* would be a synonym of *Parakeyserlingina.* However, although brachial valves of the early lyttoniids have not been found, it has been argued convincingly by Watson* and others that the brachial valve covered only the space within the marginal ridge and that when closed its edge rested on the inner face of this ridge, leaving the margins of the pedicle valve uncovered all around. If it is true that the brachial valve of these forms was lobate, then *Uralina* can hardly belong to this group at all. The type specimen, indeed, can hardly be regarded as adequate material to make definite assignment to this or any other invertebrate group.

**PARAKEYSERLINGINA FREDERICKSI King, n. sp.**

Pl. XXXI, Figs. 7–9

*Description:* Shell large, very irregular. Pedicle valve unevenly convex; one specimen from T93 has the umbonal region greatly expanded laterally and strongly convex, arching over the dorsal side to form a cup. Shell bilaminate, the external layer being marked on the outside of the shell by fine, irregular growth wrinkles, while the internal layer forms the complexly fluted septal fillet. The fillet is infolded in at least 11 pairs of lateral lobes* and a long, narrow median lobe. The lateral lobes may be inclined at about a right angle to the median lobe, but generally slant somewhat forward. In the specimen from loc. T93 above referred to (the only one in which the posterior part is preserved) the first 5 lobes are inclined forward at angles as high as 50°, and are widely spaced from each other, while the later ones lie at nearly right angles. This may indicate a repetition of phylogeny in ontogeny, for in the earlier forms *Keyserlingina* and *Poikilosakos* the lateral lobes are directed forward. Each lobe is very narrow and has a groove along its axis. The inward terminations of the lobes are high and sharp; their elevation subsides distally. The vertical axes of the lobes are inclined forward at a greater or lesser angle. The outer margin of each lobe is crenulated by low vertical ridges, each about 1 mm. wide; these extend more or less interruptedly across the space between adjoining lobes. Between each lobe and parallel to the lobes there is a low broad ridge. Adjoining lobes are connected distally by the broad, low loop of the septal fillet. The median lobe of the shell is narrow and grooved in the median line just as the lateral lobes.

Brachial valve unknown, unless a brachial valve of a lyttoniid from loc. 15, having pinnate lobes lying at about 45° to the median axis, belongs to this species.

*Observations:* The specimens from loc. T93 have been referred to in several places as *Lyttonia.** The genus *Lyttonia* is not known to occur in the Wolfcamp formation. This species cannot be satisfactorily compared with the other three members of the genus, *P. mira* (Fredericks), *P. ivanovi* (Fredericks), and *P. darvasica* (Tschernyschew), for none of these occurs in any abundance, and the few specimens figured show considerable variation. The closest resemblance appears to be with *P. ivanovi.*

*Distribution:* Wolfcamp formation (Glass Mountains) at 71?, 87 low, 93, T93 (types), and 194?. Hess formation (Glass Mountains) at 15?. Leonard formation (Glass Mountains) at 19?.

**Genus LYTTONIA Waagen***

*Diagnosis:* Shell flattened irregularly, sprawling in growth, in youth attached by the beak of the pedicle valve, but later becoming free, the point of attachment becoming covered by a strong, irregular, lamellose growth. Pedicle valve gently convex, nearly flat. On the interior of the pedicle valve there is a low but thick median septum and as many as 40 pairs of thick, pinnately arranged, rounded lateral septa, which are directed transversely and are somewhat convex toward the front; they do not unite with the median septum, but are united at their distal ends by loops of the septal fillet. The septa consist of parallel outer ridges of the inner shell layer (the septal fillet) separated by an elevated axial part formed by a thickening of the outer shell layer ("interlobal callosity" of Fredericks).

Brachial valve smaller than the pedicle valve, divided into two halves by a median slit extending back from the front margin, and again divided by lateral incisions from each side into a number of smaller lobes; the lobes are slightly convex toward the front and correspond to the spaces between the septa. Internally there is a strong median septum, which probably ends forward in the longitudinal slit, and terminates at the hinge-line in an atrophied cardinal process.

Cardinal area, delthyrium, hinge-teeth, and crura are completely lacking; dental lamellae are present but very small and thin. Muscular impressions of the pedicle valve very weak and indistinct; in the brachial valve not distinguished. Shell bilaminate, the outer layer of the pedicle valve being smooth and the inner granular.

It is probable that each pinna of the brachial valve supported a double band of lophophore extending its entire length, and that the median septum was provided with a strip of lophophore on each side. Each pinna of

*Licharew, B.: Über einen neuen Vertreter de Fam. *Lyttoniidae* aus dem Obercarbon des Ural. Comtes Rendus de l'Acad. des Sc. de Russie, 1925, pp. 4–7, text fig.

*Geol. Mag., n. s., vol. 4 (1917), p. 215.

*The word "lobe" is here used for the inward-directed fold of the septal fillet.

*As in U. of Tex. Bull. 1762, p. 16.

*Editor's Note: The name *Leptodus* Kayser has priority and Kindle (**89**, p. 109) has shown that Waagen was not justified in rejecting it and substituting the name *Lyttonia.*

the brachial valve rests along its anterior edge on the side of a septum of the pedicle valve, and ends distally at the loop connecting adjoining septa; the cavity of the shell thus forms a series of canals leading to a pair of central channels running antero-posteriorly. Each lateral canal is open to the outside by a slit running the length of the dorsal pinna, formed by the gap between that pinna and the one behind it. Probably the ciliary currents were driven into these lateral canals and by the action of the strip of lophophore on the side of the median septum moved toward the posterior end.*

Genotype: *Lyttonia nobilis* Waagen.

*Geologic age:* Middle and upper Permian.

*Observations:* The following are the differences between *Oldhamina* and *Lyttonia: Lyttonia* is ovoid in outline and shallow because little arched, whereas *Oldhamina* is more arched and subhemispherical, its beak being short and incurved. There is a well developed apical callosity in *Oldhamina*, while in *Lyttonia* it is only slightly developed. The septa in *Oldhamina* are turned strongly forward, while in *Lyttonia* they diverge from the median line at nearly a right angle. *Lyttonia* differs from *Parakeyserlingina* in having a greater number of lateral septa, which have a thickened and elevated, rather than a grooved axial part.

Several species of this genus have been recognized, but Noetling and later Hayasaka have shown that they are probably only variations of one highly variable species.

### LYTTONIA NOBILIS AMERICANUS (Girty)

Pl. XXXI, Figs. 10–13; Pl. XXXII, Figs. 1–9

1908 *Leptodus americanus.* Girty, The Guadalupian Fauna, U. S. Geol. Sur. Prof. Paper 58, pp. 212–213, pl. 4, figs. 8–8b, pl. 25, figs. 1–3a.

Delaware Mountain formation, Guadalupe Point (sta. 2931). Delaware Mountain formation, Diablo Mountains (sta. 3764). Delaware Mountain formation, southern Delaware Mountains (sta. 2969).

1908 *Leptodus guadalupensis.* Girty, *idem.*, pp. 213–215, pl. 4, figs. 6–7.

Middle Capitan, Capitan Peak (sta. 2926). Basal Capitan, hill southwest of Guadalupe Point (sta. 2906). Delaware Mountain formation, "Comanche Canyon," Glass Mountains.*

*Description:* Shell large, concavo-convex, with an ovoid outline, the posterior part being narrower than the anterior. Pedicle valve slightly and irregularly convex, with entire margin. In the posterior part the margin of the pedicle valve may grow up and arch over the posterior part of the brachial valve, or turn back to lie on the outer posterior part of the shell; the inside of this growth is thickly set with granulations. As a result the umbonal region has an irregular, gnarly aspect. Outside of the shell smooth except for strong, uneven growth lirae and low concentric undulations, which do not parallel the course of the septa within the shell. In the posterior part of one specimen (from loc. 476) there is a broken area near the posterior margin which the growth lines appear to encircle, these lines being closely crowded posterior to the broken area but separated from one another in broad curves toward the sides and front of the shell; it may be that this broken area was a point of attachment during the first part of the growth of the shell. Another specimen (from loc. T240a) appears to have had a similar point of attachment which has been overgrown by the laminar outgrowth of the posterior margin, as has been described by Noetling for *L. nobilis.* An immature specimen from loc. 54 is attached to the concave surface of a brachial valve of *Marginifera popei.* (See also Girty, pp. 214–215, pl. 4, figs. 8–8b.) In a specimen from loc. 128 the posterior lateral margins have grown around the dorsal side of the shell so as to give the posterior part a cup-like form.

Within the pedicle valve there are at least 20 pairs of lateral septa, which are arranged pinnately on opposite sides of a median septum and opposed in pairs. In plan these lateral septa arch slightly toward the front and in section they are inclined forward. They are high and sharp at their proximal ends, but are low at the distal ends, where they are connected with the adjacent septa by a loop of the marginal ridge. An immature specimen from loc. 54 shows the first invagination of the septal fillet to be distinctly double, and divided along the axis by a deep groove. The second invagination appears to be a true septum, the space between the two sides of the septal fillet being filled by an "interlobal callosity." In the specimen from T240a the first four septa are distinctly double, as in *Parakeyserlingina,* and it appears thus to repeat in its ontogeny the phylogeny. The septa increase in length from the umbonal region toward the front margin, corresponding to the increase in the width of the shell. Separating the septa are regularly disposed interspaces, each about 2 mm. broad, which bear a broad median longitudinal ridge, conspicuously lower than the adjacent septa but strongly enough developed so that a prominent groove lies on each side of it; this structure is especially noticeable in molds of the inside of the pedicle valve, in which the grooves lying against the septa have the appearance of a continuous looped ridge which might be mistaken at first for the septal fillet (See Girty's plate 4, fig. 6, for example).* In the bottom of each of the grooves lying beside the septa, there is a row of regularly arranged pits, about 0.75 mm. apart, which give rise to shallow longitudinal depressions extending across the interseptal space and terminating at the crests of the septa, to which they give a strongly crenulate aspect.

A median septum extends nearly the length of the inside of the valve; it is slightly lower than the proximal ends of the lateral septa, but is of about the same width.

Brachial valve pinnate, smaller than the pedicle valve and somewhat concave. Its lateral margins are deeply

*See Noetling, Fritz: Untersuchungen über die Familie Lyttoniidae Waagen emend Noetling, Paleontographica, bd. 51, pp. 129–153, pls. 15–18.

Also, Watson, D. M. S.: *Poikilosakos,* a remarkable new genus of brachiopods from the Upper Coal-measures of Texas, Geol. Mag., n. s., vol. 4 (1917), pp. 216–219.

*The last two localities are listed under the head of "horizon and locality" for *L. americanus,* but Girty states clearly in the text (pp. 214–215) that he had placed the Glass Mountains and Capitan specimens with *L. guadalupensis.*

*Confusion as to the actual nature of this structure led Fredericks to the conclusion that Girty's two species were *"Eolyttonia"* rather than true *Lyttonia.*

split by lateral incisions, so that the shell is divided into a series of small lobes corresponding to the interseptal spaces of the other valve. There is a broad median axis, narrowing toward the front, which is covered on the outside by coarse granulations; in the median line there is a lightly impressed furrow, on the outside, which corresponds to a median ridge on the inside. This furrow deepens toward the front and ends in a slit near the anterior margn. The lateral lobes bend downward on their posterior and anterior sides, so that they are semicircular in transverse profile. As a result, they are concave on the inner side.

Dimensions in mm. of several specimens:

| | 1 (loc. 158) | 2 (loc. 4) | 3 (loc. 152) | 4 (loc. Tx148) | 5 (loc. 128) |
|---|---|---|---|---|---|
| Length | 60 | 45 | 45+ | 50+ | 42 |
| Width | 60 | 70 | 80 | 45 | 40 |

*Observations:* The only difference that can be made out between *L. nobilis americanus* and typical *L. nobilis* from the Salt Range is the presence of a long, well developed median septum in the pedicle valve of the former. In the Indian type a median septum occurred in the anterior part of the shell and is supposed to have been resorbed in the posterior part during the growth of the shell. The other distinguishing features mentioned by Girty, as well as those differentiating *L. americanus* and *L. guadalupensis* have been well shown by Hayasaka* to be unreal, and to rest either upon the imperfection of Girty's material or upon a misunderstanding of the actual structures observed. Whether the American form should not be considered a variety of *L. richthofeni* (Kayser) rather than of *L. nobilis* is a matter of some doubt. Hayasaka and others regard the two as synonymous, but since there is no such complete description of the Chinese species as there is of *L. nobilis* in Noetling's monograph, I have thought it best to compare the American form with the better known Indian type.

Girty's identification of *Oldhamina?* from the Capitan is certainly very doubtful, for many specimens of *Lyttonia* in North America are more or less convex longitudinally but do not show the characters of *Oldhamina.*

*Distribution:* Hess formation (Glass Mountains) at 4, 102, 103, 105, 106, 108, and 222. Leonard formation (Glass Mountains) at 8–14, T9, 11b, 15, T15, 24, 26, 31, 33, 37, 38, 81, 120, 121, 123, 124, 128, T183, Tb, and D1. Word formation (Glass Mountains) at 6, 32, 48, 51, 54, 55, 58, T140, 144, T144, 145, 146, 147, 148, T148, Tx148, 152, 154 158, 171, 192, 238, 239, T239, T240a, 244, 247, 249, 250, 251, 264, 265, Ta, Tc, and Tx. Altuda member (Glass Mountains) at 57. Hess formation (Sierra Diablo) at 476, 478, and 479. Bone Canyon member (Sierra Diablo) at 484 and 491. Victorio Peak member (Sierra Diablo) at 493 and 499. Delaware Mountain formation (Sierra Diablo) at 504. Limestone of Cerro Caballo (Las Delicias) at section 1, bed 11 west of the Noria; section 1, bed 15; and section 1, bed 12.

*Paleozoic Brachiopoda from Japan, Korea, and China, Sci. Repts. of Tôhoku Imperial University, ser. 2, vol. 6, no. 1, pp. 106–107.

**LYTTONIA HORTONI King, n. sp.**

Pl. XXXIII, Figs. 18–19

*Description:* Shell large, concavo-convex, outline rounded, expanding toward the anterior end. Within the pedicle valve there are as many as 20 pairs of lateral septa, convex toward the front, arranged opposite one another on opposite sides of the median septum. The septa have about the same height throughout their entire length; they are crowned with elongated nodes 2 to 3 mm. long, separated from each other by narrow transverse saddles. The nodes consist of inward extensions of the external shell layer, while the intervening saddles are covered across with the internal shell layer. In one specimen one of the septa bifurcates about 5 mm. distal to the median line of the shell. Median septum about the same height as the lateral septa, and interrupted in the same manner.

Brachial valve pinnately split, but with dissepiments across the interlobal slits corresponding to the saddles lying between the dashes of the septa within the other valve. A groove lies in the median line, above the median septum of the pedicle valve.

Dimensions in mm. of two specimens:

| | 1 (loc. 7) | (2 (loc. 220) |
|---|---|---|
| Length | 60+ | 60 |
| Width | 110 | 97 |

*Observations:* Since this species is much less common than *L. nobilis americanus* the specific characters could not be made out as accurately as for the other species. However, the nature of the septa is so strikingly different from that in other described species of *Lyttonia* that it is easily distinguished by this character alone. Since the original looped brachial ridge appears to have been obliterated by the strong development of the invagination of the outer shell layer, which has assumed a peculiar interrupted form, this species would seem to be extraordinarily specialized.

*Distribution:* Hess formation (Glass Mountains) at 17b, 207, and 222. Leonard formation (Glass Mountains) at 7. Word formation (Glass Mountains) at 54 and T140.

**Genus LEIORHYNCHUS Hall**

*Diagnosis:* Shell rhynchonelliform, medium-sized to rather large, subovate in outline and commonly subglobular in form. Median fold and sinus well developed, the plications obsolete or nearly obsolete on the lateral slopes of the valves, more or less well developed on the fold and sinus. In the pedicle valve the hinge-teeth are small and are supported by slender, vertical dental lamellae. In the brachial valve a well defined median septum is present in the rostral part of the valve; it is divided internally to form a V-shaped crural cavity. Hinge-plate divided, the inner margin of each lateral part being supported by one of the lateral walls of the crural cavity. Cardinal process absent. Crura formed by the anterior extension of the inner margins of the two divisions of the hinge-plate.

Genotype: *Orthis quadricostata* Vanuxem.

*Geologic range:* Devonian to Permian.

*Observations:* This genus has generally not been recognized above the Mississippian, but several Pennsylvanian and Permian species are almost identical in external form

with the Devonian and Mississippian ones. Since the sole distinction of the genus from *Pugnoides* is the outer configuration, there appears to be no reason for not referring them to *Leiorhynchus*. Moreover, the outer resemblance between the different species is so strong that they appear to be genetically related. In the lower Pennsylvanian are two species—*L. carboniferum* Girty and *L. rockymontanus* (Marcou). The genus does not occur in the Cisco or in the lower Permian of Trans-Pecos Texas, the next appearance being in the Leonard formation.

**LEIORHYNCHUS BISULCATUM (Shumard)**

Pl. XXXIII, Figs. 1–8

1858 *Camarophoria? bisulcata.* Shumard, Trans. Acad. Sci. St. Louis, vol. 1, p. 296 (date of volume, 1860).
Dark Permian limestone: Guadalupe Mountains; conglomerate at the mouth of Delaware Creek.

1859 *Camarophoria bisulcata.* Shumard, *idem,* p. 394, pl. 11, figs. 2a–d.
Dark and white limestone: Guadalupe Mountains; conglomerate at the mouth of Delaware Creek.

1908 *Pugnax? bisulcata.* Girty, The Guadalupian Fauna, U. S. Geol. Sur. Prof. Paper 58, pp. 310–312, pl. 21, figs. 11–12.
Dark limestone, Pine Spring Guadalupe Mountains (sta. 2930).

1908 *Pugnax? bisulcata* var. *gratiosa.* Girty, *idem.,* p. 312, pl. 21, figs. 10–10c.
Dark limestone, Pine Spring Guadalupe Mountains (sta. 2930).

1908 *Pugnax? bisulcata* var. *seminuloides.* Girty, *idem,* pp. 312–313, pl. 21, figs. 13–16, pl. 29, fig. 9?.
Middle Capitan, Capitan Peak (sta. 2926); dark limestone, Pine Spring, Guadalupe Mountains (sta. 2930); Delaware Mountain formation, southern Delaware Mountains (stas. 2965, 2969, and 3500).

*Description:* Shell medium-sized, outline subcircular to subpentagonal, width about equal to length. Pedicle valve shallow, brachial valve generally strongly and evenly convex. Ventral beak small and erect, showing clearly the foramen and the false cardinal area. The cardinal margins form slightly more than a right angle. Dorsal beak small, rounded, and in general hidden beneath the beak of the pedicle valve. In the median posterior part of the pedicle valve there is a narrow fold, which is replaced toward the front by a broad, shallow sinus, first distinguishable near the middle of the valve; the sinus may be perfectly smooth or may bear as many as 4 low, rounded ribs, the usual number being 3. Median posterior part of the brachial valve depressed in a narrow sinus, which gives way to the front in a fold, which is high in the anterior part of the shell; the fold is smooth or is marked by as many as 5 low plications corresponding to the sulci of the sinus. Lateral slopes of both valves smooth or bearing as many as 4 low, rounded, indistinct plications. Anterior margin strongly deflected downward on each side of the sinus, but, except for angulations caused by the plications, nearly straight between these deflections.

The brachial valve has internally a median septum supporting a V-shaped crural cavity and a divided hinge-plate. The pedicle valve contains short dental lamellae.

Dimensions in mm. of several specimens:

| | 1 (Las Del. sec. 2) | 2 (Las Del. sec. 2) | 3 (Shumard's type) |
|---|---|---|---|
| Length | 15 | 16 | 13.5 |
| Width | 17 | 18 | 15 |
| Thickness | 11 | 10 | 9 |

*Observations:* There can be no doubt as to the correctness of the identification with Shumard's species. The name is here used in the broad sense in which it was originally applied, the two varieties recognized by Girty being regarded as synonyms. The variety "*gratiosa*" was established for those shells having 2 or 3 broad plications on the fold and sinus, and "*seminuloides*" for those with unplicated fold and sinus. From one calcareous concretion in the black shales northwest of Las Delicias I collected 56 specimens of *L. bisulcatum,* of which 29 would be placed in the typical variety of Girty, with more than 3 plications on the fold, 16 would go into the variety "*gratiosa,*" and 11 in "*seminuloides.*" The gradation is so complete, however, that many of these specimens would be difficult to place. Since all three of Girty's varieties were collected principally from his sta. 2930 it appears that they, too, were but stages of variation in a highly variable species. At Girty's localities, the variety "*seminuloides*" was the more abundant, while among my own specimens the typical form is the common one.

The most closely related species is *Leiorhynchus weeksi* (Girty) and its variety *nobilis.* That species is larger, has broader plications, and the umbonal part of the brachial valve is not depressed, and of the pedicle valve not elevated as in *L. bisulcatus.* Another related species is *L. rockymontanus* (Marcou), which is larger, broader, and has a less well defined fold and sinus.

*Distribution:* Leonard formation (Glass Mountains) at 128. *Waagenoceras* zone (Las Delicias) at section 1, bed 19, section 2, beds 4 and 6 (abundant).

**LEIORHYNCHUS WEEKSI NOBILIS (Girty)**

Pl. XXXIII, Figs. 9–11

1910 *Pugnax weeksi* var. *nobilis.* Girty, Fauna of the Phosphate Beds of the Park City Formation, U. S. Geol. Sur. Bull. 436, pp. 32–33, pl. 3, figs. 5–7.
Phosphate beds of the Park City formation: Wyoming and Idaho.

*Description:* Shell medium-sized, subovate to subpentagonal, width and length about equal. Ventral beak large, pointed, suberect, flattened; dorsal beak inconspicuous, strongly incurved. The umbonal regions of both valves are evenly convex. The fold of the brachial valve and the sinus of the pedicle valve originate at about mid-length; they are broad and well defined. There are three broad low plications on the fold and two in the sinus, and three less distinct plications on each lateral slope. The plications originate somewhat posterior to mid-length of the shell, and may even extend nearly to the beaks.

Dimensions in mm. of several specimens:

| | 1 (loc. 144) | 2 (loc. 265) | 3 (loc. 144) | 4 (Girty's type) |
|---|---|---|---|---|
| Length | 18 | 18 | 19 | 26 |
| Width | 18 | 17 | 20 | 24 |
| Thickness | 11 | 11 | 11 | 13 |

*Observations:* The specimens of the present collection are more like the unusual specimen figured by Girty on plate 3, fig. 7 than they are like the type specimen of the variety. As Girty pointed out in his description, his specimens are so variable that it is difficult to find natural lines of division permitting the distinction of varieties. Although typical forms of the variety *nobilis* are quite different from the very faintly plicated typical forms of *L. weeksi*, there is complete intergradation between them. The Word specimens show less variation, all of them being definitely referable to the variety. The most important distinction between *L. weeksi* and its variety and *L. bisulcatus* is that the latter has a depressed dorsal umbonal region, showing the beginnings of a sinus which soon loses its identity in the median fold.

Shimer has described a species* *(Paraphorhynchus obscurum)* from Rocky Mountain quartzite, near Banff, which resembles typical *L. weeksi* in every respect except that it is said to be radially striate, and for that reason is placed in *Paraphorhynchus*. In view of the agreement in other particulars, it seems plausible to suggest that the two species are identical, the surface ornamentation of the Canadian specimens being in some way due to their manner of preservation.

*Distribution:* Word formation (Glass Mountains) at 144 (common), 259, and 265.

### Genus PUGNOIDES Weller

*Description:* Shells rhynchonelliform, below medium size, subovate in outline, with the fold and sinus well developed. Both valves marked by rounded or subangular plications which become obsolete in the posterior part of the shell. Internal characters of both valves as in *Leiorhynchus*.

Genotype: *Rhynchonella ottumwa* White.

*Geologic range:* Mississippian to Permian.

### PUGNOIDES ELEGANS (Girty)

Pl. XXXIII, Figs. 12–13; Pl. XXXIV, Figs. 2–4

1908 *Pugnax elegans.* Girty, The Guadalupian Fauna, U. S. Geol. Sur. Prof. Paper 58, pp. 315–316, pl. 15, figs. 13–14a.

Middle Capitan, Capitan Peak (sta. 2926); base of Capitan, hill southwest of Guadalupe Point (sta. 2906).

*Description:* Shell medium-sized, wider than long, with a very spreading triangular shape and a prominent fold and sinus. The plications are angular, and are 3 in number on the fold, 2 in the sinus, with 3 or 4 on each lateral slope. The median plication on the fold may be distinctly smaller than the adjacent ones. The lateral plications are rounder and fainter than those on the fold.

*In Upper Palaeozoic Faunas of the Lake Minnewanka Section, near Banff, Alberta, Geol. Sur. Canada, Bull. 42, 1926, pp. 46–47, pl. 7, figs. 1–3.

Dimensions in mm. of several specimens:

| | 1 (loc. 106) | 2 (loc. 108) | 3 (loc. 104) | 4 (loc. Ax) |
|---|---|---|---|---|
| Length | 13 | 11 | 13 | 18 |
| Width | 14 | 14 | 18 | 22 |
| Thickness | 9 | 8 | 9 | 14 |

*Observations:* This species is distinguished from *P. swalloviana* by its more spreading, triangular shape and the constant presence of three plications on the fold and two in the sinus. It is narrower and has fewer plications on the fold and in the sinus than *P. transversus*.

*Distribution:* Hess formation (Glass Mountains) at 104, 106, and 108. Leonard formation (Glass Mountains) at 15?, T15a, and Ax. Word formation (Glass Mountains) at 154. Hess formation (Sierra Diablo) at 475 and 479.

### PUGNOIDES PINGUIS (Girty)

Pl. XXXIV, Figs. 15–17

1908 *Pugnax pinguis.* Girty, The Guadalupian Fauna, U. S. Geol. Sur. Prof. Paper 58, p. 319, pl. 21, figs. 21–21c.

Dark limestone, Pine Spring (sta. 2930).

*Description:* Shell medium-sized, subtriangular or subpentagonal, with rounded angles except at the beak. Pedicle valve shallow, beak small and erect, median sinus broad and shallow, originating anterior to mid-length, prolonged forward and upward as a tongue, the end of which lies at a right angle to the profile of the rest of the valve. Brachial valve deep, flattened over the median posterior part, strongly curved near the margins. Fold low, not distinct, except at the front. Plications of variable shape, in some specimens being low and rounded, in others rather high and angular; they originate several mm. anterior to the beaks. Their number is 3 to 5 in the fold and 2 to 4 in the sinus, and 3 to 5 on each lateral slope, becoming smaller laterally. Within the pedicle valve there are strong short dental lamellae, and in the brachial valve a short median septum.

Dimensions in mm. of several specimens:

| | 1 (loc. 124) | 2 (loc. D3) | 3 (loc. 17c) | 4 (Girty's type) |
|---|---|---|---|---|
| Length | 13 | 13 | 15 | 13 |
| Width | 13 | 15 | 18 | 14.5 |
| Thickness | 10 | 11 | 12 | 9 |

*Observations:* This species is distinguished by the combination of flattened pedicle valve, geniculation of the tongue of the sinus, deep brachial valve, and few, strong plications. The shell is relatively narrower and has fewer plications in the fold and sinus than *P. shumardianus*. It is relatively narrower than *P. elegans*, has a larger number of plications, and does not have as deep a sinus or as high a fold.

*Distribution:* Hess formation (Glass Mountains) at 17b and 17c. Leonard formation (Glass Mountains) at 26, 120, 124, and D3. Word formation (Glass Mountains) at Tx.

**PUGNOIDES SHUMARDIANUS (Girty)**

Pl. XXXIV, Figs. 13–14

1908 *Pugnax shumardiana.* Girty, The Guadalupian Fauna, U. S. Geol. Sur. Prof. Paper 58, p. 316, pl. 5, figs. 15–17c.
Middle Capitan, Capitan Peak (sta. 2926).

*Description:* Shell medium-sized, triangular, wider than long. Pedicle valve mostly rather flat, but turned sharply upward near the front. Median sinus broad, shallow, and rather well defined, prolonged upward as a tongue. Beak erect. Sinus marked by 4 angular plications, the lateral slopes by three and a faint fourth one. Brachial valve convex, elevated in the median part as a fold, which merges into the general convexity posterior to mid-length of the shell. Umbonal region flattened. Fold marked by 5 plications, the lateral slopes by 3 or 4 lower plications. None of the plications extends to the beak, the umbones being smooth. The fold and sinus are less strongly marked as seen from above than from an anterior view. Within the pedicle valve there are two strong, short dental lamellae and in the brachial valve a low median septum.

Dimensions in mm. of several specimens:

| | 1 (loc. T15a) | 2 (loc. T15a) | 3 (loc. 241) |
|---|---|---|---|
| Length | 12 | 10 | 11 |
| Width | 16 | 12 | 14 |
| Thickness | 7 | 6 | 8 |

*Observations:* This species differs from *P. swallovianus* in having a larger size, less convexity, and 5 instead of 3 plications on the fold. It has about the same number of plications as *P. transversus*, but is much narrower in proportion to length.

*Distribution:* Leonard formation (Glass Mountains) at T15a.

**PUGNOIDES SWALLOVIANUS (Shumard)**

Pl. XXXIV, Figs. 10–12

1859 *Camerophoria Swalloviana.* Shumard, Trans. Acad. Sci. St. Louis, vol. 1, p. 294, pl. 11, figs. 1a–e (date of volume, 1860).
Permian rocks of the Guadalupe Mountains in Texas and New Mexico; one of the most abundant and characteristic species of the upper white limestone.

1908 *Pugnax swalloviana.* Girty, The Guadalupian Fauna, U. S. Geol. Surv. Prof. Paper 58, pp. 314–315, pl. 15, figs. 8–12d, pl. 21, figs. 17–19?.
Middle Capitan, Capitan Peak (sta. 2926); dark limestone, Pine Spring (sta. 2930).

1910 *Pugnax osagensis* var. *occidentalis.* Girty, Fauna of the Phosphate Beds of the Park City Formation, U. S. Geol. Sur. Bull. 436, pp. 33–34, pl. 1, figs. 21–22.
Phosphate beds of the Park City formation: Montpelier, Idaho

1927 *Pugnoides osagensis* var. *occidentalis.* Girty, in Geol., etc. of Part of Southeastern Idaho, U. S. Geol. Sur. Prof. Paper 152, p. 80, pl. 28, figs. 9–11.
Phosphoria formation: Montpelier, Idaho.

*Description:* Shell small, subpentagonal, strongly biconvex, as wide as or wider than long, with the greatest width at about mid-length of the shell. Pedicle valve very convex longitudinally, folded downward in the anterior half of the shell to form a deep, sharply defined sinus, which extends forward and upward as a tongue. Beak large, high, and erect. The sinus contains 2, or rarely 1, plications, and each lateral slope bears 3 or 4 plications. Brachial valve strongly convex, but with a distinct mesial depression in front of the beak. The median fold arises posterior to mid-length of the shell, and attains a considerable height at the front margin. The fold contains 3, or rarely 2, high, angular plications, and there are 3 or 4 plications on each side. The plications of both valves originate somewhat posterior to mid-length of the shell. Within the pedicle valve there are two short strong dental lamellae, and in the brachial valve a hinge-plate which is supported on each side by short lamellae, and gives rise to the downward-curving crura at its anterior end. There is a low dorsal median septum.

Dimensions in mm. of several specimens:

| | 1 (loc. 144) | 2 (loc. T240) | 3 (Shumard's type) | 4 (Phosphoria) |
|---|---|---|---|---|
| Length | 9 | 11 | 10 | 14 |
| Width | 11 | 11 | 10 | 16 |
| Thickness | 6.5 | 8 | 7 | 9 |

*Observations:* This species is distinguished from *P. osagensis* by its lower fold and generally greater number of plications on the lateral slopes. The difference is, however, small. In the Wewoka form *P. osagensis percostata* there are as many plications on the lateral slopes, but they are smaller than in *P. swallovianus.* In *P. texanus* the median fold is broader and lower, and has 3 or 4 rather than 2 or 3 plications. *P. bidentata* differs only in being smaller, more round in outline, and generally much more gibbous. The specimens from the Phosphoria described as *P. osagensis occidentalis* do not differ in any way from the Texas forms, except that the specimens figured are somewhat larger.

*Distribution:* Word formation (Glass Mountains) at 144, 148, 162, T240, 241, 243, and Tx.

**PUGNOIDES BIDENTATUS (Girty)**

Pl. XXXIV, Fig. 1

1908 *Pugnax bidentata.* Girty, The Guadalupian Fauna, U. S. Geol. Sur. Prof. Paper 58, pp. 318–319, pl. 21, figs. 20–20c, pl. 24, figs. 17–17c.
Dark limestone, Pine Spring (sta. 2930); basal black limestone, Guadalupe Point (sta. 2920). (Type from sta. 2920.)

*Description* (after Girty): "Shell small. Ventral valve rather strongly bent longitudinally and inflected at the sides. Sinus broad and deep, occupying most of the valve; defined by a rather sharp angulation, which forms, even where undefined, the line along which the shell is inflected. A slight groove lateral to the angulation adds to its prominence and is the only indication of lateral plication. The sinus contains a low median rib not visible except toward the front margin. Beak apparently rather large, high, and erect.

"Dorsal valve strongly convex. Fold high, broad, occupying most of the shell; defined at the sides by an angular groove, which is succeeded laterally by a low ridge, making one indistinct lateral plication on each side. The fold bears a rather strong median sulcus, so that its summit is divided into two plications. The plications of both valves are subangular and moderately strong, but disappear a slight distance from the margin."

The specimens in the present collection generally have 3 plications on the fold and 2 in the sinus, but the agreement with Girty's types in other particulars makes it very probable that the identification is correct. The number of plications is variable in most species of this genus.

Dimensions in mm. of several specimens:

| | 1 (loc. 491) | 2 (loc. 16) | 3 (loc. 120) | 4 (Girty's type) |
|---|---|---|---|---|
| Length | 8 | 8 | 10 | 8 |
| Width | 9 | 9 | 10 | 7 |
| Thickness | 7 | 7 | 5.5 | 5.5 |

*Observations:* This species is very close to *P. osagensis,* but has a relatively broader fold and sinus, in some specimens fewer plications on the fold and in the sinus, and generally a more gibbous form. It is smaller, narrower, and has fewer plications than *P. texanus. P. swallovianus* is larger, more subpentagonal in outline, and less strongly convex.

*Distribution:* Leonard formation (Glass Mountains) at 4, 12, 16, 24s, 25, 26s, 38, 120, 128, and D1. Bone Canyon member (Delaware Mountains) at 491.

### PUGNOIDES TEXANUS (Shumard)

Pl. XXXIV, Figs. 5–9

1859 *Rhynchonella Texana.* Shumard, Trans. Acad. Sci. St. Louis, vol. 1, p. 393 (date of volume, 1860).
Dark (Permian) limestone: Guadalupe Mountains; conglomerate at the mouth of Delaware Creek, Texas.

1908 *Pugnax osagensis.* Girty (non Swallow), The Guadalupian Fauna, U. S. Geol. Sur. Prof. Paper 58, pp. 317–318, pl. 24, figs. 16–16b.
Dark limestone, Pine Spring (sta. 2930?); Delaware Mountain formation, Guadalupe Point (sta. 2931); basal black limestone, Guadalupe Point (sta. 2920); Delaware Mountain formation, southern Delaware Mountains (sta. 2964).

1908 *Rhynchonella? texana.* Girty, *idem.,* p. 324.

*Description:* Shell small, transversely elliptical; lateral slopes converge toward the beak at an angle of 90° to 120°. Pedicle valve gently convex, marked in the anterior half by a strong median sinus, which is deflected upward as a tongue. Brachial valve more strongly convex, mesially elevated in front. Anterior part of the shell marked by high costae, three of which generally occur on the fold and two in the sinus, and from three to five on each lateral slope. Internally there are short dental lamellae in the pedicle valve; in the brachial valve there is a divided hinge-plate, which gives rise to the downward-curving crura.

Dimensions in mm. of several specimens:

| | 1 (loc. 440) | 2 (loc. 223) | 3 (loc. 439) | 4 (Shumard's type) |
|---|---|---|---|---|
| Length | 12 | 11 | 7 | 9 |
| Width | 14 | 12 | 8 | 7.5 |
| Thickness | 10 | 6 | 5 | 5.5 |

*Observations:* This species is very variable. In the lot from loc. 223 the sinus is rather shallow, the ventral umbo well arched, and the tongue curved up rather than geniculate. In other shells, typified by those from loc. 440a, the sinus is more abruptly deepened at the front, the ventral umbo smaller, and the front of the tongue at right angles to the plane of the lateral margins. The two types seem, however, to intergrade, and there does not seem enough basis for distinction.

*P. texanus* is very close to *P. osagensis* (Swallow). It is distinguished by being generally larger, having relatively somewhat broader plications, and having a shorter tongue-like extension of the sinus of the pedicle valve. In *P. osagensis,* moreover, the lateral plication next to the sinus tends to rise high at the front, while in *P. texanus* it is rather low, so that the profile of the shell does not show a sharp angulation in the ventral anterior part as does the Pennsylvanian species.

Girty believed that Shumard's *Rhynchonella texana* must actually belong to *"Rhynchonella"* rather than *Pugnax* because of the presence in it of a "false area," which Girty says is a character quite foreign to the group of *Pugnax.* Weller, however, mentions such a feature in some Mississippian *Pugnoides.* Shumard's description agrees very completely with the present form except that he states that the posterior margins converge at an angle of 63°. If 63° is taken to be the complement of the angle between the two lateral slopes, or 117°, there is then no disagreement, and it is probable that this was Shumard's meaning. The range of the forms identified by Girty as *Pugnax osagensis* into the dark limestone of the Delaware Mountain formation suggests that other forms than that here identified as *P. texanus* were included under that designation. It is very likely that the Delaware Mountain specimens so identified were varieties of *P. swalloviana.*

The small species *P. osagensis pusilla* Girty, of the Yeso formation of New Mexico, may be the same as *P. texanus,* since it is abundant at the same horizon, but Girty's figures show it as oval in outline, while specimens of *P. texanus* of the same size are triangular.

*Distribution:* Hess formation (Glass Mountains) at 105, 208, and 223 (abundant). Leonard formation (Glass Mountains) at 235. Lower Gym formation (Hueco Mountains) at 379. Middle Gym formation (Hueco Mountains) at 414, 420, 424, and 430. Upper Gym formation (Hueco Mountains) at 437, 438, 439, 440a, and 440b. Gym formation (Hueco Mountains) at 443 and 449. Bone Canyon member (Sierra Diablo) at 491?.

### PUGNOIDES TRANSVERSUS King, n. sp.*

Pl. XXXIII, Figs. 14–17

*Description:* Shell rather large, transversely ovoid, much wider than long. Pedicle valve slightly convex, the median

*Recently described as *P. mesacostalus* Girty, Jour. Wash. Acad. Sci. Vol. 19, 1929, p. 413. (Editor's.)

third depressed in a strongly defined sinus, which originates near mid-length of the shell and extends forward some distance as a tongue of the front margin. Brachial valve evenly convex, with a sharply defined median fold, originating somewhat posterior to mid-length. Sinus marked by from 4 to 6, and the fold by from 5 to 7 angular plications. The median plication on the fold is smaller than the other 4, so that the fold has the appearance of possessing a median sinus. Each lateral pair of plications on the fold appears to have arisen from the bifurcation of a single one. Lateral plications 6 to 8 in number on each side. The plications originate about 3 mm. anterior to the beaks, and increase in number by intercalation on the lateral slopes. In the pedicle valve the dental lamellae are long and slightly diverging. In the brachial valve there is a short median septum, from the anterior end of which there arises a thickening of the inside of the shell; crura formed by the anterior extension of the inner margins of the two divisions of the hinge-plate.

Dimensions in mm. of several specimens:

| | 1 (loc. 222) | 2 (loc. 222) | 3 (loc. 222) | 4 (loc. T107) | 5 (loc. 107) |
|---|---|---|---|---|---|
| Length | 15 | 16 | 17 | 16 | 16 |
| Width | 24 | 13 | 27 | 22 | 22 |
| Thickness | 10 | 12 | 14 | 11 | 11.5 |

*Observations:* This species is distinguished by its large size, its great width, and the presence of 4 to 6 plications on the fold and in the sinus.

*Distribution:* Hess formation (Glass Mountains) at T15, 107, T107, 212, and 222.

### Genus RHYNCHOPORA W. King

*Diagnosis:* Shell rhynchonelliform, generally pentagonal ovoid in outline, the median fold and sinus well developed in the anterior half of the shell. Surface of both valves marked by simple, subangular or rounded plications which may be longitudinally grooved toward the front. Internally the dental lamellae are well developed in the pedicle valve. Brachial valve with a strong median septum in the rostral region, which is divided internally to form the walls of a crural cavity; this is covered over on its cardinal side by the undivided hinge-plate, which continues forward beyond the anterior termination of the crural cavity. Crura formed by the anterior extension of the part of the hinge-plate opposite the lateral walls of the crural cavity. Shell structure punctate.

Genotype: *Terebratula geinitziana* Verneuil.

*Geologic range:* Mississippian to Permian.

*Observations:* Weller, from whom the above diagnosis was adapted, believes that the undivided hinge-plate and complete closure of the crural cavity except anteriorly are the most important characters of the genus. He expressed uncertainty, however, as to whether these characters were shown in the genotype, a fossil of the European Permian. Likharew* has since shown that *R. geinitzianus* is internally like *R. hamburgensis* and other Mississippian species in these characters. Weller's diagnosis of *Rhynchopora* should therefore be regarded as expressing the typical characters of the genus. As noted below, *R. illinoisenses* may have a divided hinge-plate; if so, it is non-typical.

*Mém. Soc. russe Min., ser. 2, vol. 54, pp. 115–118, pl. 1.

### RHYNCHOPORA ILLINOISENSIS (Worthen)

Pl. XXXIV, Fig. 18

1884 *Rhynchonella illinoisensis.* Worthen, Illinois State Mus. Nat. Hist. Bull., no. 2, p. 24.

1914 *Rhynchopora nikitini.* Kozlowski (non Tschernyschew?), Brachiopodes du Carb. Supér de Bolivie, Ann. de Pal., vol. 9, pp. 84–86, text fig. 22, pl. 9, figs. 67–70.

1915 *Rhynchopora illinoisensis.* Girty, Fauna of the Wewoka Formation, U. S. Geol. Sur. Bull. 544, p. 85, pl. 10, figs. 12–12c. See for synonymy.

*Observations:* There can be little doubt that this species is the same as that described from Bolivia by Kozlowski as *R. nikitini* Tschernyschew. If the identification with the European species is likewise correct, then Tschernyschew's name would be a synonym of Worthen's, the latter having been proposed a year before the former. Kozlowski's specimens are described as having a divided hinge-plate, a statement contrary to that in the generic diagnosis given above. Specimens of *R. illinoisensis* from the Gaptank formation likewise seem to have this character. If this feature is characteristic of the species, then *R. illinoisensis* is unlike not only *R. taylori* of the higher beds of the Permian, but also the Mississippian forms described by Weller.

*Distribution:* Wolfcamp formation, *Uddenites* zone (Glass Mountains) at 88s and T88. Wolfcamp formation (Glass Mountains) at 90.

### RHYNCHOPORA TAYLORI Girty

Pl. XXXIV, Figs. 19–24

1908 *Rhynchopora taylori.* Girty, Fauna of Phosphate Beds of the Park City Formation, U. S. Geol. Sur. Bull. 436, p. 34, pl. 3, fig. 8.
Phosphate beds of the Park City formation, Montpelier, Idaho.

*Description:* Shell medium-sized to rather large, subpentagonal, wider than long, with the greatest width near mid-length. Convexity moderate, the brachial valve having only a very slightly convex profile along the median line, then turning sharply downward to meet the tongue-like median extension of the pedicle valve without any angulation at the front suture, the anterior part of the shell being flat. Sinus of the pedicle valve equal in width to about half that of the shell, marked by 6, or less commonly 7 or 8, regular rounded plications. Fold of the brachial valve broad, not very high, well-defined near the front. Each lateral slope is ornamented by about 8 to 11 plications, of which the outer ones are less distinct. Within the pedicle valve there are long, diverging dental lamellae. In the brachial valve a strong median septum is divided posteriorly to form the walls of the crural cavity; the latter is covered over on the cardinal side by an undivided hinge-plate. The shell texture is punctate.

Dimensions in mm. of several specimens:

| | (1 (loc. 144) | 2 (loc. T240) | 3 (Girty's type) |
|---|---|---|---|
| Length | 11 | 12 | 12 |
| Width | 14 | 14 | 14.5 |
| Thickness | 9 | — | 8.5 |

*Observations:* This species is distinguished from the very similar *R. illinoisensis* by having a proportionately narrower fold, which has 1 or 2 less plications, while the lateral slopes are proportionately broader and have 1 to 3 more plications, the total number being about the same in both species. In extreme forms the limits of variation overlap.

*Distribution:* Word formation (Glass Mountains) at 6, 46, 54, 144, T240, 243, and Tc. Upper Gym formation (Hueco Mountains) at 436. Limestone of Cerro Caballo (Las Delicias) at section 1, beds 11 and 14.

### Genus CAMAROPHORIA W. King

*Description:* Shells small to large, rhynchonelliform, rostrate, subovate to subpentagonal in outline, with well developed median sinus in the pedicle valve and fold in the brachial valve. Surface plicated or with the plications nearly or quite obsolete. In the pedicle valve the dental lamellae are continued forward into a well developed spondylium which either is supported by a median septum or rests directly upon the floor of the valve for a part or the whole of its length. In the rostral part of the brachial valve the hinge-plate is continuous and is supported by a strong median septum; between the hinge-plate and the floor of the valve the median septum also supports a concave cruralium which is continued far beyond the hinge-plate, becoming broader and more highly elevated toward the front; posteriorly the median septum passes through the cruralium for the support of the hinge-plate, but the extension of the septum above the concave surface of the cruralium soon disappears beyond the anterior margin of the hinge-plate.

Genotype: *Terebratula Schlotheimi* von Buch.

*Geologic range:* Devonian to Permian.

### CAMAROPHORIA DELOI King, n. sp.*

Pl. XXXIV, Figs. 24–27

*Description:* Shell small, transversely elliptical, strongly biconvex. Pedicle valve about equal in depth to the brachial valve, depressed in the median anterior part of the shell to form an ill-defined sinus, which extends forward and upward as a tongue at the anterior margin. Brachial valve evenly convex, elevated in the median anterior part as a low fold. The fold and sinus are marked by 4 or 5 angular plications, which originate 2 to 4 mm. anterior to the beak. Each lateral slope bears 4 or 5 plications. New plications are introduced on the lateral slopes and in one specimen at the sides of the fold and sinus. Within the pedicle valve the dental lamellae curve around to join, forming a spondylium, which is supported by a median septum.

Dimensions in mm. of several specimens:

| | 1 (loc. 376) | 2 (loc. 241) | 3 (southern Arizona) |
|---|---|---|---|
| Length | 9 | 11 | 10 |
| Width | 11 | 14 | 12 |
| Thickness | 7.5 | 8 | 8 |

*Observations:* This species is distinguished by its *Pugnoides*-like exterior, with more numerous plications than are the rule in *Camerophoria*. The shell is also smaller and more strongly convex than other species of *Camerophoria* in this fauna.

*Distribution:* Lower Gym formation (Hueco Mountains) at 376 (type) and 379. Word formation (Glass Mountains) at 241. Lower Permian, Sawmill Canyon, near Onyx Mine, Santa Rita Mts.; Snyders Hill, 18 miles southwest of Tucson, Arizona.

### CAMAROPHORIA THEVENENI Kozlowski

Pl. XXXIV, Fig. 28

1914 *Camarophoria Thevenini.* Kozlowski, Brachiopodes du Carb. Supér. de Bolivie, Ann. de Pal., vol. 9, pp. 86–88, text fig. 23, pl. 9, figs. 71–76; pl. 10, fig. 15.

*Description:* Shell small, wider than long, subtriangular. Ventral beak conical, incurved, with a triangular delthyrial foramen, in which lies the cardinal process of the other valve. Median part of the pedicle valve depressed, and extended forward as a tongue. There are 2 or 3 low plications in the sinus, which originate some distance anterior to the beak. Lateral slopes marked by one or two fainter plications. Brachial valve more strongly convex than the other, with a low median fold, which bears 3 or 4 plications. Lateral slopes marked by 2 or 3 low plications which arise near the anterior margin.

Within the pedicle valve the dental lamellae curve downward toward each other to form a spondylium, which is supported by a median septum.

Demensions in mm. of one specimen: length, 12; width, 13; thickness, 7.

*Observations:* This species, like *C. huecoensis*, has somewhat the outer form of a *Pugnoides*. It is distinguished from other species of the genus in the present fauna by its fewer and fainter plications. It is much smaller than *C. venusta.*

*Distribution:* Wolfcamp formation (Glass Mountains) at T82?, 93, and 195. Leonard formation (Glass Mountains) at 123. Lower Gym formation (Hueco Mountains) at 400.

### CAMAROPHORIA VENUSTA Girty

Pl. XXXIV, Figs. 29–31; Pl. XXXV, Figs. 1–5

1908 *Camarophoria venusta.* Girty, The Guadalupian Fauna, U. S. Geol. Sur. Prof. Paper 58, pp. 303–304, pl. 31, figs. 6–6c.

Middle Capitan, Capitan Peak (sta. 2926); Delaware Mountain formation, "Comanche Canyon," Glass Mountains.

*Description:* Shell large, subtriangular, generally wider than long. Pedicle valve shallow, with an elongate, slightly incurved, pointed beak. Delthyrium broadly triangular, nearly filled by the beak of the other valve. One specimen

*Editor's Note: This species is in part embraced in *C. huecoensis* Girty, Jour. Wash. Acad. Sci., Vol. 19, 1929, p. 412.

from loc. 174 shows small deltidial plates between the incurved beak of the brachial valve and the rounded foramen that occupies the apex of the valve. Sinus more than a third the width of the shell, deep and well-defined, arising about 8 mm. from the beak and expanding to the front margin. There are 3 to 7, but generally about 5, sharp plications in the sinus and 4 to 6 more rounded ones on each side. In some specimens some of the plications bifurcate toward the front. Brachial valve strongly convex, with a broad, well-defined fold. The sinus bears 4 to 8 plications, and each lateral slope 4 to 6.

Within the pedicle valve there are small hinge-teeth on each side of the broad delthyrium. The sides of the delthyrium are supported by plates which curve toward each other downward to the median line of the inner wall of the shell, where they join in an even curve to form a spondylium. The median septum supporting the spondylium is very greatly reduced and mainly exists as a thickening of the shell along the median line only. In the brachial valve the hinge-plate is prolonged forward into a concave elevated cruralium, which is supported by a median septum. On each side of the cruralium is attached a thin crus, which extends forward about half the length of the cruralium, and curves downward. The cruralium extends farther forward than the spondylium of the other valve, and terminates in three rounded lobes.

Dimensions in mm. of several specimens:

| | 1 (loc. 174) | 2 (loc. 107) | 3 (loc. 484) | 4 (Girty's type) |
|---|---|---|---|---|
| Length | 27 | 23 | 18 | 18 |
| Width | 34 | 26 | 27 | 19 |
| Thickness | 21 | 16 | 15 | 11 |

*Observations:* Girty's type specimen is not typical of the species as here described, being proportionately narrower and having a greater number of plications on its lateral slopes.* Some of the forms in the present collection are very close to the type, however, and grade into typical forms which are broad and coarsely plicate. Since the species is very variable, and since the type came from the Glass Mountains, it appears highly unlikely that the specimens of the present collection are not conspecific.

*Camarophoria venusta* is similar to many of the Eurasian forms from the Upper Carboniferous and Permian, but as it is a variable species, and the Eurasian Camerophorias have been rather minutely split, comparison is difficult. Probably the most similar species is *C. purdoni* Davidson. The extremely low ventral median septum may serve to distinguish *C. venusta* from some of the other species.

*Distribution:* Wolfcamp formation (Glass Mountains) at 76, 94s?, and 193?. Hess formation (Glass Mountains) at 1, 4, 106, 107, T107, 207, 222, and Tv. Leonard formation (Glass Mountains) at 4, 8–14, T9, 12, 14, 15, T15a, 16, 22, 26, 26s, 30–31, 82, 104, 105, 120, 124, 127, 128, 174, 226, 227, T233, Ax, Tb, D1, and D4. Word Formation (Glass Mountains) at 44–46, T141, 144, Tx148, 150, 155, 162, T240, 246, and 247. Bone Canyon member (Guadalupe Mountains) at 221. Bone Canyon member (Sierra Diablo) at 484. Leonard formation (Baylor Mountains) at 502. Delaware Mountain formation (Sierra Diablo) at 504. Leonard formation (Shafter) at 515. Altuda member (Glass Mountains) at 171.

*Editor's Note: The broader shells included above have been described by Girty as *C. inaequalis*. Jour. Wash. Acad. Sci., vol. 19, 1919, p. 411.

### CAMAROPHORIA? INDENTATA (Shumard)

Pl. XXXIV, Fig. 32

1859 *Rhynchonella indentata.* Shumard, Trans. Acad. Sci. St. Louis, vol 1, p. 393 (date of volume, 1860).
White (Permian) limestone: Guadalupe Mountains.

1908 *Rhynchonella? indentata.* Girty, The Guadalupian Fauna, U. S. Geol. Sur. Prof. Paper 58, pp. 321–322, pl. 15, figs. 20–20c.
Middle Capitan, Capitan Peak (sta. 2926); dark limestone, Pine Spring (sta. 2930).

*Description:* Three specimens are assigned to this species. The one from the limestone of Cerro Caballo has the following characters:

Shell medium-sized, subpentagonal, longer than wide, with the greatest width in the anterior part. Convexity low, but the shell is considerably crushed. Pedicle valve depressed in the median part into a very poorly defined sinus, which originates at about mid-length. It is prolonged somewhat forward as a tongue, but does not curve upward as sharply as in most other species of the genus. Brachial valve evenly and slightly arched, rising into a depressed fold only near the front. Surface marked by simple angular plications, which appear to originate a short distance anterior to the umbos. The median fold and sinus bear 3 or 4 plications, and each lateral slope about 7. There are also well marked concentric varices of growth. Within the pedicle valve the dental lamellae form a spondylium, which is supported by a low median septum, and in the brachial valve there is a high median septum, which presumably supports a cruralium. Length, 19 mm.; width, 17 mm.; thickness, 9 mm.

It may be seen that the description does not accord very well in some respects with that of Shumard. The low convexity of the shell is due to its having been considerably flattened by crushing. Except for the thickness, the dimensions of Shumard's type are somewhat less than those of this specimen, the length being 14 mm., the width 12, and the thickness 10. In view of the general agreement in number and strength of plications, relative size of median fold and sinus, etc., it seems best to identify the Mexican specimen questionably with Shumard's species.

The specimen from the Altuda member accords much better with Shumard's description, being much more strongly convex.

*Observations:* Girty described "short dental plates" in this species. This would at first seem to exclude it from belonging to *Camerophoria*, but since the spondylium lies near the floor of the valve, the dental plates may appear in many specimens to be quite distinct because the part near the floor of the valve, where they curve inward toward each other, is not well preserved. If *R.? indentata* is not a *Camerophoria* it would be difficult to know in what group to place it. This species resembles *Uncinuloides guadalupensis* in that the plications are small and extend nearly to the beaks, but it does not have the peculiar configuration of the front margin which characterizes *Uncinuloides*. Girty states that *R.? indentata* differs from his

species *R.? longaeva* in being larger and has a false area and flattened dorsal umbo. No specimens referable to the latter species were collected, so that I am unable to make comparison with my own specimens.

*Distribution:* Limestone of Cerro Caballo (Las Delicias) at bed 14, section 1. Word formation (Glass Mountains) at 265. Altuda member (Glass Mountains) at 57.

#### Genus UNCINULOIDES King, novum

*Diagnosis:* Shell rostrate, subtriangular in outline, with a well developed median sinus in the pedicle valve and fold in the brachial valve, anterior margin broad and flattened, lying nearly perpendicular to the longitudinal axis of the shell. Surface marked by fine, flattened costae, longitudinally grooved on the truncated anterior part of each valve. In the pedicle valve the dental plates converge toward the floor of the valve, and curve together to form a spondylium, which may rest on the inner wall of the valve or may be slightly elevated and supported by a low median septum. Hinge plate and cruralium of the brachial valve similar to those of *Camerophoria.*

Genotype: *Rhynchonella guadalupae* Shumard.

*Geologic age:* Permian.

*Observations:* This genus has in every respect the external form of *Uncinulus,* but the internal characters of *Camerophoria.* The fact that the spondylium rests on the floor of the pedicle valve or is supported by only a very low median septum is a difference from many species of *Camerophoria,* but in *Camerophoria venusta* the structure is quite comparable. A large number of species in the Permian of Eurasia have been identified as *Uncinulus.* In no instance have the internal characters of any of these genera been described in enough detail to permit certain identification with either *Uncinulus* or *Uncinuloides.* All that is stated as to the internal characters of these species is that there are diverging dental lamellae in the pedicle valve and a median septum in the brachial valve. It might at first be thought that the divergence of the dental lamellae would exclude these species from a genus in which a spondylium is present, yet an examination of the specimens from the Glass Mountains shows that the spondylium generally lies on the floor of the valve, and when the surface is cut or broken away, the dental lamellae appear to join the inner wall of the shell at some distance from each other, the true nature of the structure being apparent only when seen from the inside in an etched specimen. The Permian species assigned by various authors to *Uncinulus* are as follows:*

*U. theobaldi* Waagen, Virgal group of the Salt Range (said by Diener to be a synonym of *U. jabiensis);* Artinsk of Russia; lower Permian, *Schwagerina* horizon, Darwas, Bokhara; Permian of Ussuriland(?).

*U. jabiensis* Waagen, Chideru group of the Salt Range; Upper Permian of the Himalayas at Chitichun I and Malla Sangcha; Permian of the Residency of Djambi, Sumatra; *Waagenoceras* horizon of Basleo, Timor; *Lyttonia* horizon of the Kitakami Mountains, Japan; Upper Permian near Djulfa, Armenia (?); lower Permian, *Schwagerina* horizon, Darwas, Bokhara.

*U. posterus* Waagen, Chideru group of the Salt Range; Jarkalo, China.

*U. timorensis* (Beyrich), *Waagenoceras* horizon of Basleo, Timor; Permian of Residency of Djambi, Sumatra; Upper Permian of the Himalayas at Malla Sangcha; Lantsankiang Valley of China.

*U. hanieli* Broili, *Waagenoceras* horizon of Basleo, Timor.

*U. wichmanni* (Rothpletz), Timor.

*U. wangenheimi* (Pander), *Schwagerina* limestone of the Urals.

*U. velifer* Gemmellaro, Sosio limestone of Sicily; Trogkofel formation of the Karnic Alps.

*U. amor* Gemmellaro, Sosio limestone of Sicily.

*U. siculus* Gemmellaro, Sosio limestone of Sicily.

*U. arthaberi* Tschernyschew, lower Permian, *Schwagerina* horizon, Darwas, Bokhara; Upper Permian of Djulfa, Armenia.

I think it quite likely that all these are actually *Uncinuloides,* for the time separating the latest occurrence of undoubted *Uncinulus* from the first occurrence of these Permian species is a vast one, extending from the Lower Devonian to the Permian.

#### UNCINULOIDES GUADALUPENSIS (Shumard)

Pl. XXXV, Figs. 6–7

1858 *Rhynchonella Guadalupae.* Shumard, Trans. Acad. Sci. St. Louis, vol. 1, p. 295 (date of volume, 1859).
White (Permian) limestone: Guadalupe Mountains.

1859 *Rhynchonella Guadalupae.* Shumard, *idem.,* p. 392, pl. 11, figs. 6a–c.
White (Permian) limestone: Guadalupe Mountains.

1908 *Rhynchonella? guadalupae.* Girty, The Guadalupian Fauna, U. S. Geol. Sur. Prof. Paper 58, pp. 323–324, pl. 16, figs. 10–10b.
Top of Capitan, Capitan Peak (sta. 2966?).

*Description:* Shell medium-sized, subtriangular, with rounded basal angles, width and length about equal. Lateral margins nearly straight, diverging at about a right angle; sides flattened, with an angulation at the junction of the valves. Anterior marginal region truncated, there being no angulation where the two shells meet. Pedicle valve not very strongly convex, beak high and somewhat incurved over a distinctly defined cardinal area, which includes a triangular deltidial plate on each side of an oval delthyrium. Sinus of the pedicle valve broad and shallow, originating somewhat posterior to mid-length of the shell and curving upward through an angle of about 90° to form a tongue in the truncated anterior face of the shell. Brachial valve regularly convex, umbo flattened, curvature gentle from the beak nearly to the front margin, where the shell is deflected strongly downward to meet the tongue of the pedicle valve; median fold low and broad, well-defined only near the front. Surface marked by many fine rounded costae, which extend to the beaks, where, however, they are very small and faint. Toward the anterior part of the shell the costae become broad and flattened, and are separated by narrower striae. On the flattened anterior region of the shell the costae are medially grooved, so that they appear double. There are 9 or 10

*Many of these species are so narrowly limited that there has been much discussion as to their validity.

costae in the fold and on the sinus of a mature shell, and 15 to 19 on each side.

Within the pedicle valve the hinge-teeth are supported by dental lamellae which are produced forward more than a third the length of the valve to form a spondylium, which rests upon the floor of the valve, not being elevated on a median septum. In the brachial valve the hinge-plate is broad, and produced forward as a trough-like cruralium, which extends more than half the length of the shell and is supported in the posterior part by a high septum. From each side of the hinge-plate there arise thin downward-curving crura.

Dimensions in mm. of several specimens:

| | 1 (loc. T128) | 2 (loc. 120) | 3 (loc. D1) | 4 (R. guadalupae Sh.) |
|---|---|---|---|---|
| Length | 17 | 16 | 17+ | 14.5 |
| Width | 22 | 20 | 36 | 19 |
| Thickness | 13 | 12 | — | 12 |

*Observations:* A specimen identified by Girty was said to have a perforated hinge-plate—a character that could not be verified or disproved in the specimens of the present collection. *Rhynchonella? indentata* and *R.? longaeva* of Girty somewhat resemble *U. guadalupensis,* but seem to be considerably smaller and more coarsely plicate, and do not have the characteristic shape of *Uncinuloides.* Hardly any important difference may be seen between *U. guadalupensis* and *U. jabiensis* Waagen, the common Asiatic species. Probably the latter has somewhat stronger costae.

*Distribution:* Upper Wolfcamp formation (Glass Mountains) at 196. Upper Hess formation (Glass Mountains) at 207. Leonard formation (Glass Mountains) at 3, 7, 16, 24, 30–31, 31, 37, 83, 194, 120, 128, 231, Tb, and D1. Word formation (Glass Mountains) at 51, 54, and Tx148.

## Family SPIRIFERIDAE

### Genus SPIRIFER Sowerby

*Diagnosis* (after Weller): Shells small to very large, generally wider than long, rarely longer than wide, hinge-line straight, narrower than the greatest width of the shell and the cardinal extremities rounded, or more commonly the greatest width of the shell is along the hinge-line and the cardinal extremities are angular and more or less extended, in some conspicuously acuminate. Median sinus in the pedicle valve and fold in the brachial valve generally well developed, more rarely absent. Surface of both valves marked by radiating plications which may be simple without division from the point of origin at the cardinal margin to the anterior margin, or may divide in various ways; the plications may be present upon the lateral slopes only or upon both the lateral slopes and the fold and sinus. Besides the plications the surface may also be marked by fine radiating lirae or by fine or coarse concentric growth lines, or by both radiating and concentric markings. The pedicle valve has the beak variously elevated above the hinge-line and variously incurved, the cardinal area varying from very narrow to high, generally arched but in some nearly or quite flat, the delthyrium rather broadly triangular and open; the surface of the cardinal area is transversely striate and the inner shell layers bear a series of vertical canals at whose ends along the hinge-line the shell tissue may be produced in a row of denticles which articulate with a row of pits in the opposite valve. Internally the hinge-teeth are strong and are supported by short or undeveloped dental lamellae; the muscular area is of moderate size and may be deeply impressed, ovate or obcordate in outline, occupied in large part by the diductor scars, which are generally marked by radiating or branching furrows. The brachial valve has a very narrow cardinal area divided by a broadly triangular delthyrium; the cardinal process is a low, transverse, sessile apophysis with its surface vertically striated; the muscular impressions much less strongly marked than in the pedicle valve; the dental sockets narrow and of moderate depth, the socket plates well developed and at their ends suporting the crural bases. Crura long, straight, and slightly divergent, spiralia directed obliquely outward and backward toward the cardinal extremities; primary lamellae not united by a jugum, the position of the jugum being indicated by the presence of a pair of spine-like processes upon the primary lamellae a little in front of their junction with the crura.

Genotype: *Anomites striatus* Martin.

*Geologic range:* Silurian to Permian.

*Observations:* This genus has, like *Productus,* recently been divided into a large number of genera* but as a large number of these are of doubtful value, they are not recognized as distinct genera in the present paper.

### SPIRIFER CONDOR d'Orbigny

Pl. XXXV, Figs. 8–10; Pl. XXXVI, Fig. 1

1842 *Spirifer condor.* D'Orbigny, Voy. dans l'Amér. mér. t. 3, pt. 4, pl. 5, figs. 11–14.

1869 *Spirifer striatus* var. *multistriatus.* Toula, Sitzb. d. Kais. Acad. d. Wiss., Wien, vol. 59, p. 3, pl. 1, figs. 2–4.

1902 *Spirifer condor.* Tschernyschew, Mem. Com. Geol. Russie, vol. 16, p. 141, pp. 531–532, pl. 12, figs. 1–2, pl. 38, figs 1–2.

1914 *Spirifer condor.* Kozlowski, Brachiopodes du Carb. Supér. de Bolivie, Ann. de Pal., vol. 9, pp. 67–70, pl. 1, fig. 1; pl. 7, figs. 10–14, text fig. 16.

1914 *Spirifer condor.* H. L. F. Meyer, Carbonfaunen aus Bolivia und Perú, Neues Jahrbuch für Mineralogie, etc., vol. 37, pp. 604–610, text fig. 1.

1919 *Spirifer condor.* Douglas, Geological Sections through the Andes, Quart. Jour. Geol. Soc., vol. 70, p. 32, pl. 9, figs. 1a–c.

1929 *Spirifer condor.* Steinmann, Geologie von Perú, p. 47, fig. 42.

*Description* (modified from Kozlowski): Shell large, subtriangular, very much wider than long, the width being greater in proportion to length in younger individuals. The width may exceed twice the length. Pedicle valve somewhat more convex than the other, median part regularly arched, with a median sinus which originates at the beak and is prolonged forward at the anterior margin as a tongue. The sinus is narrow and well defined by bordering costae which are somewhat larger than those in the

*Fredericks, George: Table pour définition des genres de la Fam. *Spiriferidae* King, Bull. de l'Ac. Sci. URSS, 1926, nos. 4–5, pp. 393–423.

sinus. Beak broad and somewhat incurved over the cardinal area, which is about 10 times as wide as high and has a gently concave transverse profile. Surface of the area marked by horizontal and vertical striae, the latter being more abundant and distinct. Delthyrium triangular, somewhat wider than high, covered by deltidial plates, which are divided from one another by a suture along the median line; the anterior end of the plates is concave toward the ventral beak, the dorsal beak having fit into this depression, leaving the delthyrium open.

Brachial valve less convex and shorter than the other, with a low beak not curving over the cardinal line. Dorsal area very low. A median dorsal sinus corresponds to the fold of the other valve.

Surface of both valves covered by rounded radial costae, about 1 mm. wide, separated by grooves of about the same width, originating at the apices, where, on each valve, there are only about 4 costae on each side of a median one. Between the beak and the anterior margin each of the original costae bifurcates from 3 to 5 times, depending on the size of the specimen; the newly added costae very shortly attain the same size as the others, so that there is an average of 50 subequal costae, lying somewhat diagonal to the margin. The costae in the fold originate from the original median costa by continuous bifurcation, while those in the sinus, on the other hand, originate from the costae at the sides, rather than in the center of the sinus. Fine concentric imbrications are also present, which are distantly spaced over most of the surface but close together in the marginal region.

Within the pedicle valve there is, on each side of the delthyrium, a strong obtuse hinge tooth, below which is a thickening of the shell which is prolonged backward as a thickening of the delthyrial border until it merges with the apical callosity, which is very strongly developed, filling the entire apical region beneath the deltidial plates. Adductor muscular impressions long and narrow, extending from the apical callosity about three-fourths the length of the shell; they are separated medially by a low ridge.

Dimensions in mm. of several specimens:

| | 1 (loc. Tmm) | 2 (loc. 375) | 3 (loc. 88) | 4 (Kozlowski) |
|---|---|---|---|---|
| Length | 28 | 31 | 27 | 43 |
| Width | 56 | 63 | 52 | 80 |
| Thickness | 24 | — | — | 34 |

*Observations:* The Texas specimens agree perfectly with those described by Meyer and Kozlowski from Bolivia. Those figured by Tschernyschew from the *Cora* and *Schwagerina* beds of the Uralian show no characters by which they might be distinguished from the typical ones. The *S. cameratus* Derby from Bolivia is tentatively regarded as not synonymous, for it shows some differences and probably comes from a much lower horizon. The species described by Shumard as *S. organensis*, collected in the Organ Mountains near Ft. Filmore, New Mexico, is very probably a synonym of this species; it may have been collected from the lower Gym horizon.

This species is placed by Fredericks in the genus *Anelasma* of Ivanov,* which is distinguished from *Spirifer* by lacking true dental lamellae, having only ridges bordering the inside of the delthyrium. *S. condor* certainly fits into this genus by definition, but as there is no certainty that the species placed in *Anelasma* form a natural group, it is thought best not to recognize it.

*Distribution:* Wolfcamp formation *Uddenites* zone (Glass Mountains) at 88 and T88. Upper Wolfcamp formation (Glass Mountains) at 85, T93, 194, and 196. Lower Gym formation (Hueco Mountains) at 375, 376, 377a, 377b, 381, 387, 401, and Tmm.

### SPIRIFER MARCOUI INFRAPLICA King, n. var.

Pl. XXXVI, Figs. 2–4

*Description:* Shell large, much wider than long, with the greatest width at the hinge-line. Pedicle valve strongly convex longitudinally but of low transverse curvature. Cardinal area high and flat except at the apex, which is incurved; in large shells the sides of the area may be subparallel to the hinge-line for most of the width of the shell, then, near the lateral margins, sloping abruptly to the hinge. Beak broad, short, not greatly bent over. Sinus deep and rounded, originating near the beak. Brachial valve strongly convex on its lateral slopes, but less strongly on the median fold. Beak somewhat elevated; cardinal area rather high. Median fold high and rounded, prominent only near the front. On one specimen (from loc. Tq) the crest of the fold is not situated exactly along the median line, but somewhat to one side.

Entire surface marked by subequal rounded radial costae, of which 4 to 5 occur in the space of 5 mm. Fasciculation does not take place. From 15 to 20 costae occur on the fold and sinus and 30 to 40 on each lateral slope. Fine concentric striae give the surface a cancellated appearance.

Within the pedicle valve there are strong diverging dental lamellae.

Dimensions in mm. of several specimens:

| | 1 (loc. 499) | 2 (loc. Tq) | 3 (loc. 126) | 4 (Waagen's) | (*S. marcoui*) |
|---|---|---|---|---|---|
| Length | 38 | 45 | 32 | 58 | 53 |
| Width | 77 | 70 | 54 | 92 | 82 |
| Thickness | 27 | — | 23 | 45 | 38 |

*Observations:* No perfect specimens of this variety were found. It is very close to the widespread Eurasian species *S. marcoui*, which was thought by Waagen to be a representative of the form identified by Marcou as *S. striatus* from Pecos Village, New Mexico. Since Marcou's collection was from a much lower horizon geologically, its identity is to be regarded as doubtful. *S. marcoui infraplica* is distinguished from *S. marcoui* by having a narrower fold and sinus, the former of which is lower and the latter shallower, and the sides of the sinus are more sharply defined from the lateral slopes. A very similar species is *S. logani* from the Mississippian, which has a broader

*Ivanov: Sur la systématique et la biologie du genre *Spirifer* et de quelque brachiopodes de C-ii et C-iii du Gouv. du Moscou, Bull. de Sect. Géol. de la Soc. des Amies des N., Anthr. et G. de M., 1925, p. 109.

median fold and sinus and a lower cardinal area. Some specimens of the two might be difficult to distinguish.

*Distribution:* Leonard formation (Glass Mountains) at T9, 122, Tb, and D16. Bone Canyon member (Sierra Diablo) at 485. Victorio Peak member (Sierra Diablo) at 496 and 499. Hess formation (Finlay Mountains) at 511. Leonard formation, transition beds (Shafter) at Tq (no. 15984).

**Subgenus NEOSPIRIFER Fredericks***

*Diagnosis:* Shell similar in configuration to *Spirifer* but with the costae arranged in fascicles, generally of three. The fascicles may be distinct only in the posterior part of the shell, becoming obsolete toward the front. As in *Spirifer* there are short dental lamellae, and the muscle scars of the pedicle valve are divided by a median ridge.

Type: *Spirifer fasciger* Keyserling.

*Geologic range:* Pennsylvanian and Permian.

*Observations:* Although Fredericks selected as his type for this group a widespread Eurasian species (known in southeastern Asia generally as *Spirifer musakheylensis* Davidson), it is more particularly an American type. Whereas in Europe the middle part of the Pennsylvanian is characterized by *Choristites mosquensis*, in North America by far the most abundant spiriferid in rocks of the same age is *Neospirifer triplicatus*. In the Texas Permian fauna *Neospirifer* is the predominant type, while in Eurasia it is no more important than many other groups of spirifers.

**SPIRIFER (NEOSPIRIFER) BAKERI King n. sp.**

Pl. XXXVII, Figs. 4–6

*Description:* Shell large, subpentagonal, wider than long, with the greatest width at the hinge-line; cardinal angles acute or even very sharp, the shell being greatly extended at the hinge-line in young stages. Young specimens are relatively short in proportion to length, the increase in length taking place after the greatest width has been attained. Pedicle valve evenly convex longitudinally, the curvature decreasing gradually from the back to the anterior margin. Beak broad, short, pointed, not greatly incurved. Cardinal area low and broad, concave. Median sinus broad, evenly rounded, and rather deep, defined on each side by a high, angular ridge. From the ridges bordering the sinus there is an even slope on each side toward the ears. Brachial valve equal in convexity to the other, with a low median fold, which rises abruptly from the otherwise evenly convex shell, and has a rounded crest. Within the pedicle valve there are short dental lamellae which support strong hinge-teeth.

Surface marked by low, rounded costae, which in the posterior part increase by bifurcation, and are grouped in prominent bundles of three, in which the median costa is larger than the two lateral ones. Toward the front they do not divide, and become broad and indistinct. The costae in the fold and sinus are smaller than those on the lateral slopes and not at all bundled; toward the front the fold and sinus may be almost smooth.

Dimensions in mm. of several specimens:

| | 1 (loc. 46) | 2 (loc. 53) | 3 (loc. 7) | 4 (loc. 46) |
|---|---|---|---|---|
| Length | 37 | 29 | 20 | 22 |
| Width | 64 | 43 | 34 | 54 |
| Thickness | 22 | — | — | 17 |

*Observations:* This species is distinguished from *S. triplicatus*, *S. pseudocameratus*, and similar species by the sharp definition of the median sinus by lateral ridges and the low plications, which become indistinct toward the front. The young, relatively short specimens of this species are very close to *S. niger* Waagen from the Amb group of the Salt Range.

*Distribution:* Leonard formation (Glass Mountains) at 7. Word formation (Glass Mountains) at 46 (common, types), and 58.

**SPIRIFER (NEOSPIRIFER) COSTELLA King, n. sp.**

Pl. XXXVII, Figs. 1–3

*Description:* Shell rather large, alate in youth but approximately semicircular at maturity; hinge-line equal to the greatest breadth. Pedicle valve gently and evenly convex both transversely and longitudinally. Cardinal area high, triangular, and strongly concave. Delthyrium a broad triangle. Beak high, sharply pointed and incurved. Median sinus narrow and shallow, originating at the very apex of the beak. Dental lamellae short but strong. Brachial valve of low convexity, with a low apex and low cardinal area. Median fold low and rounded.

Surface marked by fine radial costae, originating at the apices as about 8 costae, each of which shortly splits into three parts, and toward the front margin splits several more times. About 7 costae occur in the space of 5 mm. All the costae originating from one original rib are grouped together in a raised fascicle, which is most pronounced in the median part of the shell and becomes lower and less distinct anteriorly. Generally about 7 or 8 costae compose each fascicle. The fascicles are not very strongly marked, and in some specimens are hardly noticeable. The surface is ornamented likewise by prominent lamellose erect striae of growth, which are very closely arranged and cover both valves. There are also irregular, strong imbricating marks of growth paralleling the anterior and lateral margins.

Dimensions in mm. of several specimens:

| | 1 (loc. 174) | 2 (loc. 174) | 3 (loc. 126) |
|---|---|---|---|
| Length | 35 | 27 | 20 |
| Width | 50 | 45 | 35 |
| Thickness | 21 | — | — |

*Observations:* This species is most closely related to *S. fasciger* Keyserling (=*S. musakheylensis* Davidson). The size of the costae, type of bundling, lamellose striation, and general outline are quite similar. The European species, however, has a higher fold and deeper sinus, a relatively much lower cardinal area, and a more triangular outline.

*Distribution:* Leonard formation (Glass Mountains) at 24, 123, 126, and 174 (types). It is not common at any locality.

*Fredericks, George: Études paléontologiques. 2, Les Spiriferides du carbonifére supérieur de l'Oural, Bull. du Com. Géol., vol. 38, no. 2, 1919, p. 311.

———: Table pour définition des genres de la Fam. *Spiriferidae* King, Bull. de l'Ac. Sci. URSS, 1926, no. 4–5, p. 406.

### SPIRIFER (NEOSPIRIFER) MEXICANUS LATUS King, n. var.

Pl. XXXVII, Fig. 7; Pl. XXXVIII, Fig. 1

*Description:* Shell transversely elliptical, with a very short hinge-line, equaling less than half the width of the shell; greatest width at about mid-length. Pedicle valve strongly convex, being somewhat deeper than the other valve. Beak high, pointed, and incurved over the narrow, concave cardinal area, which is only about twice as wide as high and is not well defined from the rest of the shell. Sinus shallow but well-defined, originating at the apex. Brachial valve evenly convex, with a low but distinct median fold. Both valves are covered by moderately coarse radial costae, which to some extent, especially on the brachial valve, are arranged in bundles of about 3 costae each. Of the internal characters, only the strong dental lamellae of the pedicle valve were distinguished.

Dimensions in mm. of several specimens:

| | 1 (loc. 123) | 2 (loc. 128) | 3 (loc. 491) |
|---|---|---|---|
| Length | 26 | 19 | 28 |
| Width | 32 | 22 | 34 |
| Thickness | 17 | — | — |

*Observations:* This variety is very easily distinguished by its elliptical outline and the very high, narrow, poorly defined cardinal area. The only difference from the typical form of the species is that this variety is relatively much broader. A closely related species is *S. oldhamianus* Waagen, in which the cardinal area is very similar, but the shell is broader and has a higher fold and deeper sinus.

*Distribution:* Leonard formation (Glass Mountains) at 5, 12, 26s, 120, 123 (type), 126, 128, Tg?, T233, and D1. Bone Canyon member (Sierra Diablo) at 491. Hess formation (Glass Mountains) at 8 and 104.

### SPIRIFER (NEOSPIRIFER) PSEUDOCAMERATUS Girty

Pl. XXXVIII, Figs. 4–6; Pl. XXXIX, Figs. 1–3

1859 *Spirifer cameratus.* Shumard, Trans. Acad. Sci. St. Louis, vol. 1, p. 391 (date of volume, 1860).
(Permian) sandstone and white limestone: Guadalupe Mountains.

1908 *Spirifer* sp. b. Girty, The Guadalupian Fauna, U. S. Geol. Sur. Prof. Paper 58, pp. 363–364.
Delaware Mountain formation, Guadalupe Point (sta. 2919); Delaware Mountain formation, southern Delaware Mountains (stas. 2967, 3501); Delaware Mountain formation, "Comanche Canyon," Glass Mountains.

1920 *Spirifer pseudocameratus.* Girty, in Ore Deposits of Utah, U. S. Geol. Sur. Prof. Paper 111, pp. 644–645, pl. 56, figs. 10–15.
Permian: Utah.

*Description:* Shell large, wider than long, but not alate. Greatest width at the hinge-line, the cardinal angles being acute in wide shells and more nearly rectangular in shells of greater proportionate length; the angle varies likewise according to the size of the shell.

Pedicle valve convex, especially in the umbonal region; beak small and strongly incurved. Cardinal area of moderate height, lying oblique to the plane of the shell margin. Sinus relatively narrow, of moderate depth, fairly well defined, occupied by one or two small costae. The ribs bordering the sinus bifurcate in such a way that costae extend from them forward along the flanks and even into the bottom of the sinus, attaining the same size as the original costae of the sinus, so that the latter lose their identity. Brachial valve moderately convex. Fold low and narrow but well-defined, divided by a median groove, each side bears costae which have arisen by bifurcation of the ribs bordering the median groove.

On the lateral slopes of both valves there are strongly fasciculate ribs, composed of groups of 3 or 4 costae, which may be subequal or with one costae larger than the others. There are from 3 to 5 fascicles on each side of the beak, and 4 or 5 other costae are developed distal to these toward the cardinal angles.

Within the pedicle valve the delthyrium is closed in its apical half by deltidial plates, underneath which the shell is thickened as an apical callosity. Hinge-teeth strong, their lower sides extending into the dental lamellae, which connect with the floor of the valve on each side of the deeply impressed diductor muscle scars.

Dimensions in mm. of several specimens:

| | 1 (loc. 121) | 2 (loc. 156) | 3 (loc. 174) | 4 (loc. 493) |
|---|---|---|---|---|
| Length | 42 | 52 | 36 | 40 |
| Width | 65 | 80 | 52 | 62 |
| Thickness | 28 | 32 | 22 | 27 |

*Observations:* The only difference between the Texas specimens and the types of the species, from Utah, is that the latter have a rather strongly bilobate fold on the brachial valve, the median groove being somewhat deeper than in those of the present collection. This species is distinguished from *S. cameratus* and *S. triplicatus* by being generally somewhat larger, with commonly a greater relative length, coarser, more irregular costae, and a narrower and lower fold and sinus. Girty regards the specimens from Nevada identified by Meek as *S. cameratus* as probably *S. pseudocameratus.*

*Distribution:* Leonard formation (Glass Mountains) at 8?, T9, 15?, T15?, 26, 26s, 37, T119, 120, 121 (abundant), 124, 125?, 128, T128?, 129, 174 (abundant), Tb, Tba?, D3, D5, D7, and D12. Word formation (Glass Mountains) at 6, 44–46, 47, 54, 58, 138?, T140, 144, T144, T144a, 146, 148, T148, Tx148, 150, 155, 192, 239, T239?, 241, 246, 250, 255, 256 (abundant), 257, Tu, and Tx?. Bone Canyon member (Sierra Diablo) at 484. Victorio Peak member (Sierra Diablo) at 492, 493, and 496. Victorio Peak member (Brackett Draw) at Taa. Leonard formation (Baylor Mountains) at 503. Delaware Mountain formation (Sierra Diablo) at 504. Leonard formation (Shafter) at Tj (thin-bedded zone) and 518. Limestone of Cerro Caballo (Las Delicias) at section 1, beds 11, 12 (¼ mile S of 21), 14, and 19.

### SPIRIFER (NEOSPIRIFER) HUECOENSIS King, n. sp.

Pl. XXXVIII, Figs. 2–3

*Description:* Shell large, transversely elliptical, hinge-line somewhat shorter than the greatest width, which is at

about mid-length; sides evenly rounded. Pedicle valve of low convexity. Beak low, bluntly pointed, very strongly incurved over the cardinal area, which is narrow, very sharply defined, and nearly flat in its anterior and lateral parts but very strongly incurved under the apex. Sinus shallow, originating at the beak, very broad near the anterior margin, where there is a spatulate forward prolongation. Brachial valve bearing a low median fold, which is indistinct in the posterior part but is prominently elevated near the anterior margin. Surface of both valves marked by low, rounded radiating costae, 3 or 4 of which occur in the space of 5 mm. Near the apex they are somewhat bundled, but over most of the surface they are subequal and not at all fasciculate. In the anterior part of the shell they become low, and the surface may be nearly smooth. There are also closely spaced lamellar growth lines and strong concentric varices of growth in the anterior region.

Within the pedicle valve (of a specimen from loc. 107) there is a strongly developed apical callosity, and strong hinge-teeth, which are not supported by distinct dental lamellae. The muscle scars are deeply impressed and not divided by a median ridge.

Dimensions in mm. of several specimens:

| | 1 (loc. 389) | 2 (loc. 389) | 3 (loc. 107) | 4 (loc. 107) |
|---|---|---|---|---|
| Length | 41 | 37 | 43 | 36 |
| Width | 58 | 59 | 60? | 68 |
| Thickness | 25 | 20 | 25 | 30 |

The specimen from loc. 107, no. 4, is exceptionally broad.

*Observations:* This species is distinguished from *S. mexicanus* Shumard and its varieties by its larger size, greater relative width, and low, sharply defined cardinal area.

*Distribution:* Hess formation (Glass Mountains) at 104? and 107 (common). Lower Gym formation (Hueco Mountains) at 389 (common, types).

### SPIRIFER (NEOSPIRIFER) TRIPLICATUS Hall

Pl. XXXIX, Fig. 4

1852 *Spirifer triplicata.* Hall, Stansbury's Exploration and Survey Great Salt Lake of Utah, p. 410, pl. 4, figs. 5a–e.

1914 *Spirifer cameratus.* Kozlowski, Brachiopodes du Carb. Supér. de Bolivie, Ann. Pal., vol. 9, p. 70, pl. 5, figs. 6–11.

1915 *Spirifer cameratus.* Girty, Fauna of the Wewoka Formation, U. S. Geol. Sur. Bull. 544, pp. 87–91, pl. 11, figs. 4–4b. See for synonymy.

1920 *Spirifer triplicatus.* Girty, in Ore Deposits of Utah, U. S. Geol. Sur. Prof. Paper 111, pp. 645–646, pl. 54, fig. 22.

*Observations:* According to Girty the upper Pennsylvanian fossils generally designated as *S. cameratus* should be called *S. triplicatus.* This species has stronger fasiculation of the costae, higher and more angular fold and sinus, a more triangular outline, and a relatively wider hinge-line than *S. cameratus* which occurs in the Pottsville and equivalent strata.

*Distribution:* Wolfcamp formation, *Uddenites* zone (Glass Mountains) at 88, T88, 88s, 94s, 175, 175a, and 201. Wolfcamp formation (Glass Mountains) at 75, 77, 87, 89, 91a, 93, 93s, 193, 196, and 198.

### SPIRIFER (NEOSPIRIFER) TEXANUS Meek

Pl. XXXIX, Fig. 5

1871 *Spirifer (Trigonotreta?) texanus.* Meek, Proc. Acad. Nat. Sci. Phil., vol. 23, pp. 179–181.

1921 *Spirifer texanus.* Plummer and Moore, Penn. Formations of N. Central Texas, U. of Texas Bull. 2132, p. 148, pl. 21.

*Observations:* This species is distinguished from *S. triplicatus* by its greater length in proportion to width, its short hinge-line, and its high, incurved ventral beak. The shell seems to be thicker than specimens of *S. triplicatus* of the same size.

*Distribution:* Wolfcamp formation, *Uddenites* zone (Glass Mountains) at 88, 88s, and T88. Wolfcamp formation (Glass Mountains) at 91a and 92.

### Subgenus ELIVINA Fredericks*

*Diagnosis:* Shells medium-sized to large, biconvex, the hinge-line narrower than the greatest width, the cardinal angles rounded, mesial sinus in the pedicle valve and fold in the brachial valve well developed. The pedicle valve with a fairly high cardinal area; internally the dental lamellae are well-developed and extend forward for some distance, inclosing the muscle scars, which are situated on a raised platform. There is no median ridge or septum dividing the muscle scars. Surface of both valves covered with low, broad, rounded costae, more or less arranged in fascicles. Shell impunctate.

Type: *Spirifer tibetanus* Diener.

*Geologic range:* Permian.

*Observations:* Other representatives of this subgenus are *S. rajah* Salter and *S. sulcifer* Shumard. *Elivina* was established for plicatocostae (fasciculate) shells having well-developed dental lamellae but no "euseptoid" or "euseptum." I propose to restrict the name to shells having those internal characters and the external configuration similar to that of the genotype. As the genotype was the only species assigned to this subgenus by Fredericks, this does not decrease the contents of the group as originally constituted, but gives it greater value because of the recognition of a greater number of distinctive features. *Spirifer lyra* Kutorga is similar to species of *Elivina* in all but incomplete dental lamellae, or, as Fredericks calls this structure, "delthyrial ridges." Fredericks makes *Spirifer lyra* the type of his subgenus *Eliva.* If, as seems likely, the distinguishing characters are of less importance than supposed by Fredericks, the species of the *Spirifer lyra* and *S. tibetanus* groups should be assigned to the same subgenus.

It may be said that *Elivina* is externally like *Spiriferella,* but is impunctate. The genotype of *Spiriferella, S. saranae,*

*Fredericks, G.: Études paléontologiques. 2, Les Spiriferides du carbonifére supérieur de l'oural. Bull. du Com. Geol., vol. 38, no. 2, 1919, pp. 314–315.

———: Table pour définition des genres de la Fam. *Spiriferidae* King, Bull. de l'Ac. Sci. URSS, 1926, nos. 3–4, p. 409.

differs further in having the dental lamellae coalesce over a ventral median septum ("euseptum" of Fredericks), but Weller* includes in *Spiriferella* shells having discrete, diverging dental plates. In that respect the Mississippian *Spiriferella* species resemble *Elivina*, but they still have a punctate shell.

**SPIRIFER (ELIVINA) SULCIFER Shumard**

Pl. XXXIX, Fig. 6; Pl. XL, Figs. 1–5

1858 *Spirifer sulcifera.* Shumard, Trans. Acad. Sci. St. Louis, vol. 1, p. 293 (date of volume, 1860).
White (Permian) limestone: Guadalupe Mountains.

1859 *Spirifer sulciferus.* Shumard, *idem.*, p. 391, pl. 11, figs. 3a–c.
White (Permian) limestone: Guadalupe Mountains.

1908 *Spirifer sulcifer.* Girty, The Guadalupian Fauna, U. S. Geol. Sur. Prof. Paper 58, p. 363, pl. 13, figs. 10–10b.

1908 *Spirifer* sp. a. Girty, The Guadalupian Fauna, U. S. Geol. Sur. Prof. Paper 58, pp. 362–363, pl. 21, fig. 2.
Dark limestone, Pine Spring (sta. 2930).

*Description:* Shell large, subpentagonal, length somewhat greater than the width, greatest width near or somewhat in front of mid-length, cardinal angles obtuse. Pedicle valve strongly convex, the curvature being greatest in the umbonal region and decreasing gradually forward. Cardinal area high and narrow, in large shells not perfectly triangular, being truncated at the sides, delthyrium broad triangular, closed by strongly developed deltidial plates. Median sinus broad and rather deep, originating near the apex and increasing in size toward the front. Brachial valve of much lower convexity than the other, a little wider than long, with a very low, broad median fold.

Surface of the pedicle valve marked by broad rounded costae, the largest of which border the median sinus. The costae are grouped in 3 or 4 fascicles on each side of the sinus; each fascicle consists of 3 or 4 costae, which generally do not subdivide anterior to the umbonal region. Small costae lie at the sides of the median sinus. The brachial valve bears on its low median fold generally 4 costae, which are not fasciculate, and several fascicles of costae on each side. Concentric striae of growth are likewise present on both valves. Shell impunctate.

Internally there are, in the pedicle valve, strong diverging dental lamellae, which are mostly enveloped in the apical callosity. The muscle scars lie between them on a platform, which is considerably elevated at the front. On each side of the muscular impressions the inside of the shell is covered with coarse granules.

Dimensions in mm. of several specimens:

| | 1 (loc. 246) | 2 (loc. 246) | 3 (loc. 146) | 4 (Shumard's type) |
|---|---|---|---|---|
| Length | 50 | 28 | 37 | 16.5 |
| Width | 47 | 24 | 32 | 14 |
| Thickness | 22+ | 15 | 18 | 10 |

*Observations:* Shumard's type of this species was a very small specimen, but it is indistinguishable from immature individuals of this large species. Girty suggested comparison of his *Spirifer* sp. a with *S. sulcifer*, which he knew only from Shumard's description. Study of the complete material from the Glass Mountains shows that Girty's specimen was a very typical brachial valve of *S. sulcifer*.

The external form of this species is nearly identical with that of many species of *Spiriferella*, but since *S. sulcifer* is impunctate, it cannot belong to that genus. Its closest relation is with *S. tibetanus* and *S. rajah*, which Diener has clearly shown not to be *Spiriferella*. From the former it differs in having a lower and broader cardinal area, and from the latter in having a lower dorsal median fold and a greater number of plications on the fold. The difference from *S. rajah* is, however, so small (see, for example, plate 119 of Wanner's Permische Brachiopoden von Timor) that it seems likely that *S. rajah* and *S. sulcifer* were representatives in the two continents of one varying form. This conclusion is borne out by the distribution of the two: *S. rajah* occurs in the high Permian shales in the Himalayas and in the *Waagenoceras* horizon of Timor, while *S. sulcifer* occurs abundantly in and above the *Waagenoceras* horizon of the Glass Mountains. *S. tibetanus* seems to have been an earlier form, for it is nowhere associated with *S. rajah*, occurring in the *Schwagerina* limestone of the Urals and the Trogkofel formation of the Karnic Alps.

*Distribution:* Middle and Upper Word formation (Glass Mountains) at T140, 144, T144, 146, 148, T148, Tx148, 150, 154, 158, T240a, 241, 243, T243, 246 (common), 247 (abundant), 248, 252, 255, 265, and Tc. One of the most characteristic fossils of the upper Word limestone.

**Genus SQUAMULARIA Gemmellaro**

*Diagnosis:* Shells medium-sized to large, subovate, subcircular or subovate in outline, with the hinge-line narrower than the greatest width of the shell, rounded cardinal extremities, and moderately developed or obsolete fold and sinus. Cardinal area of the pedicle valve rather small and arched, with poorly defined margins, the surface generally rounding with but little or no interruption into the lateral slopes of the valve; delthyrium rather large and open. Internally there are no dental or septal plates. Brachial valve less convex than the other, with a narrow cardinal area; internally the brachidium is similar to that of *Spirifer*. Surface of both valves covered by regular, concentric bands bearing fine prostrate spines, the tubes of which do not penetrate the full thickness of the shell.

Genotype: *Squamularia rotundata* Gemmellaro.

*Geologic range:* Pennsylvanian and Permian.

*Observations:* Species of this genus differ from those of *Reticularia* in the absence of dental or septal plates. *Reticularia* does not occur above the Mississippian.

**SQUAMULARIA GUADALUPENSIS (Shumard)**

Pl. XL, Figs. 6–8

1859 *Spirifer Guadalupensis.* Shumard, Trans. Acad. Sci. St. Louis, vol. 1, p. 391 (date of volume, 1860).
White (Permian) limestone: Guadalupe Mountains.

1908 *Squamularia guadalupensis.* Girty, The Guadalupian Fauna, U. S. Geol. Sur. Prof. Paper 58, pp. 367–369, pl. 14, figs. 4–11a.

*Mississippian Brachiopoda, Mon. 1, Ill. Geol. Sur., p. 412. 1914.

Middle Capitan, Capitan Peak (sta. 2926); basal Capitan, hill southwest of Guadalupe Point (sta. 2906); Delaware Mountain formation, Guadalupe Point (sta. 2919); Delaware Mountain formation, southern Delaware Mountains (sta. 2962); Delaware Mountain formation, "Comanche Canyon," Glass Mountains.

?1909 *Squamularia perplexa.* Girty, Fauna of the Manzano Group, U. S. Geol. Sur. Bull. 389, pp. 66–68.

Yeso formation, Mesa del Yeso; San Andres formation, San Andres.

*Description:* Shell large, ovoid, with the length about equal to or somewhat greater or less than the width, strongly biconvex; hinge-line equal to a little more than half the greatest width, which is near mid-length. Pedicle valve high, with a long, conical, incurved beak. Cardinal area triangular, three times as wide as high, poorly defined. There is no median sinus, but a slight emargination is present at the front. Brachial valve lower than the other, evenly convex, with an incurved beak which projects somewhat beyond the cardinal margin. Outer shell layer thin, generally not preserved; it is marked by narrow, regular, concentric sublamellose bands, which are marked by slender, closely crowded spine bases which originate at the posterior margin of the band and extend forward, giving rise to spines at the anterior margin of the band. If the surface is exfoliated, as is generally the case, the shell has a reticulate appearance, there being fine radial lirae at uneven intervals over the surface, which cross the concentric bands.

Dimensions in mm. of several specimens:

| | 1 (loc. 1) | 2 (loc. 106) | 3 (loc. D3) | 4 (loc. 246) | 5 (Girty) |
|---|---|---|---|---|---|
| Length | 25 | 31 | 16 | 23 | 45 |
| Width | 29 | 35 | 17 | 26 | 38 |
| Thickness | 15 | 20 | 12 | 17 | — |

*Observations:* This species is the longest ranged brachiopod in the Glass Mountains faunas. No important difference could, however, be seen between specimens from the Hess and those from the Capitan, and, while the Wolfcamp specimens are incomplete, they show very well that the surface is identical with that of the Capitan specimens. Most of the specimens in the present collection, however, seem to be rather more transverse than those described by Girty. Girty distinguishes two varieties of the species. The Glass Mountains specimens seem to be referable to the typical form.

*S. guadalupensis* is distinguished from *S. perplexa* by having very fine spines rather than large double-barrelled spines. The specimens from New Mexico identified by Girty as *S. perplexa* are, to judge from their geologic horizon and large size, probably *S. guadalupensis.* Girty stated that a separation from *S. guadalupensis* would be difficult. *S. guadalupensis* is very close to *S. indica* (Waagen), but is probably not as elongate, and it shows other minor differences in general form.

*Distribution:* Wolfcamp formation (Glass Mountains) at 93, 196, and 198. Hess formation (Glass Mountains) at 1, 24, 106 (common), T205, and 208. Leonard formation (Glass Mountains) at 7, 8, 8–14, T9, 12, 15, 26s, 82, 83, 104 (common), T119, 120 (common), T120, 121, 124, 128, 174, 231, 232, Tb, Tba, D1, D3, D12, and D22. Word formation (Glass Mountains) at 46, 54, 132, 141, 170, 246, 247, T251, 256, and 263?. Altuda member (Glass Mountains) at 43 and 57 (abundant). Bone Canyon member (Sierra Diablo) at 485. Victorio Peak member (Sierra Diablo) at 493. Leonard formation (Shafter) at 518 (common). Word? formation (Shafter) at Tp.

#### SQUAMULARIA PERPLEXA (McChesney)

1860 *Spirifer perplexa.* McChesney, Desc. New Spec. Foss., p. 43 (date of imprint, 1859).

1914 *Reticularia lineata* Mart., var. *perplexa.* Kozlowski, Brachiopodes du Carb. Supér. de Bolivie, Ann. de Pal., vol. 9, pp. 73–76, text fig. 18, pl. 1, figs. 3–4; pl. 10, figs. 19–27.

1915 *Squamularia perplexa.* Girty, Fauna of the Wewoka Formation, U. S. Geol. Sur. Bull. 544, pp. 92–94, pl. 11, figs. 1–3a. (See for synonymy.)

*Observations:* A few imperfect specimens are assigned to this species. They differ from the common *Squamularia guadalupensis* of higher horizons in their smaller average size and especially in having large double-barrelled spines; a feature excellently illustrated by Kozlowski.

*Distribution:* Wolfcamp formation, *Uddenites* zone (Glass Mountains) at 88s, 95, 201, and 203. Wolfcamp formation (Glass Mountains) at 90?.

### Genus AMBOCOELIA Hall

*Diagnosis* (After Weller): Shells small, plano-convex or slightly concavo-convex, subcircular to subsemielliptical in outline, the hinge-line generally a little narrower than the greatest width of the shell, the cardinal extremities rounded. Surface of valves smooth or with fine concentric lines of growth, rarely minutely spinose, shell substance impunctuate. Pedicle valve strongly convex with very prominent umbonal region, the beak strongly incurved, marked by a narrow, median, groove-like sinus which is generally strongest posteriorly; cardinal area well defined, concave, the delthyrium rather large and open, with incomplete deltidial plates. Internally the hinge-teeth are prominent, with recurved tips, not supported by dental lamellae; the muscular area small, the interior surface about the muscle scars strongly pitted. Brachial valve convex at the beak, becoming flat or concave forward, with a rather broad cardinal area lying in nearly a right angle to the area of the other valve; cardinal process bifurcating posteriorly, elongate, resting on the bottom of the valve except at the posterior end, the spires of the brachidium loosely coiled with few volutions, attached by long crura which are continuous with the primary lamellae; the jugum in the same rudimentary condition as in *Spirifer.*

Genotype: *Orthis umbonata* Conrad.

*Geologic range:* Silurian to Permian.

#### AMBOCOELIA GUADALUPENSIS Girty

Pl. XLI, Figs. 1–5

1908 *Ambocoelia planiconvexa* var. *guadalupensis.* Girty, The Guadalupian Fauna, U. S. Geol. Sur. Prof. Paper 58, pp. 370–371, pl. 14, figs. 12–14a.

Middle Capitan, Capitan Peak (sta. 2926) and peak north of Pine Spring (sta. 2902); Delaware

Mountain formation, southern Delaware Mountains (sta. 2936).

1910 *Ambocoelia arcuata.* Girty, Fauna of the Phosphate Beds of the Park City Formation, U. S. Geol. Sur. Bull. 436, p. 35, pl. 6, figs. 15–17.

Phosphate beds of the Park City formation: Sublette Range, Wyoming; Thomas Fork, Wyoming; Montpelier, Idaho.

*Description:* Shell large for the genus, hinge-line narrower than the greatest width, which is near mid-length. Pedicle valve highly inflated, especially in the umbonal region, with a long projecting, incurved beak. Cardinal area poorly defined, high, triangular, generally lying at an angle of about 135° to the plane of the margin, about a third occupied by the delthyrium, which is higher than wide. There is no trace of a median sinus. Brachial valve transverse, gently convex, somewhat flattened at the cardinal angles, hinge-line straight. Beak small and depressed. Surface marked only by fine concentric lines. Within the brachial valve are rather long prominent crural plates.

Dimensions in mm. of several specimens:

| | 1 (loc. 491) | 2 (loc. 265) | 3 (loc. T120) | 4 (*A. arcuata*) | 5 (*A. guadalupensis*) |
|---|---|---|---|---|---|
| Length | 17 | 14 | 13 | 15 | 11.5 |
| Width | 16 | 15 | 12 | 13 | 12.5 |
| Thickness | 10 | 8 | 7 | — | — |

*Observations:* This species averages much larger than *A. planoconvexa,* is relatively longer, has a more convex brachial valve, and lacks a median sinus. Girty mentions as differences also that *A. guadalupensis* has a lower cardinal area and longer, stronger, and more closely proximate crural plates. The specimens in the present collection agree with Girty's figures and description, but in them the cardinal area is fully as high relative to the size of the shell as in *A. planoconvexa.* In describing *A. arcuata,* Girty mentioned as a difference from *A. planoconvexa* the broader delthyrium of the former. His specimen figured on plate 6, fig. 17 shows a broad opening below the apex, but this seems to be broader than the delthyrium, which is shown in fig. 16 and is quite like that of the Texas Permian species, and certainly not conspicuously larger in proportion than that of *A. planoconvexa.* The characters mentioned in the first sentence, and which were emphasized by Girty in his discussion of *A. arcuata,* easily serve to distinguish this species. *A. arcuata* was not compared with *A. planoconvexa guadalupensis* in the original description, but the two descriptions agree so well, and the specimens in the present collection would so easily fit in either one of the descriptions that there can hardly be any doubt as to their identity.

*Distribution:* Leonard formation (Glass Mountains) at 12, 25, 31, and T120. Word formation (Glass Mountains) at 144, T240a, 265, and Tc. Middle Gym formation (Sierra Diablo) at 414. Bone Canyon member (Delaware Mountains) at 490 and 491. Bone Canyon member (Guadalupe Mountains) at 221?. Leonard formation (Shafter) at 518.

### AMBOCOELIA PLANOCONVEXA (Shumard)

1855 *Spirifer plano-convexa.* Shumard, Missouri Geol. Survey Second Ann. Rept., p. 202.

1914 *Ambocoelia planoconvexa.* Kozlowski, Brachiopodes du Carb. Supér. de Bolivie, Ann. de Pal., vol. 9, pp. 76–77, text fig. 19, pl. 1, fig. 5; pl. 10, figs. 1–14.

1915 *Ambocoelia planiconvexa* Girty, Fauna of the Wewoka Formation, U. S. Geol. Sur. Bull. 544, pp. 94–96, pl. 11, figs. 6–7b. (See for synonnymy.)

*Observations:* This species occurs only in the *Uddenites* zone. At loc. 88s most of the specimens of it are pyritized, like the ammonoids from the same locality.

*Distribution:* Wolfcamp formation, *Uddenites* zone (Glass Mountains) at 88, T88, 88s, and 94s. Wolfcamp formation (Glass Mountains) at 195?.

### Genus MARTINIA McCoy

*Diagnosis* (after Weller): Shells generally of medium size or smaller, subcircular, subovate or subelliptical in outline, the hinge-line narrow, the cardinal extremities rounded, the surface of the valves smooth or nearly so. Cardinal area of the pedicle valve small, with poorly defined lateral margins, the surface curving with little or no differentiation into the lateral slopes of the valve; delthyrium rather large. Dental and septal plates not present; muscle scars small, scarcely impressed and ill-defined. In the brachial valve the muscle scars are also inconspicuous, the crura and primary lamellae of the brachidium are elongate, the bases of the spiral coils are situated well forward and their apices are directed toward the cardinal extremities.

Genotype: *Anomites glabra* Martin.

*Geologic range:* Devonian to Permian.

### MARTINIA RHOMBOIDALIS Girty

Pl. XLI, Figs. 6–10

1908 *Martinia rhomboidalis.* Girty, The Guadalupian Fauna, U. S. Geol. Sur. Prof. Paper 58, pp. 364–365, pl. 13, figs. 11–14c.

Middle Capitan, Capitan Peak (sta. 2926); basal Capitan, hill southwest of Guadalupe Point (sta. 2906); Delaware Mountain formation, Guadalupe Point (sta. 2919); Delaware Mountain formation, southern Delaware Mountains (stas. 2935, 2962).

*Description:* Shell medium-sized to large, length equal to or somewhat greater or less than the width. Hinge-line equal to about half the greatest width. General outline rhombic. Pedicle valve rather strongly convex in the posterior part, with decreasing curvature toward the front. Beak narrow and pointed, elevated and incurved over the not very high cardinal area. Median sinus broad and shallow, originating about 20 mm. in front of the beak as a flattening, becoming broader and deeper toward the front, which extends forward in the median part as a tongue. Brachial valve subcircular to subquadrate, with a small incurved beak, and an inflated umbo. From the median line the shell falls away rapidly to the lateral slopes, leaving a rounded subangular fold the entire length of the valve. Outer surface smooth. If the shell is partly exfoliated there are fine radial lirae. Dental and septal plates absent.

Dimensions in mm. of several specimens:

| | 1 (loc. 123) | 2 (loc. 8–14) | 3 (loc. 494) | 4 (Girty's type) |
|---|---|---|---|---|
| Length | 33 | 25 | 30 | 22 |
| Width | 24 | 24 | 37 | 20 |

*Observations:* Although this species occurs in great abundance at loc. 57, there being hundreds of specimens in the rock, few can be worked out to show the valves in conjunction. The species is not very common at any locality in the lower formations. It differs from *M. wolfcampensis* in having a smaller, less strongly inflated ventral umbo. *M. shumardiana* Girty, which is distinguished from *M. rhomboidalis* by its stronger fold and sinus, is hardly more than a variety of the latter. The closest foreign relatives of *M. rhomboidalis* are *M. elongata* Waagen and *M. variabilis* Gemmellaro, especially the latter. It would be difficult to point out differences without actual specimens for comparison.

*Distribution:* Leonard formation (Glass Mountains) at 8–14, 12?, 13?, T15, 16, 21, 26s, 34, 77, 98, T119, 120, 123, 124, T128, 232, 245, D1, D12, and D16. Word formation (Glass Mountains) at 6? and 241?. Bone Canyon member (Guadalupe Mountains) at 221. Victorio Peak member (Sierra Diablo) at 494. Altuda member (Glass Mountains) at 43 and 57 (very abundant).

**MARTINIA WOLCAMPENSIS King, n. sp.**

Pl. XL, Figs. 9–13

*Description:* Shell medium-sized, about equal in length and width. The hinge-line equals about half the greatest width, which is at about mid-length. Pedicle valve strongly convex in the posterior part, becoming more flattened toward the front. Ventral umbonal region broad and highly inflated, beak strongly incurved. Cardinal area high, not well defined, mostly occupied by a broad delthyrium. Near the front of large shells there is a broad, shallow, median sinus which extends forward at the anterior margin a short distance as a tongue. Brachial valve transversely elliptical, beak small and low. This valve has very low convexity, there being a convex median region and flattened lateral slopes. Surface smooth except for irregularly spaced radial lirae on some specimens, and finer concentric lirae. Dental and septal plates are absent. In the pedicle valve the muscle scars lie in an elongate depression along the median line.

Dimensions in mm. of several specimens:

| | 1 (loc. 94) | 2 (loc. 94s) | 3 (loc. 87) | 4 (loc. 201) |
|---|---|---|---|---|
| Length | 25 | 28 | 21 | 18 |
| Width | 24 | 32 | 22 | 17 |
| Thickness | 10+ | 11+ | 15 | 12 |

There is some variation in the development of the median sinus. The specimens from locs. 75 and 87, for example, have a well developed median depression at a size at which the specimens from loc. 94 are still evenly convex.

*Observations:* This species differs from *M. rhomboidalis* Girty in having a more convex umbonal region of the pedicle valve, a less convex brachial valve, and a regularly ovoid rather than rhomboid outline. Of the foreign species, it is closest to the specimens figured by Tschernyschew as *M. triqueta* Gemmellaro, and indeed it would be difficult to point out any differences between them. However, Tschernyschew's figures show considerable difference from true *M. triqueta,* which has a much less highly inflated ventral umbo. The similarity of the Uralian specimens to *M. wolfcampensis* is much greater, and it may be questioned if they are not identical.

*Distribution:* Wolfcamp formation, *Uddenites* zone (Glass Mountains) at 88, 88s (type), T88, 94, 94s, 95, 96, 175a, 201, and 202. Wolfcamp formation (Glass Mountains) at 75, 87 low, 93, and 196.

**Genus SPIRIFERINA d'Orbigny**

*Diagnosis:* Shell spiriferoid, transverse; fold and sinus angular and not sharply differentiated from the plicae, which are few in number and angular. Anterior margin of the valves strongly serrated, with a few sharp folds, decreasing in amplitude from and including the median fold. Ventral cardinal area curved, with more or less rounded margins which curve over into the lateral slopes; beak small and incurved. Delthyrium open; internally the dental lamellae are well-developed, and diverge toward the floor of the valve, and their anterior extremities are concave. Ventral median septum well-developed, extending farther forward than the dental lamellae. In the brachial valve the spiralia are directed laterally, the primary lamellae being joined by a complete, simple transverse jugum. Shell composed of two layers, of which the inner is strongly punctate and the outer covered with fine, tubular, spinose projections, which, when broken, give the shell a pustulose appearance. Concentric, irregularly arranged lamellae of growth are generally present.

Genotype: *Terebratulites rostrata* Schlotheim.

*Geologic range:* Middle Mississippian to Lias.

*Observations:* The name *Spiriferina* is here used as restricted by North, the lamellose forms with large fold and sinus being placed in *Punctospirifer*. According to the restricted usage, the only Pennsylvanian species of *Spiriferina* (except for the Mississippian holdovers in the Bend fauna) are *S. campestris* White and *S. gonionotus* Meek (if the latter is not identical with *S. campestris).* The common *S. kentuckyensis* is placed with *Punctospirifer*. Mississippian species of *Spiriferina* (restricted) are *S. norwoodana* (Hall), *S. salemensis* Weller, and *S. spinosa* (Norwood and Pratten). Girty,* in discussing the divisions of *Spiriferina* in the American Carboniferous, pointed out the presence of the two major subdivisions now distinguished as *Spiriferina ss.* and *Punctospirifer*. He suggests a different origin for the two groups of "*Spiriferina.*" The group now known as *Punctospirifer* has been derived by Hall and Clarke from the lamellose-septate group of spirifers *(Delthyris)*. Girty compares the spinose (or true) *Spiriferina* to some of the ostiolate spirifers, but says that since the ostiolate spirifers lack a ventral median septum, an attempt to derive *Spiriferina* from them would be rather fanciful. Continuing, Girty remarks that there is a little known group of spirifers (including *S. aciculifer* and *S. schucherti)* in the early Mississippian that have a spinose surface, but it was not known whether they were or were not septate and punctate, so that "it thus remains in doubt as to whether the *spinosa* group drew its ascent

*U. S. Geol. Sur. Prof. Paper 58, pp. 372–374. 1908.

from the same or different stock from *Spiriferina transversa" (Punctospirifer)*. In 1914, however, Weller established the genus *Acanthospira (Acanthospirina* Schuchert and LeVene), with *Spirifera aciculifera* Rowley as the genotype, for spinose spirifers having a non-punctate shell and no median septum. The presence of these characters strongly suggests that *Acanthospirina* is the connecting link between the ostiolate spirifers and *Spiriferina*. The geologic range of the three forms is still further suggestive. The ostiolate spirifers are especially characteristic of the Devonian. *Acanthospirina* occurs in the Louisiana limestone, of the lower Kinderhook. The first appearance of true *Spiriferina* is in the Salem limestone. Thus we see that *Acanthospirina* not only holds an intermediate position structurally but also chronologically. In conclusion, it may be stated as likely that *Punctospirifer* and *Spiriferina* were two differently derived groups of similar external form.*

### SPIRIFERINA ANGULATA KING, n. sp.

Pl. XLII, Figs. 12–13

*Description:* Shell large, much wider than long, with the greatest width at the hinge-line, semicircular, strongly biconvex. Cardinal area of the pedicle valve of moderate height, concave, beak rather low. Shell folded into very high, angular plications; median fold and sinus not very much larger than the plications and sulci, which are 4 or 5 in number on each side. The anterior margin is deeply serrate, the median digitation being as long as the entire thickness of the shell, and the lateral ones becoming gradually less toward the wings. The shell structure is strongly punctate, and there are occasional zig-zag growth varices.

Dimensions in mm. of several specimens:

| | 1 (loc. 478) | 2 (loc. 478) | 3 (loc. 479) |
|---|---|---|---|
| Length | 17 | 13 | 14? |
| Width | 28 | 22 | 28 |
| Thickness | 18 | — | — |

*Observations:* This species is distinguished by its strongly biconvex form and very high, sharp plications. I know of no other species of the genus in which there is a combination of these characters.

*Distribution:* Hess formation (Glass Mountains) at 106 and 211. Leonard formation (Glass Mountains) at Tba. Hess formation (Sierra Diablo) at 476, 478 (types), and 479.

### SPIRIFERINA HILLI Girty

Pl. XLII, Figs. 1–6

1908 *Spiriferina hilli.* Girty, The Guadalupian Fauna, U. S. Geol. Sur. Prof. Paper 58, p. 379, pl. 30, ffis. 15–15b.
Delaware Mountain formation, "Comanche Canyon," Glass Mountains.

1915 *Spiriferina Hilli.* Haack, Permfauna aus Nordmexiko, Zeit. der deutsch. geol. Gesell., 66 Bd. (1914), pp. 495–497, pl. 39, figs. 2–9.
Pichagüa, near Las Delicias.

*Description:* Shell small, wider than long, being widest somewhat in front of the hinge-line. Cardinal area high, rather variable in its degree of inclination to the shell margin; beak incurved. Median sinus distinctly larger than the lateral plications, but not very deep. According to Girty, it may contain a low median plication. Lateral plications 4 to 6 in number, decreasing in size laterally, the two plications bordering the sinus being considerably larger than the others. Brachial valve wider than long, with a small incurved beak. Median fold somewhat larger than the lateral plications, which number from 4 to 6 on each side. Surface smooth except for unevenly distributed growth varices, which are most closely spaced near the anterior and lateral margins. Shell structure rather coarsely punctate but not as coarsely as *S. laxa*. Some of the internal characters were described by Haack.

Dimensions in mm. of several specimens:

| | 1 (loc. 208) | 2 (loc. 223) | 3 (loc. D20) |
|---|---|---|---|
| Length | 12 | 6 | 9 |
| Width | 15 | 8 | 10 |

*Observations:* This species is distinguished from *S. laxa* by its much smaller size, greater length in proportion to width, and somewhat finer punctae. Haack states that the nearest foreign relative is *S. multiplicata* (Sowerby) rather than *S. cristata*, to which it was compared by Girty. *S. laxa* is more nearly related to the latter, both those forms being large and very coarsely punctate, while *S. multiplicata* and *S. hilli* are small and finely punctate.

*Distribution:* Hess formation (Glass Mountains) at 4, 208, and 223. Leonard formation (Glass Mountains) at 123?, T205?, 224, D3, D7, D9, and D20. Leonard formation transition beds (Shafter) at Th. Limestone of Pichagüa (Las Delicias) at Cerro Pichagüa.

### SPIRIFERINA LAXA Girty

Pl. XLII, Figs. 7–11

1908 *Spiriferina laxa.* Girty, The Guadalupian Fauna, U. S. Geol. Sur. Prof. Paper 58, pp. 377–378, pl. 21, figs. 3–3b.
Dark limestone, Pine Spring (sta. 2930) and Guadalupe Point (sta. 3762s?); Delaware Mountain formation, southern Delaware Mountains (sta. 2969?).

1908 *Spiriferina welleri.* Girty, *idem*, pp. 380–382, pl. 14, figs. 17–19b.
Middle Capitan, Capitan Peak (sta. 2926); basal Capitan, hill southwest of Guadalupe Point (sta. 2906); dark limestone, Pine Spring (sta. 2930?); Delaware Mountain formation, southern Delaware Mountains (sta. 2969?); Delaware Mountain formation, Glass Mountains.

1915 *Spiriferina Haarmanni.* Haack, Permfauna aus Nordmexiko, Zeitsch. der Deutsch. Geol. Gesell., bd. 66 (1914), pp. 492–495, pl. 38, figs. 7a–c; pl. 39, figs. 1a–c.
Pichagüa, near Las Delicias.

*Description:* Shell large, wider than long or of about equal length and width, with the greatest width at the hinge-line; more or less unsymmetrical. Shape of the pedicle valve transversely semicircular. The shell is not

*See North, F. J.: *Syringothyris* and *Spiriferina*, Quart. Jour. Geol. Soc. London, vol. 76, pt. 2 (1920), pp. 162–227.

very strongly convex. Cardinal area inclined backward at about 100° from the plane of the margin, concave near the apex but nearly flat near the hinge-margin, defined from the rest of the shell by an angulation near the beak but less clearly set off toward the sides. Beak pointed and incurved. Delthyrium large, higher than wide, bordered on each side by a ridge which, where it reaches the cardinal line, forms the outer surface of the hinge-tooth. Sinus broad but not much broader or deeper than the lateral sulci. Near the front it may contain one or two small plications. On each side of the sinus there are 4 or 5 large subangular plications, becoming smaller toward the sides. Brachial valve less strongly arched than the other, with a small, low beak and a very narrow cardinal area. The fold is only slightly larger than the lateral plications, which are 4 to 6 in number on each side.

Outer shell layer dense and finely papillose, marked near the front by several subimbricating lamellae. The main part of the shell is very coarsely punctate, the punctae being easily seen with the naked eye if the outer shell layer is worn away, when the shell is given a spongy appearance.

Within the pedicle valve the hinge-teeth have the form of blunt protuberances on each side of the delthyrium; their bases extend backward into the thickened dental lamellae, which reach to the bottom of the shell only in the apical region. Median septum thin, rising to a sharp crest just in front of the hinge-line. In the brachial valve there are well defined dental sockets, bordering which are protuberances that give rise on their anterior faces to the crura. In the median posterior part of the shell there is a thick ridge, the cardinal process.

Dimensions in mm. of several specimens:

| | 1 (loc. T240) | 2 (loc. 192) | 3 (loc. 46) | 4 (*S. welleri*) | 5 (*S. laxa*) | 6 (*S. haarmanni*) |
|---|---|---|---|---|---|---|
| Length | 16 | 16 | 12 | 17 | 16 | 22 |
| Width | 16 | 17 | 20 | 24 | 23 | 32 |
| Thickness | 9 | — | — | — | — | 17 |

It may be seen that the ratio of length to width is very variable in the Glass Mountains specimens, but that in the three species here placed in synonymy it is practically identical. (The figures for *S. haarmanni* are taken from Haack's plate rather than from the text.)

*Observations:* That the names *S. laxa, S. welleri,* and *S. haarmanni* have been applied to a single species seems certain. Any of the specimens in the present collection would fit equally the description of the three supposed species, and no significant difference can be found between any of the descriptions. A very well preserved specimen collected by myself from Cerro Pichagüia, the same hill from which the type of *S. haarmanni* was collected, shows no difference from the Word specimens. Several points of difference may, however, be noted between it and the types of Haack. Haack states that there are 3 or 4 punctae in the inner layer in the space of 1 mm., near the anterior margin. In my own specimen a measurement in the anterior part of the shell gives 8 punctae in 1 mm. If Haack's figures were correct the punctae would be incredibly large. Haack says that the median septum of the pedicle valve is not very high. This certainly must be due to imperfect preservation, for my own specimen has a high septum. The large pores said by Girty to exist in the outer shell layer of *S. laxa* were not observed in the present specimens.

*Spiriferina laxa* is characterized by its large size, large, high plications, which are not much smaller than the median fold, and by the coarsely punctate inner shell layer and papillose outer layer. The most similar American species is *S. pyramidalis* Girty, of the Capitan. The differences between them mentioned by Girty are the more rounded outline of *S. pyramidalis* and the higher cardinal area. Girty himself suggested that *S. pyramidalis* might be only a young stage of *S. laxa. Punctospirifer? pulchra* is a large, coarsely punctate shell, but has a relatively much broader fold and sinus, a greater number of lateral plications, and well-marked concentric lamellae. As Haack has shown, the closest relation of the species is to *S. cristata* (Schlotheim) and *S. nasuta* Waagen. The latter differs especially in that the fold and sinus are both broader and deeper. After comparing his specimens from Pichagüia with some from the Salt Range, Haack concludes that *S. haarmanni (=S. laxa)* is the American variant of the Indian and Chinese form.

*Distribution:* Word formation (Glass Mountains) at 46, T140, 142, 144, T144, T148, Tx148, 192, T239, T240, 247, 259, and 265. Altuda member (Glass Mountains) at 43. Bone Canyon member (Sierra Diablo) at 485?. Limestone of Pichagüia (Las Delicias) at Cerro Pichagüia. The species is common only at loc. 192.

### SPIRIFERINA PYRAMIDALIS Girty

Pl. XLII, Fig. 14

1908 *Spiriferina pyramidalis.* Girty, The Guadalupian Fauna, U. S. Geol. Sur. Prof. Paper 58, pp. 378–379, pl. 14, figs. 20–21b.

Middle Capitan, Capitan Peak (sta. 2926); Delaware Mountain formation, southern Delaware Mountains (sta. 2969?).

*Description:* Shell medium-sized, much wider than long, subtriangular, widest at the hinge-line. Pedicle valve high and conical, with a broad, high, nearly flat cardinal area, which is strongly defined by angles at its junction with the sides of the shell, and lies nearly perpendicular to the plane of the margin. Delthyrium large, nearly twice as high as wide. Median sinus not greatly larger than the lateral sulci. On each side of the sinus there are 4 strong plications, becoming smaller toward the wings. Brachial valve strongly convex, beak incurved over a narrow cardinal area. The median fold is much higher than the lateral plications, resulting in a strong deflection of the anterior margin. There are 3, and a faint fourth, lateral plications. Within the pedicle valve the median septum rises to a crest near its anterior end. Surface of both valves marked by fine, closely but irregularly distributed concentric lamellae. The shell is strongly punctate. Length of brachial valve, 11 mm.; width, 21 mm.; height of ventral cardinal area, 7.5 mm.

*Observations:* This species resembles those of *Punctospirifer* in having many concentric lamellae, but they are not as closely and regularly set as in that genus, and, moreover, *S. pyramidalis* has a relatively smaller fold

and sinus and much more extended ears than do most specimens of *Punctospirifer*. It is closely related to *S. laxa*, but has a much higher and less curved cardinal area.

*Distribution:* Word formation (Glass Mountains) at 264 (one specimen).

### Genus PUNCTOSPIRIFER North

*Diagnosis:* Shell spiriferoid, wider than long, with the greatest width near the hinge-line; cardinal angles slightly rounded or subangular; cardinal area moderately high and concave. Median fold and sinus considerably larger than the plications, which are numerous, small, and rounded. Surface of both valves crossed by regularly disposed imbricating lamellae, which in adult specimens may be more or less obsolescent, especially in the posterior part of the valve. Delicate hair-like spinules of varying abundance and arrangement are also present. Shell structure fibrous and strongly punctate. Within the pedicle valve the dental lamellae are slightly divergent; median septum well developed, thickened at its base, terminating somewhat abruptly about half-way toward the anterior margin. No marked development of an apical callosity, the three vertical septa appearing as distinct structures in sections near the beak. A low median crest bisects the muscle scars of the brachial valve; spiralia large, connected by a slender jugum, which forms a shallow V-shaped process, with its apex directed ventrally and backward.*

Genotype: *Punctospirifer scabricosta* North.

*Geologic range:* Lower Mississippian to Permian (at least).

*Observations:* This genus differs from *Spiriferina* in the following respects: It is more alate. There is a greater number of plications, which are smaller and less angular, and the fold and sinus are relatively smaller. The surface is lamellose and finely spinose, rather than smooth except for irregularly arranged growth lamellae and relatively large spines. Finally, the jugum is V-shaped with the apex directed backward, rather than straight, or curved toward the front* as in *Spiriferina*. In establishing this genus, North emphasized especially the relatively greater size of the fold and sinus, rather than the lamellose surface. Following Girty* I would regard the latter character as of primary importance. Unfortunately, neither the specimens which I studied, nor those described by Weller in "Mississippian Brachiopoda" showed the structure of the jugum, but since *Punctospirifer kentuckyensis* as described by Hall and Clarke has an angular, backward pointing jugum, it seems likely that the other American forms having the external characters of *Punctospirifer* would agree in that respect. The Mississippian species of *Punctospirifer* are *P. subtexta* (White), *P. solidirostris* (White), *P. subelliptica* (McChesney), and *P. transversa* (McChesney). The only Pennsylvanian species above the Bend is *P. kentuckyensis* (Shumard).

Hall and Clarke have shown that the group now comprised in *Punctospirifer* probably arose from *Delthyris*.* As indicated in the discussion of *Spiriferina*, that genus and *Punctospirifer* may have had independent origins.

*North, F. J.: *Syringothyris* and *Spiriferina*, Quart. Jour. Geol. Soc. London, vol. 76, pt. 2 (1920), pp. 208–214.

*See Kozlowski's figure of *Spiriferina campestris* White, in Ann. de Pal., vol. 9, text fig. 17.

*U. S. Geol. Sur. Prof. Paper 58, pp. 371–373.

### PUNCTOSPIRIFER BILLINGSII (Shumard)

Pl. XLII, Figs. 15–18

1858 *Spiriferina Billingsii.* Shumard, Trans. Acad. Sci. St. Louis, vol. 1, p. 294 (date of volume, 1860). White and dark (Permian) limestone: Guadalupe Mountains, and conglomerate at mouth of Delaware Creek, New Mexico.

1859 *Spiriferina Billingsii.* Shumard, *idem.*, p. 391. White and dark (Permian) limestone: Guadalupe Mountains.

1908 *Spiriferina Billingsii.* Girty, The Guadalupian Fauna, U. S. Geol. Sur. Prof. Paper 58, pp. 374–376, pl. 13, figs. 16–19d, 21–21b, 24–24c; pl. 14, figs. 15–16. Middle Capitan, Capitan Peak (sta. 2926); dark limestone, Pine Spring (sta. 2930); Delaware Mountain formation, Guadalupe Point (sta. 2930?); Delaware Mountain formation, southern Delaware Mountains (sta. 2969?); Delaware Mountain formation, "Comanche Canyon," Glass Mountains.

*Description:* Shell medium-sized, wider than long, strongly biconvex, with the hinge-line shorter than the greatest width of the shell, cardinal angles rounded. Pedicle valve greatly arched, with a high, extended beak, which is incurved over a broad triangular, concave, poorly-defined cardinal area. Median sinus broad and angular, extending from the apex to the front. The lateral slopes each bear from 4 to 9 small rounded plications, which are simple and gradually enlarge from the beak to the front. Brachial valve evenly convex, with a strong median fold, elevated toward the front above the general convexity. The lateral slopes bear small plications corresponding to those of the pedicle valve. Surface marked by regular, strong imbricating, concentric lamellae, from 7 to 13 of which occur in the space of 5 mm.

Dimensions in mm. of several specimens:

| | 1 (loc. T148) | 2 (loc. Tx) | 3 (Shumard's type) |
|---|---|---|---|
| Length | 11 | 10 | 18.5 |
| Width | 12 | 11 | 22.5 |
| Thickness | 9 | — | 14.5 |

*Observations:* Save for the smallest size of the specimens from the Glass Mountains, there is no difference between them and the typical ones as described by Shumard and Girty. Girty recognized three varieties of *Spiriferina billingsii: S. billingsii retusa, S. evax,* and *S. sulcata,* all

*According to Miss Muir-Wood (Notes on the Silurian brachipod genera *Delthyris, Uncinulina,* and *Meristina,* Ann. and Mag. of Nat. Hist., ser. 9, vol. 15, 1925, p. 83) some of the species of *Delthyris* in the Upper Devonian and Mississippian should be placed in *Tylothyris* North. In his original description, North (Quart. Jour. Geol. Soc. London, vol. 76, pt. 2, 1920, p. 195) said that his genus *Tylothyris* differed from the Silurian *Delthyris* in its much greater size, smaller and more numerous costae, and the marked development of an apical callosity. None of these would ordinarily be regarded as valid for generic distinction. Miss Muir-Wood, however, shows that in true *Delthyris* the lamellae are fimbriate, while in *Tylothyris* no ornament has been described on the lamellae. The Upper Devonian and Lower Mississippian shells formerly classed as *Delthyris* which do not have fimbriate lamellae may, she suggests, be *Tylothyris.*

from the Capitan. None of them was recognized in the present collections. The most closely related foreign species is *P. ornata* (Waagen), which differs in having larger and fewer lateral plications. *P. billingsii* is a larger form than *P. kentuckyensis,* and, though related closely to *P.? pulchra* of the Phosphoria, has a quite different configuration.

*Distribution:* Word formation (Glass Mountains) at 54, 144, T148, 192, T239, 241, 243, Ta, Tc, and Tx. Altuda member (Glass Mountains) at 43 and 50. Not common at any locality.

#### PUNCTOSPIRIFER KENTUCKYENSIS (Shumard)

1855 *Spirifer Kentuckyensis.* Shumard, Missouri Geol. Survey Second Ann. Rept., p. 205.

1915 *Spiriferina kentuckyensis.* Girty, Fauna of the Wewoka Formation, U. S. Geol. Sur. Bull. 544, pp. 85–87, pl. 11, figs. 8–8a.

*Distribution:* Wolfcamp formation, *Uddenites* zone (Glass Mountains) at T88. Wolfcamp formation (Glass Mountains) at 89. Rare.

### Family RHYNCHOSPIRINIDAE

#### Genus HUSTEDIA Hall

*Diagnosis* (after Weller): Shells small, subovate in outline, with nearly equally convex valves, marked by rather coarse, subangular, simple plications, the shell structure punctate. Pedicle valve with a prominent umbo perforated at its apex by a subcircular foramen; cardinal area small, with sharply defined lateral margins, the delthyrium closed by deltidial plates and in contact with the foramen at its apex only. Internally the inner surface of the pseudodeltidium bears a split tube attached by its closed side to the deltidial plates, with its open side directed toward the interior of the shell. In the brachial valve the hinge-plate projects considerably beyond the hinge-line posteriorly, being recurved into the umbonal cavity of the pedicle valve, the upper face convex and elevated medially, the posterior margin sinuate and crescentic with the horns of the crescent very short; from the lateral margins of the plate arise a pair of strong lobes which bear the erect slightly recurved crura. At the base of the cardinal process and in the median line, there arises a free, slender, lingulate process which curves upward and backward with somewhat less curvature than the hinge-plate, and rises to the highest point attained by the latter; the inner surface of this process is deeply grooved and at its base it is supported by a median septum which extends for a third the length of the valve. Spiralia approximate, with 8 or 9 volutions, the apices directed laterally; jugum posterior in position, regularly curved from the points of origin on the primary lamellae forward and inward, giving rise at its center to a simple stem-like process. The posterior margins of the coils and the jugum are fimbriated.

Genotype: *Terebratula mormonii* Marcou.

*Geologic range:* Mississippian to Permian.

*Observations:* This genus is distinguished from *Eumetria* by having fimbriate spiralia and jugum and a simple rather than forked extremity of the stem-like process from the jugum. Most species have coarser plications than those of *Eumetria,* but *H. pusilla* is finely plicate.

#### HUSTEDIA BIPARTITA Girty

Pl. XLII, Fig. 19

1908 *Hustedia bipartita.* Girty, The Guadalupian Fauna, U. S. Geol. Sur. Prof. Paper 58, pp. 398–399, pl. 30, figs. 19–20a.

Dark limestone, Pine Spring (sta. 2930?) and Guadalupe Point (sta. 3762e?), Guadalupe Mountains. Delaware Mountain formation, southern Delaware Mountains (sta. 2969); Delaware Mountain formation, "Comanche Canyon," Glass Mountains.

*Description:* Shell similar to *H. meekana* but with the median plication of the brachial valve undeveloped at the beak and never reaching the size of the plications on each side, with the result that there is a median depression in the general surface of the valve. There are 11 plications on the brachial valve of the specimen of *H. bipartita* in the present collection, and the beak, as in *H. meekana,* is small, pointed, and erect. Length, 9 mm., width, 8 mm., thickness, 6 mm.

*Observations:* Perhaps this group should be regarded as no more than a variety of *H. meekana,* for in the latter species the median plication of the brachial valve may at its origin be somewhat smaller than those on each side.

*Distribution:* Leonard formation (Glass Mountains) at D2.

#### HUSTEDIA HESSENSIS King, n. sp.

Pl. XLII, Figs. 43–46

*Description:* Shell rather large for the genus, moderately convex, longer than wide. Pedicle valve evenly rounded at the front, with nearly straight posterior lateral slopes which converge to meet at the beak at an angle of about 70°. Ventral beak high, only slightly incurved; cardinal area well defined, somewhat concave, more than two-thirds as high as wide. Foramen large and subcircular. Both valves are folded into high, broad, subangular plications, averaging about 10 in number; the plications may be somewhat irregular, some of them being higher than the adjoining ones. Surface marked by fine, irregular, concentric growth lirae. Internal characters not determined.

Dimensions in mm. of several specimens:

| | 1 (loc. 478) | 2 (loc. 106) | 3 (loc. 208) | 4 (loc. 474) |
|---|---|---|---|---|
| Length | 14 | 16 | 15 | 10 |
| Width | 11 | 14 | 14 | 9 |
| Thickness | 10 | 10 | — | 5 |

*Observations:* This species is distinguished by the large size, angularity, and irregularity of its plications, its acutely pointed posterior part of the pedicle valve, its elevated ventral umbo, and its high cardinal area. A closely related, if not identical, species is *Retzia compressa* Meek, from the Permian of California. The description and figures of that species are inadequate for proper comparison.

*Distribution:* Hess formation (Glass Mountains) at 105, 106, 208, 211, 212, and 222. Hess formation (Sierra Diablo) at 469, 474, 476, 478 (types), and 479.

#### HUSTEDIA HUECOENSIS King, n. sp.

Pl. XLII, Figs. 40–42

*Description:* Shell large for the genus, outline subcircular or elongate ovoid, convexity low and about the same in

both valves. Ventral beak high, not greatly incurved over the small, well defined cardinal area. Apical foramen large and round. Surface ornamented by 16 to 18 low, subangular plications. Fold and sinus absent, though the median plication or sulcus may be slightly larger than the others.

Dimensions in mm. of several specimens:

| | 1 (loc. 440a) | 2 (loc. 440a) | 3 (loc. 440a) | 4 (loc. 440b) |
|---|---|---|---|---|
| Length | 14 | 15 | 13 | 9 |
| Width | 13 | 15 | 10.5 | 7 |
| Thickness | 8 | 7 | 9.5 | 5 |

Number 4 is probably an immature specimen of this species, and number 3 is an exceptionally narrow, thick specimen which is considerably different from the typical form of the species, but appears to be connected with the typical form by complete gradation.

*Observations:* The typical form of this species is distinguished from others in the west Texas Permian by the combination of low convexity, relatively large size, and numerous plications. The narrow forms are less easily distinguishable from *H. meekana,* but are more finely plicated. *H. hessensis* similarly has low convexity, but there are fewer and sharper plications.

*Distribution:* Upper Gym formation (Hueco Mountains) at 437, 440a, and 440b.

### HUSTEDIA MEEKANA (Shumard)

Pl. XLII, Figs. 34–39

1858 *Retzia* (?) *Meekana.* Shumard, Trans. Acad. Sci. St. Louis, vol. 1, p. 295 (date of volume, 1860).
White and dark (Permian) limestone: Guadalupe Mountains; conglomerate at the mouth of Delaware Creek.

1859 *Retzia Meekana.* Shumard, *idem,* p. 395, pl. 11, figs. 7a–b.
White and dark limestone: Guadalupe Mountains; conglomerate at the mouth of Delaware Creek.

1908 *Hustedia meekana.* Girty, The Guadalupian Fauna, U. S. Geol. Surv. Prof. Paper 58, pp. 394–396, pl. 14, figs. 22–26a; pl. 21, figs. 5–8a; pl. 24, figs. 14–14a; pl. 29, fig. 8; pl. 30, figs. 16–17.
Top of Capitan (sta. 3762a), middle Capitan, Capitan Peak (sta. 2926); basal Capitan, hill southwest of Guadalupe Point (sta. 2906); dark limestone, Pine Spring (sta. 2930), Guadalupe Point (stas. 3762b and 3762d), hill southwest of Guadalupe Point (sta. 2924); basal black limestone, Guadalupe Point (stas. 2920 and 2967); Delaware Mountain formation, southern Delaware Mountains (stas. 2969 and 3500); Delaware Mountain formation, "Comanche Canyon," Glass Mountains.

1915 *Retzia (Hustedia) Meekana.* Haack, Permfauna aus Nordmexiko, Zeitsch. der deutsch. geol. Gesell., bd. 66 (1914), pp. 497–501, text figs. 1–2; pl. 39, figs. 10a–c.
Pichagua, near Las Delicias.

*Description:* Shell medium-sized to large for the genus, longer than wide, valves subequally convex, with a subcircular brachial valve and a high, pointed pedicle valve. Hinge-line equal in width to about a fourth the width of the shell. Cardinal area of the pedicle valve well defined, a little less than twice as wide as high, regularly incurved. Apical foramen large and ovoid. On average specimens there are 12 plications on the pedicle valve and 11 on the brachial, but the last one or two on each side are very small. The plications are large and subangular, and the sulci are the reverse of the plications. There are faint growth varices, especially near the front. Shell finely punctate. The internal structures have been described in detail by Haack, who shows that they are similar to those of *H. mormonii,* except that the prongs on each side of the cardinal process are stronger and higher and the median septum shorter; moreover, the points of origin of the jugum from the beginning of each of the spiralia are said to be broader.

Dimensions in mm. of several specimens:

| | 1 (loc. 123) | 2 (loc. 46) | 3 (loc. 491) | 4 (loc. 236) | 5 (Shumard's type) |
|---|---|---|---|---|---|
| Length | 10 | 12.5 | 11.5 | 18 | 11.5 |
| Width | 9 | 11.5 | 9.5 | 19 | 10.5 |
| Thickness | 7 | 8.5 | 8.0 | — | 9.0 |

*Observations:* According to Girty, there are generally 9 plications on the brachial and 10 on the pedicle valve. The greater number mentioned in the above description may be accounted for by the fact that the plications farthest on each side are very small, and show only in very well preserved specimens. Girty has pointed out that a feature mentioned by Shumard as an important specific character, the supposed presence of small costae superimposed on the larger ones, is due to exfoliation of the outer surface. This probably is shown more in the Capitan specimens than in those of the present collection, most of which are silicified. *H. meekana* differs from *H. mormonii* in having larger and fewer plications, a more elongate and erect beak, and in attaining a larger size. Haack has shown that *H. meekana* is very close to the specimens grouped by Waagen in his species *Eumetria indica.* They are said to differ in that the latter has perhaps smaller costae, somewhat greater length than width, and a more strongly convex pedicle valve. There is, however, enough variation among the specimens of the present collection to include some such minor differences from the typical form, and it may be that Waagen's species is not even varietally distinct, as regarded by Haack. Haack suggested that the Mexican form be distinguished as var. *mexicana* because of its supposed greater arching of the brachial valve, but from his figures the distinction hardly seems justifiable. The brachial valve is not preserved in my own specimens from Mexico.

*Distribution:* Hess formation (Glass Mountains) at 4, 17a, 105, 107, T107, 223, and Tv. Leonard formation (Glass Mountains) at 7, 8, 12, 15, T15a, 19, 25, 26, 37, 82, 83, T119, 120, 123, 124, 128, 232, T233, 234, 236, Tb, Tba, D2, D3, D5, D7, D9, D17, D18, and D22. Word formation (Glass Mountains) at 44–46, 46, 51, 54, 55, T141, 144, 148, T148, Tx148, 154, 162, 168, 192, 237, T239, D239, T240a, 241, 243, T243, 245, 246, 247, 248, 253, 265, and Ta. Altuda member (Glass Mountains) at 42, 43, and 50. Bone Canyon member (Delaware Mountains) at

491. Victorio Peak member (Sierra Diablo) at 499. Leonard formation (Baylor Mountains) at 502 and 503. Delaware Mountain formation (Sierra Diablo) at 504. Leonard formation (Shafter) at 515 and 518. Limestone of Cerro Caballo (Las Delicias) at section 1, beds 11, 12, and 14. Limestone of Pichagüa (Las Delicias) at Cerro Pichagüia.

**HUSTEDIA MORMONII (Marcou)**

Pl. XLII, Fig. 33

1858 *Terebratula Mormonii.* Marcou, Geology of North America, p. 51, pl. 6, figs. 11–11c.

1914 *Hustedia mormonii.* Kozlowski, Brachiopodes du Carb. Super. de Bolivie, Ann. de Pal., vol. 9, pp. 77–78, pl. 1, fig. 2; pl. 9, figs. 1–34; pl. 10, figs. 34–35. (See for best description and figures.)

1915 *Hustedia mormoni.* Girty, Fauna of the Wewoka Formation, U. S. Geol. Sur. Bull. 544, pp. 103–104, pl. 12, figs. 5–6a. (See for synonymy.)

*Distribution:* Wolfcamp formation, *Uddenites* zone (Glass Mountains) at 88, T88, and 94. Wolfcamp formation (Glass Mountains) at 75?, 87 low, 87 high, 89, 90, 91, T91, 93, T93, 193, and 196. Lower Gym formation (Hueco Mountains) at 376, 377a, 389, 290, 395, and 401.

**HUSTEDIA MORMONII PAPILLATA (Shumard)**

Pl. XLII, Figs. 27–32

1858 *Retzia papillata.* Shumard, Trans. Acad. Sci. St. Louis, vol. 1, p. 294 (date of volume, 1860).
Dark (Permian) limestone: Guadalupe Mountains.

1859 *Retzia papillata.* Shumard, *idem.*, p. 395, pl. 11, figs. 9a–c.
Dark (Permian) limestone: Guadalupe Mountains.

1908 *Hustedia papillata.* Girty, The Guadalupian Fauna, U. S. Geol. Sur. Prof. Paper 58, pp. 397–398, pl. 30, figs. 18–18b.
Basal Capitan, hill southwest of Guadalupe Point (sta. 2906?); dark limestone, Pine Spring (sta. 2930); Delaware Mountain formation, Guadalupe Point (sta. 2967?); Delaware Mountain formation, "Comanche Canyon," Glass Mountains.

*Description:* Shell small, elongate ovate, valves subequally convex. Beak of the pedicle valve long, not greatly incurved over the small concave cardinal area, which is well defined from the rest of the shell. Apical foramen large and round. Median fold and sinus absent. Plications subangular; 12 to 18 in number on the pedicle valve and from 11 to 17 on the brachial. Surface finely papillose.

Dimensions in mm. of several specimens:

| | 1 (loc. 54) | 2 (loc. 243) | 3 (loc. Tx) | 4 (Shumard's type) |
|---|---|---|---|---|
| Length | 9.5 | 9 | 7 | 10 |
| Width | 7.5 | 7 | 5.5 | 8 |
| Thickness | 6 | 5.5 | 4 | 7 |

*Observations:* This variety is hardly distinguishable from *H. mormonii.* It seems generally to have the ventral beak less incurved than in that species. It is distinguished from *H. meekana* by being more elongate and more finely plicated. Shumard originally supposed his species to be characterized by its papillose surface, but that feature is, of course, present in any species of the genus. The greater number of plications mentioned in the above description than in the descriptions of Girty and Shumard is due to the counting of the very small plications on the sides.

*Distribution:* Leonard formation (Glass Mountains) at 124, 128, T226, 233, D3, and D7. Word formation (Glass Mountains) at 6, 46, 54, T140, 143, 144, 238, 239, 243, T251, 264, Tu, and Tx.

**HUSTEDIA PUSILLA (Girty)**

Pl. XLII, Figs. 20–26

1908 *Pugnax? pusilla.* Girty, The Guadalupian Fauna, U. S. Geol. Sur. Prof. Paper 58, pp. 319–320, pl. 24, figs. 18–18b.
Basal black limestone, Guadalupe Point (sta. 2967).

*Description:* Shell small, moderately convex, somewhat longer than wide. Pedicle valve evenly rounded at the front, with nearly straight posterior lateral slopes which meet at the beak at somewhat less than a right angle. Cardinal area well defined, about twice as wide as high, not strongly convex. Apical foramen large and round. Some specimens have a well-marked, broad median deflection of the front margin and even a trace of a median fold in the brachial valve and sinus in the pedicle valve, but in other specimens the margin is straight except for the zigzag of the plications. The shell is folded into fine, subrounded plications, averaging about 16 in number, which originate at the beak.

Within the brachial valve the crura project ventrally, then recurve sharply to join the primary lamellae, which first extend dorsally parallel to the crura, then curve about parallel to the roof of the brachial valve. On each side the jugum arises from the distal side of the primary lamella, curves ventrally around the lamella and extends proximally into a small plate which lies parallel to the plane of the primary lamella; the ventral side of this plate gives rise to a lamella which extends ventrally, and near the middle of the posterior part of the shell joins the lamella of the jugum which arises from the other primary lamella at an acute angle; the united lamellae of the jugum then are produced obliquely toward the apical part of the valve in a sharply pointed spur. The spiralia are coiled at least 6 times. The edges of the spiralia and the posterior process of the jugum are set with fine spines.

Dimensions in mm. of several specimens:

| | 1 (loc. 46) | 2 (loc. T236) | 3 (loc. 44–46) | 4 (loc. 46) |
|---|---|---|---|---|
| Length | 8 | 6.5 | 6 | 12 |
| Width | 6.5 | 6 | 6 | 9.5 |
| Thickness | 4 | 3.5 | 2.5 | — |

*Observations:* This species is distinguished externally by its fine plications and by the occasional presence of a median fold and sinus. Internally the mode of origin of the jugum from the primary lamellae is different from that described in other species of the genus, in which the jugum appears to arise directly from the ventral side of the primary lamella on each side. Otherwise the internal characters are like those of other species of *Hustedia*

rather than like *Uncinella*, to which there is considerable external resemblance. The specimens of the present collection are undoubtedly the same species as Girty's *Pugnax? pusilla*. The apical foramen figured in Girty's specimen would hardly be present in a rhynchonellid.

*Distribution:* Wolfcamp formation (Glass Mountains) at 195a?. Hess formation (Glass Mountains) at T82a and D2. Word formation (Glass Mountains) at 44–46, 46, 51, 53, 54, T236, and 241?. Limestone of Cerro Caballo (Las Delicias) at section 1, bed 14, west of the Noria. Limestone of Pichigüia (Las Delicias) at Cerro Pichagüia.

## Family ATHYRIDAE

### Genus COMPOSITA Brown

*Diagnosis* (after Weller): Shells small or medium-sized, subovate, subquadrangular to subpentagonal in outline, biconvex, with the fold and sinus developed in the anterior part of the shell or in some extending backward to the umbonal region; both the fold and sinus may be marked by a rather sharp median sulcus. The surface of the valves smooth or marked only by concentric growth-lines which are not extended into lamellae. Beak of the pedicle valve incurved so as to conceal the delthyrium, but the foramen generally exposed, encroaching upon the umbonal region of the valve. Internally the hinge-teeth are prominent, recurved at the tips, and supported by short but rather stout dental lamellae which are not produced forward about the muscular area; between the dental lamellae is a deep, transversely striated pedicle cavity, and in front of this an ovate muscular area extending about half the length of the valve; the diductor scars are faintly developed, and the adductors occupy a narrow, more sharply defined central area. In the brachial valve the dental sockets are broad and deep, the socket walls are connected by a hinge-plate which is extended posteriorly beyond the margin of the valve into the umbonal cavity of the opposite valve; the lateral margins are thickened and are produced forward into the crura. Spiralia and jugum similar to those of *Athyris*.

Genotype: *Spirifer ambiguus* Sowerby.

*Geologic range:* Mississippian to Permian.

### COMPOSITA EMARGINATA AFFINIS Girty

Pl. XLIII, Figs. 12–17

1908 *Composita emarginata* var. *affinis*. Girty, The Guadalupian Fauna, U. S. Geol. Sur. Prof. Paper 58, p. 389, pl. 15, figs. 6–7b.

Middle Capitan, Capitan Peak (sta. 2926); Delaware Mountain formation, southern Delaware Mountains (stas. 2969? and 3500?); Delaware Mountain formation, "Comanche Canyon," Glass Mountains.

*Description:* Shell large, suboval to subpentagonal, moderately convex, both valves being of about the same depth. The proportion of length to width varies considerably, but all specimens are longer than wide, and relatively more elongate than the average of specimens of *C. subtilita*. Beak of the pedicle valve elevated and incurved, with a round foramen. Near the front of most specimens there is a shallow median depression at the front of the pedicle valve, which is more or less prolonged as a tongue. In the brachial valve the shell is regularly arched transversely, except (in most specimens) at the front, which is slightly more elevated, corresponding to the sinus of the other valve. The fold and sinus are developed only in the anterior part of the shell, and generally only at a much larger size than in *C. subtilita*. Surface smooth except for a few lamellose concentric elevations near the front. Some specimens have a semblance of indistinct radial lirae. Within the pedicle valve there is a deep, longitudinally striated pedicle cavity between the short dental lamellae; in front of it there is an ovate, rather faintly defined area containing the flabellate diductors and the narrow adductor scars. The surrounding pallial region is pitted. In the brachial valve the hinge-plate is large and similar to that of *C. subtilita*.

Dimensions in mm. of several specimens:

| | 1 (loc. 247) | 2 (loc. 247) | 3 (loc. 247) | 4 (loc. 145) | 5 (Girty's specimen) |
|---|---|---|---|---|---|
| Length | 33 | 31 | 31 | 32 | 26 |
| Width | 29 | 25 | 29 | 25 | 20 |
| Thickness | 17 | 16 | 18 | 19 | 14 |

*Observations:* Were it not for Girty's statement that "this form graduates more or less completely into *Composita emarginata*" I would have no hesitation in recognizing it as distinct. Characteristic forms are quite different from the typical Capitan species, which is characterized by the presence of a median sinus on both valves, which causes the anterior margin to recede. No such character was seen in any of the specimens here examined. Specimens from the Capitan labeled by Girty as *Composita emarginata* seem, however, to agree better with the form here described. *C. emarginata affinis* differs from *C. subtilita* in its lower relative convexity and the development of the fold and sinus only in the anterior part of large shells, at a size much larger than in typical *C. subtilita*. This variety is slightly larger than average specimens of *C. subtilita*. It is rather difficult to distinguish *C. emarginata affinis* from some forms of *C. subtilita*.

*Distribution:* Middle and upper Word formation (Glass Mountains) at 47, 144, T144, 145, T148, Tx148, 150, 159?, 162, 192, 240?, T240, 243?, 246, 247 (abundant), 248, 257, and 259?. Altuda member (Glass Mountains) at 43, 50?, and 47?. Delaware Mountain formation (Sierra Diablo) at 504. Limestone of Cerro Caballo (Las Delicias) at section 1, beds 12 and 14.

### COMPOSITA MEXICANA (Hall)

Pl. XLIII, Figs. 1–11

1857 *Terebratula mexicana*. Hall, Emory's Rept. U. S. and Mexican Boundary Surv., vol. 1, pl. 20, fig. 2. (Carboniferous: Texas?)

1908 *Composita mexicana*. Girty, U. S. Geol. Sur. Prof. Paper 58, pp. 389–390.

Delaware Mountain formation, "Comanche Canyon," Glass Mountains.

1908 *Composita mexicana* var. *guadalupensis*. Girty, *idem*, p. 390, pl. 24, figs. 11–13b.

Basal black limestone, Guadalupe Point (stas. 2920 and 2967).

1909 *Composita mexicana.* Girty, Fauna of the Manzano Group, U. S. Geol. Sur. Bull. 389, p. 68, pl. 8, fig. 1.

Yeso formation, Mesa del Yeso; San Andres formation, San Andres.

*Description:* Shell small, subpentagonal, widest at about two-fifths the distance from the front margin to the beak; width about equal to the length. Ventral beak large and inflated. Sinus strongly developed toward the front, producing a strong forward and upward deflection of the anterior margin. Brachial valve rather strongly convex, bearing near the front a high, narrow fold, which is accentuated by a strong depression of the shell on each side. Concentric growth lamellae are unevenly distributed over the surface, and may be closely crowded near the front. There is considerable variation in the configuration of this species. The convexity is variable, and the length may exceed the width. The amount of development of the fold and sinus varies greatly, some shells having hardly any at all, and thus being indistinguishable from some small specimens of *C. subtilita*, while others are very strongly emarginate at the front, both the median fold and sinus and the bordering reflexions of the shell forming waves in the margin.

Dimensions in mm. of several specimens:

| | 1 (loc. 436) | 2 (loc. 397) | 3 (loc. 223) | 4 (loc. 439) | 5 (loc. 451) |
|---|---|---|---|---|---|
| Length | 11 | 17 | 15 | 13 | 14 |
| Width | 10 | 15 | 13 | 14 | 14 |
| Thickness | 7 | 11 | 10 | 9 | 7 |

*Observations:* It is hardly feasible to attempt to separate from typical *C. mexicana* the form distinguished by Girty as *C. mexicana guadalupensis*, the only distinguishing features of which are the lower fold and shallower sinus, and lower convexity. There is so much variation in the species that a number of varieties might be recognized, but their distinction would hardly be profitable. *C. mexicana* typically is quite different from *C. subtilita*, but intermediate types exist. It is much smaller and has a much more strongly marked fold and sinus.

*Distribution:* Hess formation (Glass Mountains) at 1, 112, 113, and 223 (abundant). Leonard formation (Glass Mountains) at 123 and 233. Lower Gym formation (Hueco Mountains) at 376, 377b, 379, 389, 381, 390, and Tzzaa. Middle Gym formation Hueco Mountains) at 418, 419, 420, and 424. Upper Gym formation (Hueco Mountains) at 436b, 437b, 438, 439, 440a, 440b, 442, 443, 447, and 449. Gym formation (Diablo Plateau) at 446 and 451. Hess formation (Sierra Diablo) at 462 and Te (abundant). Victorio Peak member (Sierra Diablo) at 499. Leonard? formation (Shafter), thin-bedded zone, at Tj.

### COMPOSITA MIRA (Girty)

Pl. X, Fig. 13; Pl. XLIV, Figs. 3–8

1877 *Athyris roissyi.* Meek, U. S. Geol. Expl. 49th Par. Final Rept., vol. 4, pt. 1, p. 82, pl. 9, figs. 3–3b.

Carboniferous limestone: Ruby Group; Wachoe Mountains; Mahogany Peak; Egan Range, Nevada.

1899 *Athyris mira.* Girty, Geology of the Yellowstone National Park, U. S. Geol. Sur. Mon. 32, pt. 2, p. 570.

1926 *Composita mira* Girty, in Ore Deposits of Utah, U. S. Geol. Sur. Prof. Paper 111, pp. 646–647, pl. 56, figs. 16–16a.

Permian: Utah.

*Description:* Shell medium-sized to large, subcircular, not very strongly convex. Beak of the pedicle valve broad and incurved, with a round apical foramen. Beak of the brachial valve small and obtusely pointed. Near the front of the pedicle valve there is a low, narrow sinus, not greatly extended forward as a tongue. In the brachial valve there is no break in the regularity of the transverse convexity except at the front, where there is a low elevation corresponding to the sinus of the other valve. The surface is marked by strong, irregularly disposed varices of growth, which, because of the small amount of forward extension of the sinus, have the form of regular curves, the forward projection corresponding to the development of the sinus being present only in the anterior part.

Dimensions in mm. of several specimens:

| | 1 (loc. 174) | 2 (Las Delicias, sec. 1, bed 11) | 3 (Meek's type) |
|---|---|---|---|
| Length | 23 | 27 | 42 |
| Width | 22 | 27 | 43 |
| Thickness | 14 | — | 20 |

*Observations:* This species is similar in a general way to *C. subtilita*, but typical forms of the two are quite different. The outline is more rounded, the fold and sinus are less strongly developed, and the ventral sinus is not prolonged as far forward as a tongue, the convexity is lower, and the growth varices are stronger. In addition, the type specimen is much larger than most specimens of *C. subtilita* and has more numerous growth lamellae, but in these respects the Texas and Mexican specimens show little difference from *C. subtilita*. There is a great deal of variation in this species within certain limits. The relative convexity and the strength of the sinus are especially variable, but they all show greater similarity to *C. mira* than to *C. subtilita*. *C. mira* differs from *C. emarginata affinis* mainly by its rounded rather than elongate outline; indeed it is quite likely that it is a species intermediate between *C. subtilita*, which occurs as high as the Leonard, and *C. emarginata affinis*, which first appears in the middle Word.

*Distribution:* Leonard formation (Glass Mountains) at 13, 21, 123, 124, 126, 128, 174, Tb, D3, and D17. Word formation (Glass Mountains) at 44–46, 46, 51, 54, 144, T144a, 153, and T233. Upper Wolfcamp formation? (Sierra Diablo) at 460. Hess formation (Sierra Diablo) at Te. Bone Canyon Member (Sierra Diablo) at 484. Leonard formation (Baylor Mountains) at 503. Word formation (Sierra Diablo) at 506. Leonard formation (Shafter) just above brecciated zone, at Td. Limestone of Cerro Caballo (Las Delicias) at section 1, beds 11 and 13–17, and section 2, between El Tordillo and Cerro Caballo on bed 12.

### COMPOSITA PERSINUATA (Meek)

Pl. XLIII, Figs. 18–19

1877 *Athyris? persinuata.* Meek, U. S. Geol. Expl. 40th Par. Final Rept., vol. 4, pt. 1, pp. 81–82, pl. 9, figs. 4–4b.

Light-colored Carboniferous limestone of White Pine Mountains, Nevada.

*Description:* Shell rhomboid, strongly biconvex, somewhat longer than wide, lateral margins rounded, shell margin deeply sinuous. Pedicle valve strongly convex in the umbonal region, depressed in front into a narrow, rounded median sinus, which is not very deep but projects forward and upward at the front as a long tongue; the sinus originates in the anterior part of the shell. Brachial valve about as deep as the other, elevated at the front in a narrow, high, rounded median fold, which originates a little in front of the middle of the shell and extends forward somewhat at the front margin, where it meets the tongue of the other valve at less than a right angle. Surface marked by concentric imbrications, which are few and not very strongly developed. The specimen from loc. 55 is exceptionally narrow, as shown by the following table giving in mm. the dimensions of several specimens:

| | 1 (loc. 241) | 2 (loc. 155) | 3 (loc. 55) | 4 (Meek's specimens) |
|---|---|---|---|---|
| Length | 22 | 22 | 16 | 32 |
| Width | 20 | 21 | 11 | 39 |
| Thickness | 15 | 15 | 10 | 24.5 |

*Observations:* *C. persinuata* is distinguished from *C. mira*, which occurs at the same horizons by being smaller, more elongated, more strongly convex, and having a stronger fold and sinus. The main difference from *C. subtilita* is its much narrower fold and sinus. The only difference between the Glass Mountains specimens and the type from Nevada is the much smaller size of the former. The good agreement of the other characters indicates that the size is a feature of lesser importance.

*Distribution:* Word formation (Glass Mountains) at 55, 155, T239, and T251. The species is not common at any locality.

### COMPOSITA SUBTILITA (Hall)

Pl. XLIII, Fig. 20; Pl. XLIV, Figs. 1–2

1852 *Terebratula subtilita.* Hall, Stansbury's Exped. Great Salt Lake, Rept., p. 409, pl. 4, figs. 1–2b.

1909 *Composita subtilita?* Girty, Manzano Group of the Rio Grande Valley, U. S. Geol. Sur. Bull. 389, pp. 68–72.

Yeso formation, Mesa del Yeso; San Andres formation, Fra Cristobal; Abo sandstone, Abo Canyon.

1914 *Seminula argentea.* Kozlowski, Brachiopodes du Carb. Supér. de Bolivie, Ann. de Pal., vol. 9, pp. 79–82, text fig. 20, pl. 11, figs. 1–46. (See for best complete description and figures of this species.)

1915 *Composita subtilita.* Girty, Fauna of the Wewoka Formation, U. S. Geol. Sur. Bull. 544, pp. 96–101, pl. 12, figs. 4–4c; pl. 5, fig. 7; pl. 6, fig. 13. (See for synonymy.)

*Observations:* This species is common at many localities in the Wolfcamp, Hess, and Leonard formations. In the Gym formation it is represented by small specimens which are associated in places with *C. mexicana* but have the fold and sinus less strongly or even very faintly developed. Since the only difference from *C. mexicana* is the faint development of the sinus, it may perhaps be proper to regard the Gym types as a variety intermediate between typical *C. subtilita* and *C. mexicana*, an idea borne out by the occurrence there of *C. subtilita* only in the lower part of the formation, while *C. mexicana* is most common in the upper part. The upper Hess formation of the Glass Mountains contains an abundance of *C. subtilita* which are entirely similar both externally and internally to average specimens of *C. subtilita*. In the Wolfcamp formation at loc. 456 there are two exceptionally large specimens, which are somewhat deformed.

*Distribution:* Wolfcamp formation, *Uddenites* zone (Glass Mountains) at 88, 201, 202, and 203. Wolfcamp formation (Glass Mountains) at 89, 90, 92, 93, and 196. Hess formation (Glass Mountains) at 4, 100?, 104, 106, 107 (abundant), T107, 178?, 207, 208, 211, and 223. Leonard formation (Glass Mountains) at 3, 7, 8, 8–14, T9, 12, 13, 15, 16, 19, 20, 23, 24, 26s,* 31, 35, 37, 38, 82, 83, 119, 120, 232, Tba, D2, D3, and D20. Hess formation (Sierra Diablo) at 469, 474, 479, and Te. Bone Canyon member (Sierra Diablo) at 483. Victorio Peak member (Sierra Diablo) at 494. Leonard formation (Baylor Mountains) at 502. Gym formation (Wylie Mountains) at 507. Gym formation (Carrizo Mountains) at 510. Lower Gym formation (Hueco Mountains) at 389, 395 (abundant), 401, 402, 404, 405, 407, and 409. Leonard formation (Shafter) at 514, above transition beds at Ti, Cieneguita beds at Tg.

### COMPOSITA SUBTILITA ANGUSTA King, n. var.

Pl. XLIV, Figs. 9–10

*Description:* Under this designation are included some shells of variable size but similar configuration. The specimens from the *Uddenites* zone, which were taken as the types, are rather small, but those from loc. 456 are larger, and two of them are very large shells for the genus, but both these latter were crushed and distorted so that their reference to this species is very uncertain. Shell considerably longer than wide, subpentagonal, with the greatest width at about mid-length. Pedicle valve evenly rather strongly convex longitudinally; beak broad and incurved. The transverse profile is regularly convex, or slightly depressed in the median part near the anterior margin which is prolonged forward as a tongue. There can hardly be said to be a true sinus. Brachial valve strongly and evenly convex, elevated at the front to correspond to the tongue of the other valve. The surface is smooth except for low varices of growth near the margins and faint, irregular radial marks which occur on some more or less exfoliated specimens, and therefore probably do not have the nature of surface markings. Of the internal characters only the dental lamellae of the pedicle valve were made out.

Dimensions in mm. of several specimens:

*Identified by Schuchert (Am. Jour. Sci., ser. 5, vol. 14, p. 395, 1927) as *Notothyris schucherti*.

| | 1 (loc. 199) | 2 (loc. 199) | 3 (loc. 199) | 4 (loc. 456) |
|---|---|---|---|---|
| Length | 17.5 | 16 | 13 | 25 |
| Width | 13 | 13 | 11.5 | 19 |
| Thickness | 10 | — | 7 | 14.5 |

*Observations:* The specimens referred to this variety agree in being narrow, rather strongly convex, and with an extended anterior tongue but no true median sinus. These features distinguish it especially from *C. subtilita.* It is quite different from *C. mexicana,* which is much more deeply sinuate and generally smaller in size. Some of the lower Gym specimens referred to *C. subtilita* show a certain general resemblance to *C. subtilita angusta* in the amount of development of the fold and sinus, etc., but are relatively considerably broader, and thus much closer to if not identical with *C. subtilita.*

*Distribution:* Wolfcamp formation, *Uddenites* zone (Glass Mountains) at 95, 168, 199 (types), and 201. Wolfcamp formation (Glass Mountains) at T93? and 193. Upper Wolfcamp formation (Sierra Diablo) at 456 (abundant, but generally crushed). Leonard formation (Glass Mountains) at 124?.

## Family TEREBRATULIDAE

### Genus DIELASMA W. King

*Diagnosis:* Shell terebratuliform. Pedicle valve with or without a median sinus, and with well developed dental lamellae internally, foramen large and encroaching on the umbonal part of the valve, beak strongly incurved. Brachial valve generally without median fold; internally there may be crural plates which are separate from the dental socket plates, or the crura may be supported on the septal plates; the free part of the brachidium is short with diverging descending lamellae. In the inner posterior part of the valve there is a concave, transverse plate for muscular attachment, which joins the inner surface of the crural plates or the dental socket plates a little above their bases, resting against the inner surface of the valve along the median line for a part or the whole of its length, or free throughout; where attached along the median line a pair of slender cavities, triangular in cross-section, converge from the general cavity of the shell toward the beak, but where, on the other hand, the transverse plate is not attached along its median line, there is a single, broad, low cavity beneath the plate extending toward the apex; the plate extends forward to a greater or less distance beyond the attachment of the crural plates and is pointed, rounded, or emarginate in front, its surface marked by concentric wrinkles parallel with its anterior margin which are generally discontinuous along the median line.

Genotype: *Terebratula elongatus* Schlotheim.

*Geologic range:* Devonian to Permian.

*Observations:* Weller's diagnosis of the genus, from which the above has been modified, is too specific to include some of the varieties of internal structures of the group. As there is some question as to the structures in the genotype it is thought best to change the diagnosis enough to admit such species as *D. problematicum wordensis.*

Weller* states that "the crural plates are separated from the socket walls, one of the most essential features of *Dielasma* as here defined." In *P. biplex* the crural plates are undeveloped, the crura being supported on the septal plates. This seems to be true also of *D. bovidens* as figured by Kozlowski, and of *D. breviplicatum* Waagen. Waagen believed that in the last named species there was also a median dorsal septum; if his observation was correct, it is an important difference from most other species of the genus. The mode of junction of the septal plates with the roof of the valve is quite different from that in both *D. breviplicatum* and *D. bovidens,* in which the septal or muscle-bearing plates join one another in the median line but do not meet the roof of the valve.

### DIELASMA PROBLEMATICUM WORDENSE King, n. var.

Pl. XLIV, Fig. 16

1915 *Dielasma* cf. *biplex.* Haack, Permfauna aus Nordmexiko, Zeit. der deutsch geol. Gesell., bd. 66 (1914), p. 502, pl. 39, figs. 12a–b.

Pichagua, near Las Delicias.

*Description:* Shell medium-sized, longer than wide, subpentagonal, widest at about mid-length. Pedicle valve of low, even longitudinal curvature. The transverse profile in the posterior part of the shell is flattened or somewhat depressed in the middle, then bent dorsally at an acute angle to meet the margin of the other valve without any angulation. At about mid-length the angulations at the sides merge into the lateral margin, the entire transverse profile of the shell is slightly concave, and the two valves meet at an acute angle. At the front the shell curves upward toward the fold of the other valve. There is a low median angulation corresponding to the median sulcus of the fold of the brachial valve. Brachial valve of low longitudinal convexity but strongly bent transversely and much deeper than the pedicle valve. The median part of the valve is a high crest from which the shell falls away steeply on each side. At about mid-length there originate two rounded folds, which increase in size toward the front; between them is a median sulcus.

Dental lamellae of the pedicle valve short and strong. Two long dental sockets originate on each side of the cardinal process; the socket plates diverge from the apex and are attached to the roof of the valve along the cardinal margins. Joined to the proximal sides of the socket plates are septal plates, which descend obliquely toward the median line of the valve, but in the posterior part join before they reach the roof of the valve, while farther forward they become fixed to the roof but gradually diverge from each other so that they no longer meet. On the dorsal faces of the septal plates are the upper parts of the crura, which originate at the cardinal process and extend across the septal plates, projecting from them toward the front. There are no crural plates supporting them; they rest solely on the septal plates. The crura descend nearly straight to about mid-length of the shell, diverging somewhat. The crosspiece is not preserved in the specimen studied.

*Mississippian Brachiopoda, Ill. Geol. Sur. Mon. 1, pp. 256–257.

Dimensions in mm. of one specimen from T144: length, 19; width, 12; thickness, 9.5.

*Observations:* This variety is remarkable for having the two plications on each side of the fold of the brachial valve. It differs from typical *D. problematicum* (Davidson) in having its greatest width near rather than posterior to mid-length.

*Distribution:* Word formation (Glass Mountains) at T144 (one specimen). Limestone of Pichagüia (Las Delicias) at Cerro Pichagüia (two specimens).

#### DIELASMA BOVIDENS (Morton)

Pl. XLIV, Fig. 11

1836 *Terebratula bovidens.* Morton, Am. Jour. Sci., (1), vol. 29, p. 150, pl. 2, fig. 4.

1903 *Dialasma bovidens.* Girty, Carb. Formations and Faunas of Colorado, U. S. Geol. Sur. Prof, Paper 16, pp. 409–411, pl. 7, figs. 11–11a. (See for synonymy.)

1914 *Dielasma bovidens.* Kozlowski, Brachiopodes du Carb. Supér. de Bolivie, Ann. de Pal., vol. 9, pp. 88–90, text fig 24, pl. 9, figs. 61–65. (See for best description.)

*Distribution:* Wolfcamp formation, *Uddenites* zone (Glass Mountains) at 95, 199, 201, and 203. Wolfcamp formation (Glass Mountains) at 91a. Lower Gym formation (Hueco Mountains) at 395 and 397.

#### DIELASMA SHAFTERENSE King, n. sp.

Pl. XLIV, Fig. 21

*Description:* Shell large, spatulate, longer than wide, with the greatest width at about mid-length. In the anterior half of the shell the outline is semicircular; the posterior half has nearly straight sides which converge to the beak at an angle. The longitudinal convexity of the pedicle valve is low and even, being greatest at the beak and decreasing toward the front. Almost at the beak there originates a slight median depression which broadens and deepens toward the front, where it occupies the entire width of the shell, so that the transverse profile is strongly and evenly concave there. Brachial valve almost straight in longitudinal profile but strongly convex transversely, with the median part elevated; beak broad and extending back below the ventral umbo. Surface of both valves smooth; shell finely punctate.

Within the pedicle valve there are strong diverging dental lamellae. In the brachial valve there is a median muscle-bearing plate which is attached along the median posterior part of the shell and joins the inner surfaces of the crural plates a little above their bases.

Length, 29 mm.; width, 24 mm.; thickness, 13 mm.

*Observations:* This species is distinguished from *D. bovidens* by having the greatest width near mid-length rather than near the front, in having rounded rather than straight lateral margins in the pedicle valve (except in the posterior part), and in being proportionately much broader. It has the median sinus developed much farther from the beak than in *D. spatulatum*, and is relatively much broader than *D. prolongatum* Girty. The closest resemblance is with some Uralian species, such as *D. giganteum* Tschernyschew, which is, however, a much larger shell with a less deep sinus.

*Distribution:* Leonard formation, Cieneguita beds (Shafter) at Tf (no. 15953).*

#### DIELASMA SPATULATUM Girty

Pl. XLIV, Figs. 17–20

1859 *Terebratula elongata.* Shumard, Trans. Acad. Sci. St. Louis, vol. 1, p. 392 (date of volume, 1860). (Permian): Guadalupe Mountains.

1908 *Dielasma spatulatum.* Girty, The Guadalupian Fauna, U. S. Geol. Sur. Prof. Paper 58, pp. 330–331, pl. 16, figs. 3–4c.
Top of Capitan, Capitan Peak (sta. 2966?); Middle Capitan, Capitan Peak (sta. 2926); dark limestone, Pine Spring (sta. 2930), Guadalupe Point (sta. 3762d?); Delaware Mountain formation, southern Delaware Mountains (sta. 2962?).

1908 *Dielasma cordatum.* Girty, The Guadalupian Fauna, U. S. Geol. Sur. Prof. Paper 58, pp. 331–332, pl. 16, figs. 2–2c.
Middle Capitan, Capitan Peak (sta. 2926).

*Description:* Shell medium-sized, spatulate, longer than wide, with the greatest width toward the front. Pedicle valve not very convex, strongly bent upward at the lateral margins. Beak large, projecting over that of the other valve. At about mid-length of the shell a slight concavity appears along the median line, which broadens but remains shallow and ill-defined, and does not at any point occupy most of the width of the shell; the sinus causes some upward deflection of the front margin. Brachial valve of very low convexity, with a gentle, even, transverse arching except for a downward deflection of the shell at the lateral margins. Beak small and not incurved. There is a low elevation at the front corresponding to the sinus of the other valve. Surface smooth except for fine concentric growth varices, which are most closely crowded toward the margins. Shell finely punctate. Within the pedicle valve there are strong dental lamellae.

Dimensions in mm. of several specimens:

| | 1 (loc. 144) | 2 (loc. T240) | 3 (loc. T240) | 4 (type *D. spat.)* | 5 (type *D. cor.)* |
|---|---|---|---|---|---|
| Length | 21 | 18 | 16 | 20 | 19 |
| Width | 17 | 15 | 13.5 | 16.5 | 14.5 |
| Thickness | 10 | 10 | 9.5 | 10 | 8.5 |

*Observations:* Girty stated that in *D. cordatum* the median sinus was narrower than in *D. spatulatum*. The specimens in the present collection are more or less intermediate, and could as well be placed in one or the other of Girty's species. The good agreement of the two in other respects, as well as their occurrence together, makes their identity hardly doubtful. As Girty has noted, *D. spatulatum* is very similar to *D. truncatum* Waagen.

*Distribution:* Leonard formation (Glass Mountains) at Ax?. Word formation (Glass Mountains) at 46, 144, 148, 171, 239, T239, T240, 243, 255, and Tc. Word formation, yellow limestone (Shafter) at Tp. (no. 16013).*

*Identified as *Dielasma bovidens* in Geol. Shafter Silver Mines Dist., U. of Texas Bull. 24, p. 14.

*Probably the form identified as *Seminula argentea* Shepard? in the "Geol. of the Shafter Silver Mine Dist." U. of Tex. Bull. 24, p. 23.

**DIELASMA? sp.**

This includes a heretogeneous group of terebratulids, most of which are almost certainly, and the others probably *Dielasma.* Those from T233 and Taa are broad and of low convexity, those from 241 are large and thick, and those from locs. 8–14, D1, and D3 are narrow and deep.

*Distribution:* Leonard formation (Glass Mountains) at 8–14, 12, T233, Ax, D1, and D3. Word formation (Glass Mountains) at 241. Victorio Peak member (Brackett Draw) at Taa.

**DIELASMA SCHUCHERTI MINOR King, n. var.**

Pl. XLIV, Figs. 12–14

*Description:* Shell small, outline longitudinally subelliptical, tapering toward the beak. Pedicle valve of low, even convexity, beak broad and incurved, pierced by a round foramen. The median anterior part is somewhat prolonged forward and upward as a tongue, and is marked by a low median plication and two lateral ones, which are developed only at the very front. Brachial valve more strongly arched than the other, somewhat elevated near the front and marked by two plications corresponding to the sulci between the plications of the other valve. Surface smooth, shell finely punctate. The internal characters are indeterminable, though in some specimens dental lamellae seem to be distinguishable. Length of a specimen from loc. 239: 8 mm.; width, 6.5 mm.; thickness, 4 mm.

*Observations:* A comparison of these shells with actual specimens of *Dielasma schucherti** from the Whitehorse sandstone shows that the only difference between them is the smaller size of the Word specimens, average specimens of the latter being only about half as large as those from Oklahoma.

*D. schucherti* and its variety are very similar to *D. problematicum wordense,* but are relatively much broader and of lower convexity. In general outline they resemble *Notothyris schucherti* Girty, of the Delaware Mountain and Capitan, but have a distinct fold and sinus as in typical *Dielasma,* and have one less plication in the anterior part of the shell than does Girty's species. The generic reference of *Notothyris schucherti* is very doubtful. The genus *Notothyris* is based on internal characters, which were not determined by Girty for his species. Moreover, the external resemblance of *N. schucherti* to *Dielasmina guadalupensis* is so great that it hardly appears to be more than a variety of that species.

Beede's species is very close to *Dielasma biplex* Waagen, but is easily distinguishable by minor differences in configuration. *Dielasma schucherti minor* is still further distinguished from Waagens' species by its much smaller size.

*Distribution:* Middle Word formation (Glass Mountains) at 144, 239 (abundant), and D239.

**Genus DIELASMINA Waagen**

*Diagnosis:* Similar to *Dielasma,* but instead of having a well marked fold and sinus, folded into a series of subequal plications limited to the marginal part of the shell, without any distinct fold or sinus.

Genotype: *Dielasmina plicata* Waagen.

*Geologic range:* Permian.

**DIELASMINA GUADALUPENSIS Girty**

Pl. XLIV, Fig. 15

1908 *Dielasmina guadalupensis.* Girty, The Guadalupian Fauna, U. S. Geol. Sur. Prof. Paper 58, pp. 333–334, pl. 16, figs. 6–7a; pl. 21, figs. 22–22a.
Capitan formation, Capitan Peak (sta. 2926); dark limestone, Pine Spring (sta. 2930).

1915 *Dielasmina Guadalupensis.* Haack, Permfauna aus Nordmexiko, Zeit. der deutsch. geol. Gesell. bd. 66 (1914), p. 501, pl. 39, figs. 11a–c.
Pichagua, near Las Delicias.

*Description:* Shell small, inflated, longer than wide, with the greatest width at mid-length. Pedicle valve strongly and evenly convex longitudinally, with a small incurved beak. The transverse profile is evenly of low convexity except for two narrow, rounded plications which originate at about mid-length, and two indistinct marginal folds. Brachial valve of low, even convexity, in the anterior part folded into 3 strong, rounded plications. Surface smooth. Within the pedicle valve are two strong, short dental lamellae. Dimensions in mm. of the only specimen: length, 13; width, 10; thickness, 8.

*Observations:* The specimen in the present collection is unlike Girty's type in being smaller and having the plications well developed at a size when the other is smooth or only faintly plicated. The differences do not, however, appear to justify a distinct name. The differences between *D. guadalupensis* and *D. perinflata* (Shumard) are so small that it hardly seems proper to regard them as distinct. If not, the latter name would be used, as it is the older. *D. guadalupensis* and *D. perinflata* are less strongly plicated than most Eurasian species of the genus.

*Distribution:* Upper Hess formation (Glass Mountains) at 223.

INDEX TO LOCALITIES*

1 Hess: Small hill 1 mile northwest of the summit of Iron Mountain.
2 Wolfcamp, upper: Section 14, shale, west of Iron Mountain.
3 Leonard: 1 mile west of the summit of Iron Mountain, about 50 feet above the base.
4 Hess?: .6 mile north of hill 5021, Altuda Quadrangle.
5 Leonard: Section 14, below soft shale of the Clay slide, about 250 feet below the top. Same as 301?
6 Word, lower: near Clay slide, section 14.
7 Leonard: Section 14, bed 16.
8 Hess: Section 14 (contains some lower Leonard fossils also).
9 Leonard: Section 14, below middle limestone layer of bed 9.
10 Leonard: Section 14, middle limestone layer of bed 9.
11 Leonard: Section 14, between middle limestone layer of bed 9, and top of bed 12.
12 Leonard: Section 14, top of bed 12.
13 Leonard: Section 14, base of bed 14.
14 Leonard: Section 14, lower part of bed 15.

*Beede, J. W.: Invertebrate Paleontology of the Upper Permian Red Beds of Oklahoma and the Panhandle of Texas, Kans. Univ. Science Bull., vol. 4, 1907, pp. 148–150.

*References to bed and section numbers in the Glass Mountains are to stratigraphic sections described in Part One of this memoir. Collections from these localities were made by R. E. King and P. B. King.

15 Leonard: Section 15.
16 Leonard: Section 15, slightly above the base, or near the base.
17 Hess: Projecting spur of range just west of the Marathon-Sullivan Ranch road; the lower part, 17a, the middle 17b, the upper 17c.
18 Hess: Section 12.
19 Leonard: Basal, section 12, bed 1.
20 Leonard: Section 12, bed 1.
21 Leonard: Section 12, near middle of bed 24.
22 Leonard: Section 12, middle of bed 24.
23 Leonard: Section 12, near middle of bed 24, below hill 4920, and north of locality 21.
24 Leonard: Section 12, some distance above middle of bed 24, but below top of hill 4920.
25 Leonard: Section 12, upper part of bed 24, southwest side of hill 4920.
26 Leonard: Section 12, bed 26, southwest side of hill 4920.
27 Leonard: Section 12, beds 19 to 24, southwest side of hill 4920.
28 Leonard: Section 12, middle of bed 24, southwest side of hill 4920.
30 Leonard: Section 12, bed 28, south of Sullivan Peak.
31 Leonard: Section 12, bed 34, south of Sullivan Peak.
31a Leonard: Section 12, bed 36, south of Sullivan Peak.
32 Word, lower: South of Cathedral Mountain.
34 Leonard: .5 mile south of tank at locality 4555.
35 Hess: .5 mile east-northeast of hill 4902, southern Altuda quadrangle, at base of escarpment.
36 Leonard: Section 11, bed 4, on hill 5300.
37 Leonard: Section 15, bed 10, tank west of north end of Iron Mountain.
38 Leonard: Basal, .25 mile east of hill 5300.
39 Hess: Section 10, top of bed 1.
40 Leonard, upper: Southeast of James Ranch.
41 Capitan, Altuda member: East side of Altuda Mountain.
42 Capitan, Altuda member: Valley northeast of Altuda.
43 Capitan, Altuda member: Top of Cathedral Mountain, east end.
44 Word, lower limestone: Section 12, south of Sullivan Peak.
45 Word, lower limestone: Section 12, .40 mile southwest of 44.
46 Word, lower limestone: Section 12, .25 mile south of 44 and .2 mile east of 45.
47 Word, middle: West side of Gilliland Canyon, below hill 5939.
48 Word, middle limestone: .5 mile northeast of locality 4869, north-northwest of Iron Mountain.
49 Capitan: .5 mile north-northwest of point 6311, north of Cathedral Mountain.
50 Capitan: Altuda member, northeast part of Del Norte range south of James Ranch.
51 Word: Lower limestone, northeast of Clay slide.
52 Leonard: Section 12, bed 37, lower part; calcareous sandstone, bearing ammonoids, south of Sullivan Peak.
53 Word: Lower, section 11, beds 1 and 2, 1.5 mile southwest of Sullivan Peak.
54 Word: Lower, section 11, bed 3, 1.5 mile southwest of Sullivan Peak.
55 Word: Lower, 1.1 mile south-southwest of hill 5935, at about the same horizon as locality 54.
56 Capitan: .4 mile north-northwest of hill 5935, west end of Cathedral Mountain.
57 Capitan: Altuda member, 1.5 mile northeast of Altuda.
58 Word: Lower, south side of Cathedral Mountain.
59 Word: Upper, section 10, beds 8 and 9.
60 Gilliam: .7 mile east of hill 4870 south of Gilliland Canyon.
61 Capitan: Altuda member, .25 mile east of hill 5672, at southwest end of Glass Mountains.
62 Word: Upper, from loose blocks lying on downslumped Vidrio 7 miles southeast of Altuda.
63 Word: Upper, east slope of Sullivan Peak.
69 Wolfcamp: Lower shale, just to north of igneous plug, Hess Ranch horst.
70 Wolfcamp: Base of range about 1 mile west of Sullivan Ranch-Marathon road.
71 Wolfcamp: At base of escarpment between Sullivan Ranch-Marathon road and a point .5 mile to west.
72 Wolfcamp: At base of escarpment about .3 mile west of Sullivan Ranch-Marathon road.
73 Wolfcamp: Hill just west of Sullivan Ranch-Marathon road.
74 Wolfcamp: Between hill west of Iron Mountain and the Sullivan Ranch-Marathon road.
75 Wolfcamp: .75 mile west-northwest of the summit of Iron Mountain.
76 Wolfcamp: Upper, westernmost outcrop of Wolfcamp on Leonard Mountain; bed containing large *Schwagerina.*
77 Wolfcamp: Upper, lowest beds above basal conglomerate, about .2 mile east of locality 76.
78 Leonard: Float on south side of Leonard Mountain.
79 Wolfcamp: Northwest of the summit of Iron Mountain, .2 mile south of locality 1.
82 Wolfcamp or Leonard: Float on middle of south side of Leonard Mountain.
83 Leonard: Float at east base of Leonard Mountain.
84 Wolfcamp: *Uddenites* zone on northeast side of Leonard Mountain, near small igneous intrusion.
85 Wolfcamp: South middle part of Hess Ranch horst, west end.
86 Wolfcamp: South side of Hess Ranch horst, west end.
87 Wolfcamp: One mile west by north of Wolfcamp, Section 23, bed 12 of Wolfcamp section and above. Differentiated into two parts, low and high, the lower part being approximately equivalent to bed 13 and the upper to the higher beds of the section at Wolf Camp.
88 Wolfcamp: Section 24, *Uddenites* member, bed 1b at Wolf Camp.
89 Wolfcamp: Section 24, bed 4, at Wolf Camp; containing a few brachiopods.
90 Wolfcamp: Section 24, bed 8, at Wolf Camp; not very fossiliferous.
91 Wolfcamp: Section 24, bed 12, at Wolf Camp.
91x Wolfcamp, Hess, or Leonard: Loose pieces picked up on bed 12, section 24, the fossils of which indicate a Hess or Leonard age.
92 Wolfcamp: Section 24, bed 14, at Wolf Camp.
93 Wolfcamp: Section 24, bed 9, and float from the next few higher beds on the side of the arroyo northeast of Wolf Camp.
94 Wolfcamp: *Uddenites* bed, about .5 mile west of locality 95, below hill 4815, and northeast of Wolf Camp.
95 Wolfcamp: Section 27, bed 13, at hill 4752; *Uddenites* bed.
96 Wolfcamp: Lowest beds near Gap Tank, about .25 mile east of the edge of the Hess Canyon quadrangle.
91 Wolfcamp: Basal beds about .25 mile southeast of Gap Tank.
98 Hess: West end of Glass Mountains in Monument Spring quadrangle, about 20 feet above the Wolfcamp.
99 Hess: East end of Dugout Mountain, along the downfaulted spur.
100 Hess: West end of Dugout Mountain.
101 Hess: Upper, top of Leonard Mountain, in the line of section 17.
102 Hess: Near top of formation, section 17.
103 Hess or Leonard: Float on Wolfcamp at foot of escarpment about 1 mile west of the Marathon-Sullivan Ranch road.

104 Leonard: Northeast of Hess Ranch, on north side of the escarpment, 1.2 mile west of hill 5725.
105 Hess: .5 mile northeast of hill 5305 on west end of the Hess Ranch horst.
106 Hess: On hill 5305 from beds lower than those from which collection was made at locality 105.
107 Hess: Upper, on escarpment west of Hess Canyon fault, from upper fossiliferous horizon, immediately below a conspicuous layer of massive limestone; *Perrinites compressus* horizon.
108 Hess: Upper, same horizon as 107, between and south of hills 5767 and 5821.
109 Hess: Upper, float from the beds above the upper fossiliferous horizon, collected upon the fossiliferous bed.
110 Hess: Lower, bed 5, section 23, escarpment northwest of Wolf Camp.
111 Hess: Uppermost, south of the road and a short distance west of the Word Ranch.
112 Hess: Upper, about .5 mile south of the forks of Hess Canyon, 1.25 miles southwest of Word Ranch.
113 Hess: Middle, section 27, bed 6, on hill 4752, near top of layer of nodular limestone.
114 Hess: Middle, about 1 mile southeast of hill 5233, eastern Hess Canyon quadrangle.
115 Hess: Middle, .5 mile northwest of hill 4762 on escarpment, eastern Hess Canyon quadrangle.
117 Hess: Upper, between hills 5233 and 5035, on the east side of the long valley, eastern Hess Canyon quadrangle.
118 Hess: Upper, .5 mile northeast of hill 5233, at elevation 5000, eastern Hess Canyon quadrangle.
119 Leonard: Upper, below and to west of Clay Slide.
120 Leonard: Middle, .6 mile east of hill 4910, east of Clay Slide; lower *Perrinites* horizon.
121 Leonard: Middle, section 17, north of Leonard Mountain, bed 13 near top.
122 Hess or Leonard: Limestones below lower Leonard shale on Leonard Mountain, section 17.
123 Leonard: Section 17, bed 14; richly fossiliferous horizon near middle of Leonard formation.
124 Leonard: Lower, hill south of forks of Hess Canyon, south of road to Word Ranch.
125 Leonard: Upper, about .5 mile south of hill 5611, south of Hess Canyon.
126 Leonard: Above conglomerate bed near the Word Ranch house.
127 Leonard: Directly above conglomerate bed southwest of Split Tank, 1.5 mile northeast of Word Ranch house.
128 Leonard: .2 to .3 mile northeast of Split Tank, 1.5 mile northeast of Word Ranch house; also from the same horizon about a mile east-northeast of Split Tank; the Tank is on the line of section 26.
129 Leonard: Upper, north of hill 4627, northeast corner of Hess Canyon quadrangle.
130 Word: Middle, Del Norte Mountains, near the "I" in "Mountains" as represented on the topographic sheet, Monument Spring quadrangle; yellow limestone between lower and middle limestones.
131 Word: Lower limestone, south side of Altuda Mountain.
132 Word: First limestone, section 17, bed 1.
133 Word: First limestone, section 17, upper part of bed 1.
134 Word: Lower, section 17, bed 3.
135 Word: Third limestone, section 17, bed 7c, mountain north of Leonard Mountain.
136 Word: Third limestone, Section 17, bed 7f, mountain north of Leonard Mountain.
137 Word: Third limestone, Section 17, mountain north of Leonard Mountain.
138 Word: Third limestone, Section 17, top of hill south of junction of Road and Gilliland Canyons.
139 Word: Float at junction of Road and Gilliland canyons.
140 Word: First limestone, .5 mile south of hill 5874, west side of Gilliland Canyon.
141 Word: Lower limestone, above Clay slide, section 14; same locality as number 6.
142 Word: Fourth limestone, about .5 mile northeast of Willis Ranch.
143 Word: Fourth limestone, east and west of hill 5578, Hess Canyon.
144 Word: Second limestone, .5 mile north of hill 5611, Hess Canyon.
145 Word: Second limestone, 1.5 mile northwest of Word Ranch house, .5 mile west of hill 5507, Hess Canyon.
146 Word: Third limestone, south and west of south part of Comanche outlier north of Word Ranch house.
147 Word: First limestone, in channel of Hess Canyon near Leonard-Word contact.
148 Word: Fourth limestone, northeast of Word Ranch house.
149 Word: Fourth limestone, northeast of Word Ranch and east of 148.
150 Word: Fourth limestone, east of Comanche outlier north of Word Ranch house; collections extend for about a mile along the strike.
151 Leonard: Near Split Tank, collected by McGonagill family, same as 128.
152 Word: Upper, several miles east-northeast of Word Ranch house.
153 Word: First limestone, south side of hill 5611, west side of Hess Canyon, section 23.
154 Word: Upper, east from hill 5360 to big fault 5 miles northeast of Word Ranch.
155 Word: Upper, east of big fault 5 miles northeast of Word Ranch, as far as hill 4902; mainly collected from west side of latter hill. Large flat gastropods present at this locality could not be collected.
156 Word: Upper, across valley from hill 5461; Hess Canyon quadrangle.
157 Word: Upper, large *Fusulina* from east side of big fault.
158 Word: Upper, chert on hill 4732, Hess Canyon quadrangle.
159 Word: Upper, chert east of hill 4800, northeast part of Hess Canyon quadrangle.
160 Delaware Mountain formation: West of Guadalupe Point in hills just to north of highway from Van Horn to Carlsbad.
161 Capitan: Altuda member, south side of Altuda Mountain.
162 Word: Upper, cherty limestone east of Comanchean outcrop north of Word Ranch, approximately the same as 150.
163 Capitan: Middle of Altuda member, bed 5, section 16, at south end of Old Blue Mountain.
164 Capitan: Gilliam member; lower 300 feet in Gilliland Canyon just north of Igneous plug.
165 Capitan: Gilliam member; upper part 1 mile east-southeast of B. M. 3948, and .75 mile northwest of hill 4870.
168 Wolfcamp: Section 24, Gray Limestone member at Wolf Camp.
169 Wolfcamp: About 250 yards south of locality 84, the ammonoid locality on the east side of Leonard Mountain.
169a Wolfcamp: Just above Gray Limestone member, section 24, Wolf Camp.
170 Word: Upper, thick prominent limestone 700 feet below top north of Dugout Mountain.
171 Word: Lower limestone, northwest of Dugout Mountain.
173 Word: Lower limestone, capping hill between B. M. 4873 and 4869, in Gilliland Canyon.
174 Leonard: Section 27, .25 mile east of hill 5157, in arroyo.
175 Wolfcamp: Below Gray Limestone member, northeast of locality 88.

175a Wolfcamp: Yellow-brown limestone in shale underlying Gray Limestone member northeast of locality 88.
176 Word: Middle, west side of Gilliland Canyon, west of Road Canyon.
179 Capitan: Gilliam member, 2.1 miles east of Easterwood Ranch, on top of ridge.
180 Word: Float in northern Hess Canyon.
181 Wolfcamp: Range east of Wolf Camp hills, and southeast of hill 4815.
182 Wolfcamp: Wolf Camp section, section 24, shale of bed 5.
183 Leonard: Section south of Altuda Mountain, section 5, bed 8.
184 Capitan: Gilliam member?; south of old Beatty Place, from a thick bed crowded with *Fusulina.*
185 Capitan: Gilliam member; .75 mile north of head of Spring Canyon.
186 Capitan: Altuda member; northern Altuda Mountain, 1.5 mile north of locality 41.
190 Wolfcamp: Northeast of Gap Tank, from shale containing *Schwagerina,* directly below Hess conglomerate.
191 Hess: Northeast of hill 4402, Van Horn quadrangle.
192 Word: Fourth limestone; north of junction of Road and Gilliland Canyons, section 17.
193 Wolfcamp: Upper, south of high point on Dugout Mountain.
194 Wolfcamp: Upper, perhaps including some basal Hess, from float on south side of high point of Dugout Mountain.
195 Wolfcamp: Lower shale; .5 mile southwest of hill 5816, Hess Ranch horst.
196 Wolfcamp: Upper; .5 mile northeast of hill 5305, Hess Ranch horst.
198 Wolfcamp: Upper fossiliferous bed in the graben near the middle of the Hess Ranch horst, and the same bed to the west of the graben.
199 Wolfcamp: *Uddenites* member; .35 mile northeast of hill 5060, east of Wolf Camp; basal brown limestone about 15 feet above Gaptank limestone.
200 Wolfcamp: Gray Limestone member, lower part, .35 mile southwest of hill 4952, northeast of Wolf Camp.
201 Wolfcamp: Brown limestone near base of hill, .3 mile southwest of hill 4815, 4 miles northeast of Wolf Camp.
202 Wolfcamp: Directly below base of Hess, .8 mile west of east end of Hess Canyon quadrangle.
203 Wolfcamp: Beds below Gray Limestone member, on range of foothills at east edge of Hess Canyon sheet, within the quadrangle.
204 Wolfcamp: .5 mile south of Allison and Gilbert ranch.
205 Hess: At foot of cliff of Dugout Mountain, 1 mile southeast of summit.
206 Hess: Uppermost, 1 mile northeast of Lenox.
207 Hess: Upper, west side of fault on spur north of high point of Leonard Mountain.
208 Hess: .5 mile north of Hess Ranch, at west end of limestone ridge.
209 Hess: 150–200 feet below top, northeast end of Leonard Mountain, directly below Leonard outlier on east side of fault, below and south of locality 207.
210 Hess: White limestone at top of section on Hess Ranch horst, north of the northeast end of igneous intrusion.
211 Hess: .75 mile north-northeast of hill 5305, on Hess Ranch horst.
212 Hess: Upper fossiliferous horizon, the same as that of localities 107 and 108, .6 mile west-southwest of hill 5725, northwest of Wolf Camp.
213 Hess: Middle part, from dolomite layer about halfway up scarp south of hill 5725, 3.5 miles east-northeast of Hess Ranch.
214 Hess: Section 26, on Meeks Ranch, southeast of hill 5567 on face of escarpment, beds 11, 12, and 13.
215 Hess: Upper fossiliferous horizon, .5 mile east-southeast of hill 5632, at top of section 27.
216 Hess: Middle part, 4.25 miles east-northeast of Gap Tank.
217 Hess: Middle part, about 4 miles west of east edge of Hess Canyon quadrangle; layer containing *Fusulina.*
218 Hess: Middle part, 1.5 mile east-northeast by east of Allison and Gilbert Ranch, along old washed-out road.
219 Hess: Basal layers at easternmost outcrops 3.5 miles east of Allison and Gilbert ranch, from layers of sandstone.
220 Hess: Upper part on cuesta capped by Hess limestone west of the fault bounding the west side of Dugout Mountain.
221 Leonard: Bone Canyon member, near contact with Delaware Mountain formation near Bone Spring, Guadalupe Mountains.
222 Hess: Upper fossil horizon, same as that of localities 107, 108, and 212, on scarp east of head of east fork of Hess Canyon about .25 mile below and southeast of hill 5767.
223 Hess: Uppermost, 1.1 mile southwest of Word Ranch house.
224 Leonard: First limestone member, or slightly below, at west end of Dugout Mountain.
225 Leonard: Just below base of Word at old Payne Ranch, from highest limestone layer of the Leonard.
226 Leonard: First Limestone member, in central part of Dugout Mountain.
227 Leonard: Second Limestone member on Dugout Mountain.
228 Leonard: Horizon of First Limestone, east of fault at east end of Dugout Mountain.
229 Leonard: Lower 200 feet, west end of range north of Lenox.
230 Leonard: First Limestone member, at top, 2 miles northeast of Lenox.
231 Leonard or Uppermost Hess: Limestone containing chert pebbles directly below Leonard shale on south face of Leonard Mountain; same as locality 122.
232 Leonard: Lower limestone on top of Leonard Mountain, above first bed of siliceous shale.
233 Leonard: Basal, on spur north of high point of Leonard Mountain, on east side of fault at east end of the Mountain.
234 Leonard: .6 mile west-northwest of hill 5726, south of Hess Ranch horst.
235 Leonard: Upper part, 1 mile west of Word Ranch house, section 24.
236 Leonard: Uppermost, .3 mile southeast of hill 5674, north of Leonard Mountain.
237 Word: Lower limestone near latitude 30°10′, Del Norte Mountains.
238 Word: Middle limestone, .5 mile south of south side of big basin in northern Del Norte Mountains.
239 Word: Third Limestone member, on south side of Road Canyon, west of divide separating the drainage of Gilliland and Hess canyons.
240 Word: Third Limestone member, capping mountain north of Leonard Mountain along the line of section 18, southeast of hill 5453.
241 Word: First limestone member on mountain north of Leonard Mountain, southeast of hill 5453.
242 Word: Third Limestone member, from bed below uppermost white limestone at east end of range north of Leonard Mountain, on the north slope of hill 5453.
243 Word: Third Limestone member, .5 mile southeast of hill 5803, north side of Road Canyon.
244 Word: Fourth Limestone member, north of Willis Ranch.

245 Leonard: Downfaulted block south of Hess Ranch horst, .6 mile west-northwest of hill 5726, from bituminous limestone; same as 234?
246 Word: Fourth Limestone member, north side of Hess Canyon, east of hill 5543.
247 Word: Fourth Limestone member, east side of Hess Canyon, near its angle.
248 Word: Fourth Limestone member, on east side of Hess Canyon, near northernmost outcrop of the formation.
249 Word: Basal dolomite, 1 mile west of Word Ranch.
250 Word: Second Limestone member, section 23, east of Hess Ranch horst; same as 264.
251 Word: First Limestone member, .8 mile west of Word Ranch on section 24.
252 Word: Fourth Limestone member, on line of section 24 west of Comanche outlier north of Word Ranch.
253 Word: Upper part of Fourth Limestone, north of Word Ranch, east and west of Comanche outlier for a short distance.
254 Word: Upper part of Fourth Limestone, 1 mile northeast of Comanche outlier north of Word Ranch.
256 Word or Leonard: Lower part of hill lying in the middle of a broad valley in the northeasternmost corner of the Hess Canyon quadrangle.
257 Word: Middle and upper beds on hill in middle of broad valley in the northeasternmost corner of Hess Canyon quadrangle.
258 Hess: Near summit of Sierra Madera.
259 Word: .35 mile southeast of hill 4108, north-central part of Sierra Madera.
260 Word: .6 mile west of hill 4054, east-central Sierra Madera.
261 Word: .5 mile west-northwest of point 3663, northeast of Sierra Madera; from float.
262 Word: .3 mile south of hill 4600, the high point of Sierra Madera.
263 Word: About .5 mile northeast of point 4588, Sierra Madera.
264 Word: Second Limestone member east of Hess Ranch horst, west side of Hess Canyon, same as 250.
265 Word: .8 mile northwest of hill 5079, northwestern Hess Canyon quadrangle, float lying on the surface of the upper Vidrio.
266 Capitan: Tessey member, .75 mile north of point where many pelecypods were collected (locality 267) north of Hess Canyon quadrangle.
267 Capitan: Tessey member, 5.1 miles west of west side Sierra Madera quadrangle, and 1.8 mile north of north side of Hess Canyon quadrangle, about .07 mile east of north end of a Comanche mesa lying on the east side of the road to Sanderson Cabin. From a bed a foot thick, containing a great abundance of *Pleurophorus*, above a layer of anhydrite which outcrops .5 mile to the east on the west side of a fault.
268 Bissett: Section 24, bed 5, near highest exposures of formation 3 miles north of Warren Ranch, on west side of Hess Canyon.
269 Word: Float .5 mile east of Skinner's gate on the Marathon-Fort Stockton highway.
270 Leonard: Northeast of summit of Dugout Mountain, on Hess cuesta.
301 Leonard: Mid-part of Clay Slide, same as locality 6.
303 Delaware Mountain formation: Limestone in lower part of formation, below Bone Spring, Guadalupe Mountains.
376 Lowest Gym above Canyon-Cisco limestone 3.3 miles west of Helms Peak.
377a Lower Gym: Section west of Powwow Canyon, bed 1.
377b Lower Gym: Section west of Powwow Canyon, bed 2.
378 Lower Gym: Sand-mantled hills 6 miles west-northwest of Helms' West Well.
379 Lower Gym: Basal beds on hill 5 miles N 80° W of the summit of Helms Peak (north of Carlsbad road).
380 Uppermost lower Gym: 5.5 miles west-southwest of Hueco Tanks.
381 Lower Gym: Section west-southwest of Hueco Canyon, bed 1.
382 Lower Gym: Section west-southwest of Hueco Canyon, bed 2.
383 Lower Gym: South of Hueco Tanks road on northwest side of the Hueco anticline, on the east side of the hills.
384 Lower Gym: East end of hills north of Hueco Tanks road, west of Hueco Tanks.
385 Lower Gym: Slopes of Juan Peak.
386 Lower Gym: Section on Juan Peak, bed 9.
387 Lower Gym: Marl underlying the bed capping Juan Peak.
388 Lower Gym: Beds capping Juan Peak.
389 Lower Gym: Ridge east of Juan Peak, from beds stratigraphically above that capping Juan Peak.
390 Upper part of lower Gym: Southern part of Lefthand Canyon.
391 Lower Gym: 1.3 miles north 10° east of hill 5345, south of Hueco Tanks road.
392 Lower Gym: Section just south of the entrance to Hueco Canyon, bed 2.
393 Lower Gym: Limestone inlier in igneous sill .7 mile notrh of hill 5345 El Paso quadrangle, due east of Hueco Tanks.
394 Lower Gym: .8 mile north of hill 5345, south of the Hueco Tanks road, collected directly above the igneous sheet.
395 Lowest Gym: West of hill 5345, El Paso quadrangle south of Hueco Tanks road.
396 Lower Gym: South end of hill 5345, section southeast of Hueco Tanks, bed 1.
397 Lower Gym: Section southeast of Hueco Tanks, bed 2.
398 Lower Gym: Section southeast of Hueco Tanks, bed 5.
399 Lower Gym: Section southeast of Hueco Tanks, bed 8.
400 Basal lower Gym: Southwest corner of Neville Mountain.
401 Lower Gym: 1.2 mile south-southwest of hill 5345, El Paso quadrangle.
402 Lower Gym: Top of hill 1.9 mile south of hill 5345.
403 Lower Gym: 2 miles south-southwest of hill 5345, El Paso quadrangle.
404 Lower Gym: Second range of hills north of Powwow Canyon, 3 miles south of hill 5345, El Paso quadrangle.
405 Basal lower Gym: .8 mile west of point halfway between "O" and "U" of "Mountains," El Paso quadrangle.
406 Lower Gym: North side of Powwow Canyon.
407 Lowest Gym: East wall of upper Powwow Canyon.
408 Lower Gym: 3.8 miles northwest of hill 5436, Cerro Alto quadrangle.
409 Lower Gym: .5 mile northeast of hill 5426, Cerro Alto quadrangle, north of place where section was measured.
410 Lowest Gym: Coyote Tanks, southeast end of Hueco Mountains.
411 Lowest Gym: Outlier on Strawn .4 mile east of west end of Cerro Alto and 1.7 mile south of 31°50′ lat. line.
412 Lower Gym: 2.2 miles south-southeast of b. m. 4523, southwestern Cerro Alto quadrangle.
413 Lower Gym: Summit 4659, south of Padre Mine Canyon.
414 Middle Gym: Summit of highest hill in mountains west of Hueco Tanks, 6 miles west-southwest of Hueco Tanks.
415 Middle Gym: 3.4 miles west and a little south of Hueco Tanks.
416 Middle Gym: Section west-southwest of Hueco Canyon, bed 12.
417 Middle Gym: Lower middle part, 2.2 miles northwest of the summit of Cerro Alto.

418 Lower middle Gym: Above sill, metamorphosed limestone 2 miles south-southwest of southeast corner of Cerro Alto.
419 Middle Gym: Limestone in marly beds above metamorphosed limestone 2 miles southwest of the southwest corner of Cerro Alto.
420 Middle Gym: Southwest foot of Cerro Alto.
421 Middle Gym: Northwest Cerro Alto quadrangle.
422 Middle Gym: 1.4 mile south of summit of Cerro Alto, below *Pugnoides* horizon.
423 Middle Gym: 2 miles south of summit of Cerro Alto, not far below the Upper Gym.
424 Middle Gym: 2.6 miles south 5° east of summit of Cerro Alto, about 75 feet below the Upper Gym.
425 Middle Gym: 2 miles northwest of hill 5700 near west border of Cerro Alto quadrangle, from black limestone near the same horizon as beds just above the sill to the east.
426 Middle Gym: 2 miles north of hill 5700, near west border of Cerro Alto quadrangle.
427 Middle Gym: Just above sill on line of section south of Cerro Alto in basin.
428 Middle Gym?: Southwest of Neville Mountain; possibly in part from Upper Gym.
429 Middle Gym: 1.2 mile east of east summit of Neville Mountain, northwestern Cerro Alto quadrangle.
430 Middle Gym: North of Powwow Canyon near line of section.
431 Middle Gym: Section north of Powwow Canyon, bed 1 of Middle Gym.
432 Middle Gym: Section north of Powwow Canyon, bed 6, directly below Upper Gym.
433 Middle Gym: 1.4 mile west-southwest of b. m. 5149, southwest central Cerro Alto quadrangle.
434 Lower part of Middle Gym: 1.9 mile southwest of b. m. 5149, northwest center of Cerro Alto quadrangle.
435 Upper Gym: 3 miles north of entrance to Hueco Canyon.
436a Upper Gym: Highest beds on section on south side of Hueco Canyon, easternmost Hueco Canyon exposure near Lincoln Tank.
436b Lower Upper Gym: West of Daily Mountain (on east side of road to Bassett Ranch), 1.6 mile west of the summit.
437a Lower upper Gym—about 1.5 miles south of summit of Cerro Alto.
437b Basal upper Gym: South side of Daily Mountain.
438 Upper Gym: 2 miles south of the summit of Cerro Alto, above the red beds.
439 Upper Gym: 1.7 mile south of summit of Cerro Alto.
440a Upper Gym: Summit of Deer Mountain, northwestern Cerro Alto quadrangle .
440b Upper Gym: Limestone capping hill 5700, northwestern Cerro Alto quadrangle.
441 Lower Upper Gym: Below red beds south of Deer Mountain.
442 Gym: 2 miles north-northwest of Dagger Tank, Cerro Alto quadrangle.
443 Lower upper Gym?: 2 miles east of b. m. 4984, 2 miles northeast of summit of Rancheria Mountain.
444 Lower upper Gym?: 2 miles due east of b. m. 4984, 2 miles northeast of summit of Rancheria Mountain, from a level 40 feet above that of 443.
445 Gym: 1.8 mile north of Big Tank (west end of Buckhorn Draw), Cerro Alto quadrangle.
446 Gym: 1.5 mile north of central part of Buckhorn Draw, Cerro Alto quadrangle.
447 Gym: 2.6 miles east-southeast of Panther Tank, Cerro Alto quadrangle.
448 Gym: Hills south of Borrego Tank.
449 Gym: Hills north of tank halfway between Borrego and Shakespeare Tanks.
450 Gym: .2 mile north of first "d" in "Abandoned Tanks," Cerro Alto quadrangle.
451 Gym: Cuesta above sill, west side of Sierra Tinaja Pinta, near old Miller Ranch.
452 Gym: Hills 6 miles south-southwest of the southwest corner of Sierra Tinaja Pinta.
453 Upper Gym: 1 mile southeast of Deer Mountain above red beds.
454 Lower Gym: *Schwagerina* from beds west of fault on west side of Franklin Mountains.
455 Middle Gym: Dark limestone on west side of Franklin Mountains west of main western fault.
456 Upper Wolfcamp: .5 mile southeast of diabase hill 5039, Van Horn quadrangle.
457 Upper Wolfcamp: South of Marble Quarry Canyon.
458 Upper Wolfcamp: Above Strawn limestone southeast of Marble Quarry, bed 12.
459 Upper Wolfcamp: South side of entrance to Victorio Canyon.
460 Wolfcamp: South of Victorio Peak, above Millican.
461 Hess?: Southeast of Apache Peak, arroyo north of hill 4520, float.
462 Hess: Southeast of Apache Peak, halfway up hill 4520.
463 Hess: Southeast of Apache Peak, limestone at east foot of hill 4520.
464 Hess: .3 mile southwest of hill 4520, east of Apache Peak, on east side of fault.
465 Bone Canyon: 1 mile northeast of summit 6630, south of Apache Peak.
466 Hess: Lower cliff, 1.5 mile east of summit 6630, near Marble Quarry.
467 Hess?: Float, 1 mile east of Marble Quarry.
468 Hess: Lower, base of section south of Marble Quarry, float.
469 Hess: Section south of Marble Quarry, bed 14.
470 Hess?: Float, entrance to Victorio Canyon.
471 Lower Hess: Victorio Peak section, bed 13.
472 Hess: Victorio Peak section, bed 14.
473 Hess: Victorio Peak section, bed 15.
474 Hess: Victorio Peak section, bed 16.
475 Hess: Victorio Peak section, bed 18.
476 Hess: Victorio Peak section, beds 20–21.
477 Hess: Victorio Peak section, beds 23–24.
478 Hess: Lower two fossiliferous layers in upper part of Hess, 1.5 mile east of b. m. 6140, Sierra Diablo south of Victorio Peak.
479 Hess: Upper fossiliferous layer in upper part of Hess, 1.5 mile east of b. m. 6140, Sierra Diablo south of Victorio Peak.
480 Hess: Upper part of Hess 1.3 mile south of b. m. 6140, cherty horizon.
481 Hess: Float, 1 mile southeast of b. m. 5560, Baylor Mountains.
482 Hess: Below summit 5560, southeastern Baylor Mountains.
483 Bone Canyon: Apache Peak section; *Fusulina* from the lower part of the member and gastropods from bed 6, below the upper limestone of Apache Peak.
484 Bone Canyon: Shale northeast of summit 6345 (southwest of Victorio Peak) and underlying the limestone which is exposed on that summit.
485 Bone Canyon: Lower part of lower shale above Hess limestone, section south of Victorio Peak.
486 Bone Canyon: Upper shale, section south of Victorio Peak.
487 Bone Canyon: Victorio Peak section, bed 25.
488 Bone Canyon: Black limestone below the shale underlying hill 6345, south of Victorio Peak.
489 Bone Canyon: Conglomerate, about 150 above base, .8 mile southwest of b. m. 6140, southwest of Victorio Peak.
490 Bone Canyon: Section in Delaware Mountains north of Van Horn quadrangle, bed 2.
491 Bone Canyon: Section in Delaware Mountains north of Van Horn quadrangle, bed 5.
492 Victorio Peak: 8.5 miles south of Dos Alamos.

493 Victorio Peak: Upper beds, below Delaware Mountain section 11.5 miles south of Dos Alamos.
494 Victorio Peak: .5 mile west of east edge of Salt Basin quadrangle, in beds dipping from escarpment east into the flat.
495 Victorio Peak: White limestone just east of Salt Basin sheet in beds dipping into flat from the escarpment.
496 Victorio Peak: Victorio Peak section, 150 feet above base of bed 30.
497 Victorio Peak: .6 mile northeast of summit 6630.
498 Victorio Peak: Upper grey limestone at top of section south of Marble Quarry.
499 Victorio Peak: Victorio Peak section, bed 29.
500 Victorio Peak: Same as 496.
501 Victorio Peak: Victorio Peak section, about 459 feet above base of bed 30.
502 Leonard: Section on Baylor Mountains, beds 5–6, mostly bed 5.
503 Leonard: Section on Baylor Mountains, beds 8–9, mostly bed 9.
504 Delaware Mountain: 6 miles south of Dos Alamos.
505 Delaware Mountain: About 30 feet above its base, 8.5 miles south of Dos Alamos.
506 Delaware Mountain: Section 11.5 miles south of Dos Alamos, bed 3.
507 Gym: Section at west side of Wylie Mountains, lower part.
508 Gym: Section at west side of Wylie Mountains, upper part.
509 Gym: Wylie Mountains 3 miles south of Wildhorse.
510 Gym: Lower beds of the Permian 3 miles east of Dalberg, Carrizo Mountains.
511 Hess: Conglomerate bed, Finlay Dome, south of "Fossil Hill."
512 Leonard: Shales just below ammonoid bed, "Fossil Hill," northwestern Finlay Dome.
513 Leonard: "Fossil Hill," Finlay Dome.
514 Leonard: Limestone beds in dark shale near Ross Mine, Shafter quadrangle.
515 Leonard: Limestone in upper beds near divide east of Ross Mine.
516 Leonard: Lower part of section northwest of Shafter Mine.
517 Leonard: Second main limestone northwest of Shafter Mine, on line of section.
518 Leonard: North of Shafter silver mine, probably including some fossils from the basal Word limestone.
519 Leonard: 1.5 mile northwest of Ojo Bonito.
520 Chupadera: 3.5 miles south-southeast of McVeigh Ranch, northeastern Salt Basin quadrangle (collected by Mr. Shapleigh Gray).

## LOCALITIES IN UNIVERSITY OF TEXAS COLLECTIONS*

Ta Word: "Between stations 60 and 62, Green Canyon," Glass Mountains.
Tb Leonard: North side of top of Leonard Mountain, "near stations 80 and 81"; north face of Leonard Mountain, "thin-bedded ledges just below station 81 and above heavy ledge, interbedded with sandstone."
Tba Leonard: North base of Leonard Mountain.
Tc Word: Middle limestone in Hess Canyon, "upper half of of section D, Hess Tank"; "stations 1 to 2 in section northeast of camp at Hess Tank"; "hill below beginning of section D, Hess Tank"; "stations 3 and 4 in section northeast of camp at Hess Tank"; "samples from end of section, after station 7, taken northeast of camp at Hess Tank." Probably collected .5 mile south of hill 5578; as identified by Beede some of these fossils are listed in Udden's section 6, bed 1.
Td Leonard: Horizon in "Cibolo" just above brecciated zone of Cibolo Ranch, hill just east of Cibolo Creek, associated with *Lyttonia*.
Te Gym: Upper beds on 3 Mile Mountain near Van Horn.
Tf Leonard: "Cieneguita beds" 10 miles north-northwest of Shafter.
Tg Leonard: "Cieneguita beds," Shafter, collections 56, 57, and 62.
Th Leonard: Transition beds of "Cibolo," 6 miles north-northwest of Shafter.
Ti Leonard: Above transition beds of "Cibolo," 5 miles northwest of Shafter.
Tj Word: Thin-bedded zone of "Cibolo," 4–5 miles north-northwest of Shafter.
Tk Word: Thin-bedded zone of "Cibolo," 4–5 miles north-northwest of Shafter.
Tl Leonard: Brecciated zone of "Cibolo," 4 miles north-northwest of Shafter.
Tm Leonard: "Cibolo limestone," 4–6 miles northwest of Shafter.
Tn Leonard: "Cibolo limestone," 6–10 miles northwest of Shafter.
To Leonard: "Chinati series" north of Chinati Mountains.
Tp Word: Yellow limestone of "Cibolo limestone," 3 miles north-northwest of Shafter.
Tp Leonard: Transition beds of "Cibolo limestone," 3–4 miles northwest of Shafter.
Tr Leonard?: 10 miles west of Decie (Wedin) Ranch.
Ts Lower Hess: Dugout Mountain.
Tt Hess: About two-thirds mile north of Gap Tank.
Tu Word?: "Numbers 33 and 34."
Tv Hess?: "About the same horizon as the first hogback north of Leonard Mountain."
Tx Word: Middle Word limestone in Hess Canyon, fossils etched free.
T9 Leonard: "Top of ridge 2½ miles northeast of Wedin's oil derrick," "about 3 miles north 20° east of Wedin's oil derrick," "north side of Round Point ridge, 3 miles north of Wedin's oil derrick," "base of hill 2 miles north 20° east of Wedin's oil derrick," "about halfway between first day's section and Wedin's land."
T9a Leonard: 3 miles west of Iron Mountain "above 200 feet of shale, station 3, upper part," "top of bluff, section of October 27, station 3, 1½ mile east (?) of Iron Mountain."
T15 Lower Leonard: 1 mile west of Iron Mountain.
T15a Lower Leonard: "Basal part of section ¾ mile west-southwest of Skinner's West Ranch," top ledge of first ridge west of Iron Mountain.
T41 Leonard: "Near Bird Mine, 200–400 feet below base of dolomite cliff."
T82 Wolfcamp or Hess: Just above the conglomerate "at station 85," south side of Leonard Mountain.
T82a Leonard?: South face of Leonard Mountain.
T85 Wolfcamp?: 1½ mile east-southeast of Hess Tank, uplift northeast of Hess Ranch.
T88 Wolfcamp: *Uddenites* zone at Wolf Camp.
T88a Wolfcamp: Base of *Uddenites* zone.
T90 Middle Wolfcamp: Just below T93.
T91 Wolfcamp: "1 mile northwest of station 3, sheet 2; layer of limestone in shale"; "grey limestone of very fine texture containing shells of ostracods"; "stations 21 and 22."
T93 Middle Wolfcamp: On stream bank northeast of Wolf Camp.
T107 Hess: 2½ miles north 65° west of Wolf Camp.
T108 Upper Hess: Fossiliferous horizon about 150 feet stratigraphically below the top of the cuesta north of Wolf Camp.
T119 Leonard: Shale slide 4 miles northeast of Decie Ranch.

*Where a number is suffixed (as T9) the number corresponds approximately to the locality listed under that number in the foregoing list.

T120 Upper Leonard: Hill east of shale slide "about midway between first day's section and Wedin (Decie) land."
T128 Leonard: Near Word Ranch.
T141 Word: Limestone above shale slide.
T144 Word: Mostly below middle Word ammonoid horizon, Bowman's section 2.5 miles up valley from Hess Tank.
T144a Word: Ravine 3 miles east of Hess Tank.
T148 Word: Tank ½ mile north of Word Ranch and beds above tank.
Tx148 Word: Near the tank ¼ mile north of Word Ranch.
T183 Leonard: South side of Altuda Mountain, near granite hill.
T205 Hess: Basal, 2(?) miles west-northwest of Old Payne Ranch House, Dugout Mountain.
T226 Middle Leonard: Dugout Mountain, "Payne Ranch."
T232 Leonard: Top of Leonard Mountain.
T233 Leonard: South side of Leonard Mountain, "station 83."
T238 Word limestone: Southern part of Permian outcrop on Del Norte range.
T239 Middle Word: ½ mile southeast of junction of Road and Gilliland Canyons.
T240 Middle Word: Mountain 1½ miles north 55° west of old Hess Ranch House, "number 24, section 1," collected by Beede, Bybee, and Christner.
T240a Middle Word: Top of the first mountain north of Leonard Mountain "sta. 74"; "middle of the yellow sandstone between stations 74 and 75."
T243 Middle Word: Road Canyon, "stations 70 and 70–71,"
T251 Lower Word: ¼ mile southwest of Word Ranch on road to Leonard Mountain.
Taa Victorio Peak: Brackett Draw.
Γzz, aa Gym: Western Wiley Mountains.
Tmm Wolfcamp: Top of Beede's section K, 4.5 miles south of Hueco Tanks.

## LOCALITIES IN DELO COLLECTION

The exact localities of this collection are not known. D1 is from the Leonard on Leonard Mountain; D2, D3, D4, D5, D7, D8, D9, D10, D11, D12, D13, D14, D15, D16, D17, D18, D19, and D21 are from various localities in the Leonard formation in the western part of the Glass Mountains; D6 is from the Hess formation; and D239 and part of D3 are from the Word formation. None of these numbers has any relation at all to the numbers in the remainder of the collection, except D239, which is the junction of Road and Gilliland Canyons, middle Word limestone, corresponding to localities 239 and T239.

## LOCALITIES IN SCHUCHERT COLLECTION

The localities in this collection have been numbered to correspond with the numbers in the King and The University of Texas collections. The more important localities are:

23S Middle Leonard: South of Sullivan Peak.
26S Middle Leonard: South of Sullivan Peak.
88S Wolfcamp: *Uddenites* zone at Wolf Camp.
93S Middle Wolfcamp: Stream bank northeast of Wolf Camp.

## LOCALITIES IN R. F. BAKER COLLECTION

This small collection was made from several localities, particularly 88, 93, the hills southwest of Hueco Tanks, the Gym in Hueco Canyon, and locality 520.

## BIBLIOGRAPHY*


1. Abich, H. 1878—Geologische Forschungen in den Kaukasischen Ländern, Theil i. Eine Bergkalkfauna aus der Araxesenge bei Djoulfa in Armenien. Th. iii. Wien. (Not seen.)
2. Arthaber, G. von. 1900—Das jüngere Paläozoicum aus der Araxes-Enge bei Djulfa. In F. Frech and G. von Arthaber, "Ueber das Paläozoicum in Hocharmenien und Persien." *Beitr. z. Pal. und Geol. Öst.-Ungarns und des Orients*, Bd. 12, pp. 161–308. Wien und Leipzig. (Not seen.)
3. Bassler, R. S. 1929—The Permian Bryozoa of Timor. *Paläontologie von Timor*, lief. 16, pt. 28. Stuttgart.
4. Bayan. 1874—Fossiles paléozoïques de Chine. *Bull. Soc. Géol. France*, sér. 3, vol. 2, pp. 409–415. Paris.
5. Beede, J. W. 1907—Invertebrate paleontology of the Upper Permian Red Beds of Oklahoma and the Panhandle of Texas. *Kansas University Science Bull.*, vol. 4, pp. 115–171. Lawrence.
6. Beede, J. W., and Kniker, H. T. 1924—Species of the genus Schwagerina and their stratigraphic significance. *University of Texas Bull.* 2433. Austin.
7. Böse, Emil. 1916—Contributions to the knowledge of Richthofenia in the Permian of West Texas. *University of Texas Bull.* 55. Austin.
8. Böse, Emil. 1917—The Permo-Carboniferous ammonoids of the Glass Mountains, West Texas, and their stratigraphical significance. *University of Texas Bull.* 1762. Austin.
9. Böse, Emil. 1920—On ammonoids from the Abo sandstone of New Mexico and the age of the beds which contain them. *Amer. Jour. Sci.* (4), vol. 49, pp. 51–60. New Haven.
10. Broili, F. 1916—Die Permischen Brachiopoden von Timor. *Paläontologie von Timor*, lief. 7, pt. 12. Stuttgart.
11. Caralp, J. 1903—Le Permien de l'Ariége, ses divers faciés, sa faune marine. *Bull Soc. Géol. France*, sèr. 4, vol. 3. Paris. (Not seen.)
12. Chao, Y. T. 1927—Productidae of China, Part I; Producti. Geol. Surv. of China, *Pal. Sinica*, ser. B, vol. 5, fasc. 2. Peking.
13. Chao, Y. T. 1928—Productidae of China, Part II. Chonetinae, Productinae, and Richthofeninae. Geol. Surv. of China, *Pal. Sinica*, ser. B, vol. 5, fasc. 3. Peking.
14. Colani, M. 1924—Nouvelle contribution a l'Étude des fusulinidés de l'extréme-Orient. *Mém. du Serv. géol. de l'Indochine*, vol. 11, fasc. 1. Hanoi.


*Most of the works dealing with the stratigraphy and paleontology of the marine Permian and high Carboniferous rocks of the world are mentioned here, as well as a few other more or less relevant papers. As a rule, where a work contains nothing which is not discussed more fully in a later publication, and is therefore definitely superseded, that work is not mentioned. Thus the works of Rothpletz and Beyrich on Timor are regarded as superseded by the various parts of the *Paläontologie von Timor*.

**15.** Darton, N. H. 1916—Geology and Underground Water of Luna County, New Mexico. *U. S. Geol. Surv. Bull.* 618. Washington.

**16.** Darton, N. H. 1925—A Resumé of Arizona Geology. *Bull.* 119, *Ariz. College of Mines and Eng.* Tucson.

**17.** Darton, N. H. 1929—"Red Beds" and Associated Formations in New Mexico, with an outline of the Geology of the State. *U. S. Geol. Surv. Bull.* 794. Washington.

**18.** Davidson, T. 1858–1863—British Fossil Brachiopoda. Vol. 2, Permian and Carboniferous Species. *Palaentographical Society*, London.

**19.** Deprat, J. 1912—Étude des Fusulinidés de Chine et d'Indochine et Classification des Calcaires á Fusulines. *Mém. du Serv. géol. de l'Indochine*, vol. 1, fasc. 3, pt. 3. Hanoi-Haiphong.

**20.** Deprat, J. 1913—Les Fusulinidés des Calcaires Carbonifériens et Permiens du Tonkin, du Laos et du Nord-Annam. *Mém. du Serv. géol. de l'Indochine*, vol. 2, fasc. 1, mém. 2. Hanoi-Haiphong.

**21.** Deprat, J. 1914.—Étude comparative des Fusulinidés d'Akasaka (Japon) et des Fusulinidés de Chine et d'Indochine. *Mém. du Serv. géol. de l'Indochine*, vol. 3, fasc. 1, mém. 3. Hanoi-Haiphong.

**22.** Deprat, J. 1915—Les Fusulinidés des Calcaires carbonifériens et Permiens du Tonkin, du Laos et du Nord-Annam. *Mém. du Serv. géol. de l'Indochine*, vol. 4, fasc. 1, mém. 4. Hanoi-Haiphong.

**23.** Derby, O. A. 1874—On the Carboniferous Brachiopoda of Itaitúba, Rio Tapajos, Prov. of Pará, Brazil. *Bull. Cornell University*, vol. 1, no. 2, pp. 1–63. Ithaca.

**24.** Derby, O. A. 1894—The Amazonian Upper Carboniferous Fauna. *Jour. of Geol.*, vol. 2, pp. 480–501. Chicago.

**25.** Diener, C. 1897—The Permocarboniferous fauna of Chitichun No. 1. Mem. Geol. Surv. India, *Pal. Indica*, ser. 15, Himalayan Fossils, vol. 1, pt. 3. Calcutta. (Not seen.)

**26.** Diener, C. 1897.—Ueber ein Vorkommen von Ammoniten und Orthoceren im südtirolischen Bellerophonkalk. *Sitz. Ber. K. Ak. d. Wiss. Wien*, vol. 106. (Not seen.)

**27.** Diener, C. 1899—Anthracolithic Fossils of Kashmir and Spiti. Mem. Geol. Surv. India, *Pal. Indica*, ser. 15, Himalayan Fossils, vol. 1, pt. 2. Calcutta.

**28.** Diener, C. 1901—Ueber die systematische Stellung der Ammoniten des südalpinen Bellerophonkalkes. Central bl. f. Min. (Not seen.)

**29.** Diener, C. 1903—Permian Fossils of the Central Himalayas. Mem. Geol. Surv. India, *Pal. Indica*, ser. 15, Himalayan Fossils, vol. 1, pt. 5. Calcutta.

**30.** Diener, C. 1910—Die Brachiopodenfauna des Bellerphonkalkes von Schaschar und Schönbrunn. *Jahrb. d. k. k. geol. Reichsanst.*, Bd. 60, Heft. 2, pp. 289–309. Wien. (Not seen.)

**31.** Diener, C. 1911—Anthracolithic Fossils of the Shan States. Mem. Geol. Surv. India, *Pal. Indica*, n. s., vol. 3, no. 4. Calcutta. (Not seen.)

**32.** Diener, C. 1915—The Anthracolithic Faunae of Kashmir, Kanaur and Spiti. Mem. Geol. Surv. India, *Pal. Indica*, n. s., vol. 5, no. 2. Calcutta.

**33.** Diener, C. 1927—Leitfossilien des marinen Perm. Leitfossilien, lief. 5. Berlin.

**34.** Douglas, J. A. 1914—Geological Sections through the Andes of Peru and Bolivia: I—From the Coast at Arica in the North of Chile to La Paz and the Bolivian 'Yungas.' *Quart. Jour. Geol. Soc. London*, vol. 70, pp. 1–53.

**35.** Douvillé, H. 1909—Sur quelques Brachiopodes á test perforé: *Syringothyris* du Sud Oranais, *Spiriferella* de la Steppe des Kirghises et *Derbya* du Salt Range. *Bull. Soc. Géol. France*, Paris, sér. 4, vol. 9, pp. 144–157. (Not seen.)

**36.** Dunbar, C. O. 1924—Kansas Permian insects. Part I, The geologic occurrence and the environment of the insects. *Amer. Jour. Sci.* (5), vol. 7, pp. 171–209. New Haven.

**37.** Dunbar, C. O., and Condra, G. E. 1927—The Fusulinidae of the Pennsylvanian System in Nebraska. *Nebraska Geol. Surv. Bull.* 2, ser. 2. Lincoln.

**38.** Du Toit, A. L., and Reed, F. R. C. 1927—A geological comparison of South America with South Africa. Carnegie Institution of Washington, Publication no. 381. Discussed by Schuchert, C., The Continental Displacement Hypothesis as viewed by Du Toit. *Amer. Jour. Sci.*, vol. 16, 1928, pp. 266–274. Reply by Du Toit, *ibid.*, vol. 17, 1929, pp. 179, 183.

**39.** Enderle, J. 1900—Ueber eine anthracolithische Fauna von Balia Maaden in Kleinasien. *Beitr. z. Pal. und Geol. Öst-Ungarns und des Orients*, bd. 13, heft. 2, pp. 49–109. Wien und Leipzig. (Not seen.)

**40.** Etheridge, R. 1872—Description of the Palaeozoic and Mesozoic Fossils of Queensland. *Quart. Jour. Geol. Soc. London*, vol. 38, pp. 317–360.

**41.** Etheridge, R. Jun., 1914—Palaeontological Contributions to the Geology of Western Australia. *West. Austr. Geol. Surv. Bull.* 58, ser. 5, no. 10, Western Australian Carboniferous Fossils, chiefly from Mount Marmion, Lennard River, West Kimberley. Perth.

**42.** Fliegel, G. 1901—Ueber Obercarbonische Faunen aus Ost- und Südasien. *Palaeontographica*, bd. 48, pp. 91–136. Stuttgart.

**43.** Frech, F. 1895—Palaeozoische Faunen aus Asien und Nordafrika. *N. Jahrb. f. Min.*, bd. 2, pp. 47–67. Stuttgart.

**44.** Frech, F. 1900—Zur Kenntniss der mittleren Paläozoicum in Hocharmenien und Persien. In F. Frech and G. von Arthaber, "Ueber das Paläozoicum in Hocharmenien und Persien." *Beitr. z. Pal. und Geol. Öst-Ungarns und des Orients*, bd. 12, pp. 183–207. Wien und Leipzig. (Not seen.)

**45.** Frech, F. 1901—*Die Dyas.* Lethaea Geognostica, bd. 2, lief. 3. Stuttgart. (Not seen.)

46. Frech, F. 1911—China, Ergebnisse Eigener Reisen und darauf gegründeter Studien, von F. F. v. Richthofen, vol. 5, Obere Neodyas. Berlin.
47. Fredericks, G. 1915—Notes Paléontologiques. 1. Sur les *Productus* du Carbonifére supérieur et de l'Artinskien. *Mám. du Comité Géologique*, n. s., 103, pp. 1–63. St. Petersburg.
48. Fredericks, G. 1915—Reticularia M'Coy and Squamularia Gemmellaro, pp. 290–294. (All Russian.)
49. Fredericks, G. 1916—The Palaeontological Notes. 2. On Some Upper Palaeozoic Brachiopoda of Eurasia. *Mém. du Com. Géol.*, n. s., 156. Petrograd.
50. Fredericks, G. 1917—Addenda to the Dissertation "Paleontological Notes. 2. Some Upper Paleozoic Brachiopoda of Eurasia." (All Russian.)
51. Fredericks, G. 1918—On Some Ammonoids of the Upper Carboniferous of the Ural Mountains. *Ann. Soc. Paléont. Russie*, vol. 3, pp. 73–94. (All Russian.)
52. Fredericks, G. 1919—Note sur l'origine de l'appareil septal de *Lyttoniinae*. *Ann. Soc. Paléont. Russie*, vol. 3, pp. 108–110. (All Russian.)
53. Fredericks, G. 1924—Upper Paleozoic of the Ussuriland. I. Brachiopoda. *Rec. Geol. Com. Russian Far East*, no. 28 (1923). Vladivostok. (All Russian.)
54. Fredericks, G. 1925—Upper Paleozoicum of the Ussuriland. II. Permian Brachiopoda of Cape Kalouzin. *Rec. Geol. Com. Russian Far East*, no. 40 (1924). Vladivostok. (All Russian.)
55. Fredericks, G. 1925.—New *Lyttoniinae* from upper palaeozoicum of the Oural. *Bull. Soc. Oural. Sci. Nat.*, pp. 59–65. (All Russian.)
56. Fredericks, G. 1925—The Ufa Plateau in the Krasnoufimsk District. A Preliminary Report. *Bull. Com. Géol. Leningrad*, vol. 44, pp. 521–530.
57. Fredericks, G. 1927?—New *Lyttoniinae* from the Upper Carboniferous of Krasnoufimsk, pp. 83–89. (In English.)
58. Fredericks, G. 1926—Synopsis of the Genera of the family *Spiriferidae* King. *Bull. de l'Acad. des Sci. de l'URSS*, nos. 3–4, pp. 393–432. (All Russian.)
59. Fredericks, G. 1928—Communication for the Classification of the Genus *Productus*. *Bull. Com. Géol.*, t. 46, no. 7. Leningrad.
60. Geinitz, H. B. 1861—Dyas oder die Zechsteinformation und das Rothliegende, heft 1, Die animalischen Ueberreste der Dyas. Leipzig.
61. Gemmellaro, G. G. 1888—La afuna dei calcari con Fusulina della Valle del Fiume Sosio nella Provincia di Palermo. Fasc. 1. Palermo. (Not seen.)
62. Gemmellaro, G. G. 1896—Sopra due nuovi generi di Brachiopodi provenienti dai Calcari con Fusulina della Provincia di Palermo. Palermo.
63. Gemmellaro, G. G. 1898–1899—La Fauna dei Calcari con Fusulina della Valle del Fiume Sosio nella Provincia di Palermo. Fasc. 4, pt. 1, Molluscoidea. Palermo.
64. Gerth, H. 1921—Die Anthozoén der Dyas von Timor. *Paläontologie von Timor*, lief. 9, pt. 16. Stuttgart.
65. Gerth, H. 1926—Die Korallenfauna des Perm von Timor und die Permische Vereisung. *Overd. uit Leidsche Geol. Meded.*, d. 2, af. 1, pp. 7–14.
66. Gerth, H. 19——Die Spongien aus dem Perm von Timor. *Paläontologie von Timor*, lief. 16, pt. 28.
67. Girty, G. H. 1903—The Carboniferous Formations and Faunas of Colorado. *U. S. Geol. Sur. Prof. Paper* 16. Washington.
68. Girty, G. H. 1908—The Guadalupian Fauna, *U. S. Geol. Sur. Prof. Paper* 58. Washington.
69. Girty, G. H. 1908—On Some New and Old Species of Carboniferous Fossils. *Proc. of the U. S. Nat. Mus.*, vol. 34, pp. 281–303. Washington.
70. Girty, G. H. 1910—The Fauna of the Phosphate Beds of the Park City Formation in Idaho, Wyoming, and Utah. *U. S. Geol. Sur. Bull.* 436. Washington.
71. Girty, G. H. 1915—Fauna of the Wewoka Formation of Oklahoma. *U. S. Geol. Sur. Bull.* 544. Washington.
72. Girty, G. H. 1915—Invertebrate Paleontology, in Hinds and Greene, The Stratigraphy of the Pennsylvanian Series in Missouri. *Mo. Bur. of Geol. and Mines*, vol. 13, ser. 2, pp. 263–376. Jefferson City.
73. Girty, G. H. 1920—Carboniferous and Triassic Faunas, in Butler, Loughlin, Heikes, etc., *U. S. Geol. Sur. Prof. Paper* 111, pp. 641–657. Washington.
74. Grabau, A. W. 1920—A New Species of Eurypterus from the Permian of China; A Lower Permian Fauna from the Kaiping Coal Basin. *Bull. Geol. Sur. of China*, no. 2, pp. 61–85. Peking.
75. Grabau, A. W. 1923–1924—Stratigraphy of China, Part I, Palaeozoic and Older. *Geol. Sur. of China*. Peking.
76. Grönwall, K. A. 1917—The Marine Carboniferous of Northeast Greenland and its Brachiopod Fauna. *Medd. om Grönland*, vol. 43, pp. 509–618. Kobenhavn.
77. Haack, W. 1914—Ueber eine marine Permfauna aus Normexiko nebst Bemerkungen über Devon daselbst. *Zeitschr. d. deutsch. geol. Gesellsch.*, bd. 66, pp. 482–504. Berlin.
78. Hall, J., and Clarke, J. M. 1892—An Introduction to the Study of the Genera of Palaeozoic Brachiopoda. *Palaeont. of New York*, vol. 8, pts. 1–2. Albany.
79. Haniel, C. A. 1915—Die Cephalopoden der Dyas von Timor. *Paläontologie von Timor*, lief. 3, pt. 6. Stuttgart.
80. Haug, E. 1894—Les Ammonites du Permien et du Trias. Remarques sur leur classification. *Bull. Soc. Géol. France*, sér. 3, vol 20 Paris. (Not seen)

**81.** Hayasaka, I. 1922—Palaeozoic Brachiopoda from Japan, Korea, and China. Pt. 1: Middle and Southern China; pt. 2: Upper Carboniferous Brachiopoda from the Hon-wei-ko Coal Mines, Machuria. *Science Repts. of the Tohoku Imperial University*, ser. 2, vol. 6, no. 1. Sendai.

**82.** Hayasaka, I. 1922—Some Permian Brachiopods from the Kitakami Mountains. *Jap. Jour. Geol. and Geog.*, vol. 1, no. 2, pp. 51–70. Tokyo.

**83.** Hayasaka, I. 1923—Some Permian Fossils from the Kitakami Mountains. *Jap. Jour. Geol. and Geog.*, vol. 2, no. 4, pp. 107–116. Tokyo.

**84.** Hayasaka, I. 1924—Fossils in the Roofing Slate of Okachi, Prov. Rikuzen. *Jap. Jour. Geol. and Geog.*, vol. 3, no. 2, pp. 45–54. Tokyo.

**85.** Hayasaka, I. 1925—On Some Brachiopods from the Lyttonia Horizon of the Kitakami Mountains. *Jap. Jour. Geol. and Geog.*, vol. 4, nos. 3–4, pp. 89–103. Tokyo.

**86.** Karpinsky, A. 1889—Ueber die Ammoneen der Artinsk-Stufe und Einige mit denselben Verwandte Carbonische Formen. *Mém. de l'Acad. Imp. des Sciences de St. Petersbourg*, ser. 7, t. 37, no. 2.

**87.** Karpinsky, A. 1890—Zur Ammoneen-Fauna der Artinsk-Stufe. *Bull. de l'Acad. Imp. des Sciences de St. Petersbourg*, t. 1, pp. 65–80.

**88.** Keyte, I. A., Blanchard, W. G., and Baldwin, H. L. 1927—Gaptank-Wolfcamp problem of the Glass Mountains, Texas. *Jour. Paleontology*, vol. 1, pp. 175–178.

**89.** Kindle, E. M. 1926—The Occurrence of the Genus Leptodus in the Anthracolithic fauna of British Columbia. *Trans. Royal Soc. of Canada*, ser. 3, vol. 20, sect. 4, pp. 109–111. Ottawa.

**90.** King, P. B., and King, R. E. 1928—The Pennsylvanian and Permian Stratigraphy of the Glass Mountains. *Univ. of Texas Bull.* 2801, pp. 109–145. Austin.

**91.** King, W. 1850—A Monograph of Permian Fossils. *Palaeontograph. Soc.* London.

**92.** Koken, E. 1906—Productus Purdoni im perm von Kaschmir. *Central bl. f. Mineralogie*, etc., pp. 129–131. Stuttgart.

**93.** Koninck, L. G. de. 1847—Recherches sur les Animaux Fossiles. Premiére Partie. Monographie des Genres Productus et Chonetes. Liége.

**94.** Koninck, L. G. de. 1898—Descriptions of the Palaeozoic Fossils of New South Wales. *Mem. of the Geol. Sur. of New South Wales*, Pal., no. 6. Sidney.

**95.** Koker, E. J. 1924—Anthozoa uit het Perm van het eiland Timor. *Jaarb. v. h. Minjnwezen.* (Not seen.)

**96.** Kozlowski, R. 1914—Les Brachiopodes du Carbonifére supérieur de Bolivie. *Ann. de Paléontologie*, t. 9. Paris.

**97.** Krotow, P. 1888—Geologische Forschungen am Westlichen Ural-Abhange in den Gebieten von Tscherdyn und Ssolikamsk. *Mém. du Comité Géol.*, vol. 6, no. 1. St. Petersbourg.

**98.** Lee, W. T., and Girty, G. H. 1909—The Manzano Group of the Rio Grande Valley, New Mexico. *U. S. Geol. Sur. Bull.* 389. Washington.

**99.** Likharev, B. 1920—Review of the Literature of the Upper Permian of European Russia for the Ten Years 1910–1919. Pp. 1–48. (All Russian, not seen.) Peterburg.

**100.** Likharev, B. 1924—Note sur la Faune des Calcaires Permiens du Bassin de la Riviére Ouftiouga, District de Kadnikov, Gouv. de Vologda. *Bull. Com. géol. Leningrad*, vol. 39, pp. 547–562. (Not seen.)

**101.** Likharev, B. 1925—Zur Frage über das Alter der Perm-Kalksteine der Onega-Dwina Wasserscheide. *Mém. Soc. russe Min.*, ser. 2, vol. 54, pp. 108–151.

**102.** Likharev, B. 1924—Un Nouveau représentant des Brachiopodes du paléozoïque supérieur de Caucase du Nord. *Bull. Com. géol. Leningrad*, vol. 43, pp. 713–721.

**103.** Likharev, B. 1922—Restes des Nautiloidea dans les dépôts du Permien supérieur du bassin de la riviére Waga. *Bull. Com. géol. Leningrad*, vol. 41, pp. 53–68.

**104.** Likharev, B. 1924—Quelques bryozoaires du Permien supérieur du gouvernement de Vologda. *Bull. Com. géol. Leningrad*, vol. 43, pp. 1011–1036.

**105.** Likharev, B. 1925—Ueber einen neuen Vertreter der Fam. *Lyttoniidae* aus dem Obercarbon des Ural. *C. R. de l'Acad. des Sci. de Russie*, pp. 4–7.

**106.** Likharev, B. 1926—Sur la Morphologie et la Biologie des Lyttonides. *Ann. Soc. palent. Russie*, vol. 4, pp. 125–136. (All Russian, not seen.)

**107.** Likharev, B. 1928—Ueber einige seltene und neue Brachiopoden aus dem Unterperm des nördlichen Kaukasus. *Palaeont. Zeitsch.*, bd. 10, pp. 258–289. Berlin.

**108.** Mansfield, G. R. 1927—Geography, Geology, and Mineral Resources of Part of Southeastern Idaho, with Descriptions of Carboniferous and Triassic Fossils by G. H. Girty. *U. S. Geol. Sur. Prof. Paper* 152. Washington.

**109.** Mansuy, H. 1908—Contribution a la Carte Géologique de l'Indochine. Paléontologie. Hanoi-Haiphong.

**110.** Mansuy, H. 1913—Faunes des calcaire a Productus de Indochine. *Mém. du Service Géol. de l'Indochine*, vol. 2, fasc. 4, vol. 3, fasc. 3. Hanoi-Haiphong. (Not seen.)

**111.** Mansuy, H. 1916—Nouvelle contribution a l'étude des faunes des calcaire a Productus de l'Indochine. *Mém. du Service Géol. de l'Indochine*, vol. 5, fasc. 4, pp. 27–38. Hanoi-Haiphong. (Not seen.)

**112.** Mansuy, H. 1920—Fossiles des calcaire ouralopermiens du Tran Ninh, Laos. *Mém. du Service Géol. de l'Indochine*, vol. 7, fasc. 1, pp. 49–58. Hanoi-Haiphong. (Not seen.)

**113.** Marcou, J. 1858—Geology of North America, with two reports on the Prairies of Arkansas

and Texas, the Rocky Mountains of New Mexico, and the Sierra Nevada of California. Zurich.

**114.** Meek, F. B. 1864—Palaeontology of California, vol. 4, sect. 1. Description of the Carboniferous Fossils, pp. 3–16. Philadelphia.

**115.** Meek, F. B. 1871—Descriptions of new species of fossils from Ohio and other Western States and Territories. *Proc. Acad. Nat. Sci. Phil.*, pp. 159–184.

**116.** Meek, F. B. 1872—Report on the Palaeontology of Eastern Nebraska; with some remarks on the Carboniferous Rocks of that district. *Final Report of U. S. Geol. Surv. Neb.*, pp. 83–264.

**117.** Meek, F. B. 1876—Report on the Palaeontological Collections of the Expedition. Explorations Across the Gt. Basin of Utah in 1859 by Capt. J. H. Simpson. Appendix J, pp. 337–373. Washington.

**118.** Meek, F. B. 1877—Palaeontology. *Rept. of the U. S. Geol. Expl. of the 40th Parallel, Prof. Pp. of the Engineer Dept., U. S. Army*, no. 18, vol. 4, pt. 1, pp. 1–50. Washington.

**119.** Meek, F. B., and Worthen, A. H. 1873—Palaeontology. Description of Invertebrates from Carboniferous System, in "Geology and Palaeontology," *Geol. Surv. Illinois*, vol. 5, pp. 323–619.

**120.** Meyer, H. L. F. 1914—Carbonfaunen aus Bolivia und Peru. *Neues Jahrb. f. Min.*, etc., bd. 37, pp. 590–652. Stuttgart.

**121.** Meyer, O. E. 1922—Brachiopoden des Perm und Untercarbon der Residentschaft Djambi (Sumatra). *Verh. v. h. Geol.-Mijnbouwkundig Genootschap v. Nederland en Kolonien*, d. v., pp. 203–222, 's-Gravenhage.

**122.** Moore, R. C. 1920—Geology of Kansas. Oil and Gas Resources of Kansas, pt. 2; *State Geol. Sur. of Kansas Bull.* 6. Lawrence.

**123.** Muir-Wood, H. M. 1928—The British Carboniferous Producti. II.—*Productus (sensu stricto); semireticulatus and longispinus* groups. *Mem. Geol. Surv. Gt. Britain*, Palaeontology, vol. 3, pt. 1, pp. 1–217. London.

**124.** Netschajew, A. W. 1911—Die Fauna der Perm-Ablagerungen vom osten und vom Aeussersten Norden des Europaeischen Russlands. *Mém. du Com. Géol.*, St. Petersbourg, n. s., liv. 61.

**125.** Newberry, J. S. 1861—Geological Report, in Lieut. Ives' Report upon the Colorado River of the West. Washington.

**126.** Nikitin, S. 1890—Dépôts Carboniféres et Puits Artésiens dans la région de Moscou. *Mém. du Com. Géol.*, St. Petersbourg, vol. 5, no. 5.

**127.** Nötling, F. 1901—Beiträge zur Geologie der Salt Range, inbesondere der permischen und triassischen Ablagerungen. *N. Jahrb. f. Min.*, Beilagebd. 14, pp. 369–471. Stuttgart. (Not seen.)

**128.** Nötling, F. 1903—Ueber die Skulptur des *Productus abichi Waagen. Centralbl. f. Mineral.*, etc., pp. 529–531. Stuttgart. (Not seen.)

**129.** Nötling, F. 1904—Ueber Medlicottia Waag. und Episageceras n. g. aus den permischen und triassischen Schichten Indiens. *Neues Jahrb. f. Min.*, etc., bd. 19. Stuttgart. (Not seen.)

**130.** Nötling, F. 1904—Ueber das Alter der Otoceras-Schichten von Rimkin Paiar in Himalaya. *Neues Jahrb. f. Min.*, etc., bd. 18, pp. 528–555. Stuttgart. (Not seen.)

**131.** Nötling, F. 1904—Ueber ben Bau und die Organisation der Lyttoniidae Waagen. *Verhandl. d. Deutsch. Zool. Gesellschaft*, pp. 103–122.

**132.** Nötling, F. 1905—Untersuchungen über die Familie Lyttoniidae Waagen emend Noetling, Paleontographica, bd. 51, pp. 129–153. Stuttgart.

**133.** Ozawa, Y. 1927—Stratigraphical Studies of the Fusulina Limestone of Akasaka, Province of Mino. *Jour. of the Faculty of Science, Imp. Univ. of Tokyo*, sect. 2, vol. 2, pt. 3, pp. 121–167. Tokyo.

**134.** Ozawa, Y. 1927—Some Carboniferous Fossils collected in Manchuria and Korea. *Jap. Jour. Geol. and Geog.*, vol. 5, no. 3, pp. 77–94. Tokyo.

**135.** Plummer, F. B., and Moore, R. C. 1921—Stratigraphy of the Pennsylvanian formations of North-Central Texas. *Univ. of Texas Bull.* 2132. Austin.

**136.** Prout, H. A. 1857—Description of new species of Bryozoa from Texas and New Mexico, collected by Dr. George G. Shumard, geologist of the United States expedition for boring artesian wells along the 32d parallel, under the direction of Capt. John Pope, U. S. Corps of Top. Eng. *Trans. St. Louis Acad. Sci.*, vol. 1, pp. 228–235.

**137.** Reed, F. R. C. 1927—Palaeozoic and Mesozoic Fossils from Yun-nan. Mem. Geol. Surv. India, *Pal. Indica*, n. s., vol. 19, mem 1. Calcutta.

**138.** Roemer, E. 1880—Ueber eine Kohlenkalk-Fauna der Westküste von Sumatra. *Palaeontographica*, bd. 27, pp. 1–11. Stuttgart.

**139.** Romer, A. S. 1928—Vertebrate Faunal Horizons in the Texas Permo-Carboniferous Red Beds. *Univ. of Texas Bull.* 2801, pp. 67–107. Austin.

**140.** Schellwien, E. 1892—Die Fauna des karnischen Fusulinenkalks. *Paleontographica*, bd. 48, pp. 91–136. Stuttgart.

**141.** Schellwien, E. 1898—Die Auffindung einer permocarbonischen Fauna in den Ostalpen. *Verh. k. k. geol. Reichsanst.*, no. 16, pp. 358–363. Wien.

**142.** Schellwien, E. 1900—Beiträge zur Systematik der Strophomeniden des oberen Paläozoicum. *Neues Jahrb. f. Min.*, etc., bd. 1, pp. 1–15. Stuttgart.

**143.** Schellwien, E. 1900—Die Fauna der Trogkofelschichten in den Karnischen Alpen und den Karawanken. I. Theil, die Brachiopoden. *Abh. der k. k. Geol. Reichs-Anst.*, bd. 16, heft. 1. Wien.

**144.** Schellwien, E. 1911—Paläozoische und triadische Fossilien aus Ostasien; Futterer's Durch Asien, vol. 3. (Not seen.)

**145.** Schmidt, H. 1924—Die carbonischen Goniatiten Deutschlands. *Jahrb. der Preuss. Geol. Landesandt*, bd. 45.

**146.** Schuchert, C. 1897—American Fossil Brachiopoda. *U. S. Geol. Sur. Bull.* 87. Washington.

**147.** Schuchert, C. 1906—The Russian Carboniferous and Permian Compared with those of India and America. A Review and Discussion. *Am. Jour. Sci.* (4), vol. 22, pp. 29–46, 143–158. New Haven.

**148.** Schuchert, C. 1918—On the Carboniferous of the Grand Canyon of Arizona. *Amer. Jour. Sci.* vol. 45, pp. 347–361.

**149.** Schuchert, C. 1927—The Pennsylvanian-Permian Systems of Western Texas. *Amer. Jour. Sci.* (5), vol. 14, pp. 381–401. New Haven.

**150.** Schuchert, C. 1928—Review of the late Paleozoic Formations and Faunas, with Special Reference to the Ice-Age of Middle Permian Time. *Bull. Geol. Soc. Am.*, vol. 39, pp. 769–886.

**151.** Shumard, B. F. 1858—Notice of new fossils from the Permian strata of New Mexico and Texas, collected by Dr. George G. Shumard, geologist of the United States Government expedition for obtaining water by means of artesian wells along the 32d parallel, under the direction of Capt. John Pope, U. S. Corps Top. Eng. *Trans. St. Louis Acad. Sci.*, vol. 1, pp. 290–297.

**152.** Shumard, B. F. 1859—Notice of fossils from the Permian strata of Texas and New Mexico, obtained by the United States expedition under Capt. John Pope for boring artesian wells along the 32d parallel, with descriptions of new species from these strata and the Coal Measures of that region. *Trans. St. Louis Acad. Sci.*, vol. 1, pp. 387–402.

**153.** Shumard, B. F. 1863—Descriptions of some new Palaeozoic fossils. *Trans. St. Louis Acad. Sci.*, vol. 2, pp. 198–213.

**154.** Smith, J. P. 1903—The Carboniferous Ammonoids of America. *Mon. U. S. Geol. Sur.*, vol. 42. Washington.

**155.** Smith, J. P. 1927—Permian Ammonoids of Timor. *Jaarb. v. h. Mijn. in Ned.-Indië, Verh.*, I, 's-Gravenhage.

**156.** Smith, J. P. 1929—The Transitional Permian Ammonoid Fauna of Texas. *Am. Jour. Sci.*, vol. 17, 63–80. New Haven.

**157.** Stefano, G. D. 1914—Le Richthofenia dei calcari con Fusulina di Palazzo Adriano nella valle del fiume Sosio. *Palaeontographica Italica*, vol. 20, pp. 1–27. (Not seen.)

**158.** Steinmann, G. 1929—Geologie von Peru. Heidelburg.

**159.** Stoyanow, A. A. 1910—On the Character of the Boundary of Palaeozoic and Mesozoic near Djulfa. *Mem. Imp. Russ. Miner. Soc.*, vol. 47, pt. 1, pp. 61–135.

**160.** Stoyanow, A. A. 1915—On some Permian Brachiopoda of Armenia. *Mém. du Com. Géol.*, n. s., liv. 3, Petrograd.

**161.** Stoyanow, A. A. 1926—Notes on Recent Stratigraphic Work in Arizona. *Am. Jour. Sci.* (5), vol. 12, pp. 320–324. New Haven.

**162.** Stuckenberg, A. 1898—Allgemeine Geologische Karte von Russland, Blatt. 127. *Mém. du Com. Géol.*, vol. 1 , no. 1. St. Petersbourg.

**163.** Stuckenberg, A. 1905—Die Fauna der obercarbonischen Suite des Wolgadurchbruches bei Samara. *Mém. du Com. Géol.*, n. s., liv. 23. St. Petersbourg. (Not seen.)

**164.** Tesch, P. 1923—Trilobiten aus der Dyas von Timor und Letti, *Paläontologie von Timor*, lief. 11, pt. 21. Stuttgart.

**165.** Tchernow, A. 1907—L'étage d'Artinsk. 1. Ammonoidés des bassins de Jaïva de Kosva et de Tchoussovaïa. Livr. 1. *Bull. de la Soc. Imp. des Natur. de Moscou*, Année 1906. (Not seen.)

**166.** Thomas, I. 1910—The British Carboniferous Orthotetinae. *Mem. of the Geol. Surv. Gt. Britain*, Palaeontology, vol. 1, pt. 2, pp. 83–134. London.

**167.** Thomas, I. 1914—The British Carboniferous Producti. 1.—Genera *Pustula* and *Overtonia*. *Mem. of the Geol. Surv. Gt. Britain*, Palaeontology, vol. pt. 4, pp. 197–366. London.

**168.** Toula, F. 1875—Permo-Carbon-Fossilien von der Westküse von Spitzbergen (Belsund, Cap Staratschin, Nordfjord). *Neuen Jahrb. für Mineral.*, pp. 1–37. Stuttgart.

**169.** Trenchmann, C. T. 1917—Some Remarkably Preserved Brachiopods from the Lower Magnesian Limestone of Durham. *Geol. Mag.*, vol. 48, pp. 538–543. London.

**170.** Tschernyschew, T. 1889—Allgemeine Geologische Karte von Russland, Blatt. 139. Beschreibung des Central-Urals und des Westabhanges. *Mém. du Com. Géol.*, vol. 3, no. 4. St. Petersbourg.

**171.** Tschernyschew, T. 1902—Die obercarbonischen Brachiopoden des Ural and des Timan. *Mém. du Com. Géol.*, vol. 16, no. 2, lief. 1–2. St. Petersbourg.

**172.** Tschernyschew, T. 1914—Die Fauna der oberpalaeozoischen Ablagerungen des Darvas, lief. 1. *Mém. du Com. Géol.*, n. s., liv. 104. St. Petersbourg.

**173.** Tschernyschew, T., and Stepanow, P. 1916—Obercarbonfauna von König Oscars und Heibergs Land. Rept. of the Second Norwegian Arctic Expedition in the "Fram," 1898–1902, vol. 4, no. 34. Kristiania.

**174.** Tzwetaev, M. 1888—Cephalopodes de la section supérieure du Calcaire Carbonifére de la *Russie* Centrale. *Mém. Com. Géol.*, vol. 5, no. 3. St. Petersbourg. (Not seen.)

**175.** Waagen, W. 1879–1887—Salt-Range fossils. Productus limestone group. Mem. Geol. Surv. India, *Pal. Indica*, ser. 13, vol. 1. Calcutta.

**176.** Waagen, W. 1889–1891—Salt-Range fossils. Geological results. Mem. Geol. Surv. India, *Pal. Indica*, ser. 13, vol. 4, pts. 1–2. Calcutta.

**177.** Wanner, J. 19 —Die permischen Echinodermen von Timor. I. Teil. *Paläntologie von Timor*, lief. 7, pt. 12. Stuttgart.

**178.** Wanner, J. 1922—Die Gastropoden und Lamellibranchiaten der Dyas von Timor. *Paläontologie von Timor*, lief 11, pt. 18. Stuttgart.

**179.** Wanner, J. 1924—Die permischen Echinodermen von Timor. II. Teil. *Paläontologie von Timor*, lief. 14, pt. 23. Stuttgart.

**180.** Wanner, J. 1926—Die Marine Permfauna von Timor. Steinmann-Festschrift, *Geologische Rundschau*, bd. 17a. Berlin.

**181.** Vogl, V. 1913—Die Palaeodyas von Mrzla-Vodica in Kroatien (Karstgebiet). *Jahrb. d. k. Ungar. geol. Reichsanst.*, bd. 21. Budapest. (Not seen.)

**182.** Watson, D. M. S. 1917—*Poikilosakos*, a remarkable new genus of Brachiopods from the Upper Coal-measures of Texas. *Geological Magazine*, n. s., decade 6, vol. 4, pp. 212–219.

**183.** Weller, S. 1898—A Bibliographic Index of North American Carboniferous Invertebrates, *U. S. Geol. Sur. Bull.* 153. Washington.

**184.** Weller, S. 1914—The Mississippian Brachiopoda of the Mississippi Valley Basin. *Illinois Geol. Sur. Mon.* 1. Urbana.

**185.** White, C. A. 1877—Report on the invertebrate fossils collected in portions of Nevada, Utah, Colorado, New Mexico, and Arizona by parties of the expeditions of 1871, 1872, 1873, and 1874. *U. S. Geog. Surv. West of 100th merid.*, vol. 4, pt. 1, pp. 1–219. Washington.

**186.** White, C. A. 1891—The Texas Permian and its Mesozoic types of Fossils. *U. S. Geol. Sur. Bull.* 77. Washington.

**187.** Whitfield, R. P. 1908—Notes and observations on Carboniferous Fossils and semifossil shells brought home by members of the Peary Expedition of 1905–06. *Bull. Amer. Mus. Nat. Hist.*, New York, vol. 24, pp. 51–58.

**188.** Wiman, C. 1914—Ueber die Karbonbrachiopoden Spitzbergens und Beeren Eilands. *Nova Acta Regiae Soc. Scien. Upsaliensis*, ser. 4, vol. 3, no. 8. Upsala.

**189.** Yabe, H. 1908—A Contribution to the Genus Fusulina, with notes on the Fusulina limestones from Korea. *Jour. Coll. Sci., Imp. Univ. Tokyo*, vol. 21, 5. (Not seen.)

**190.** Yakowlew, N. 1912—Die Fauna der Oberen Abtheilung der Palaeozoischen Ablagerungen im Donez-Basin. III. Die Brachiopoden. *Mém. du Com. Géol.*, n. s., liv. 70. St. Petersbourg.





FIG. 1. CORRELATION CHART.





FIG. 2. TABLE OF FOSSILS







FIG. 3. TABLE OF FOSSILS



| SPECIES | WOLFCAMP: Upper(Los Zorros-Glass Mts) | WOLFCAMP: Middle&Upper(Word-Glass Mts) | WOLFCAMP: Upper(N.C.-Sierra Diablo) | GYM: HUECO MOUNTAINS Lower | GYM: HUECO MOUNTAINS Middle | GYM: HUECO MOUNTAINS Upper | GYM: Diablo Plateau | GYM: Wylie Mountains | GYM: Carrizo Mountains | HESS: Glass Mountains | HESS: Sierra Diablo | HESS: Finlay Mountains | LEONARD: Glass Mountains | LEONARD: SIERRA DIABLO Bone Canyon Member | LEONARD: Victorio Peak Member | LEONARD: Bone C Member-Cutoff Mbr | LEONARD: Baylor Mountains | LEONARD: Finlay Mountains | LEONARD: Shafter Area | WORD AND DELAWARE MTS: Glass Mountains | WORD AND DELAWARE MTS: Sierra Diablo | WORD AND DELAWARE MTS: Delaware-Guadalupe Mts | WORD AND DELAWARE MTS: Shafter Area | CAPITAN: GLASS MOUNTAINS Altuda Member | CAPITAN: GLASS MOUNTAINS Gilliam Member | CAPITAN: GLASS MOUNTAINS Tessey Member | CAPITAN: Guadalupe Mountains | Bissett Formation | LAS DELICIAS: Limestone of Cerro Castillo | LAS DELICIAS: Wargentinoceras Horizon | LAS DELICIAS: Limestone of Pichaguia | NEW MEXICO: Abo Formation | NEW MEXICO: Yeso Formation | NEW MEXICO: San Andres Formation | Kaibab (Arizona) | Phosphoria(Idaho,Utah,etc) | Pennsylvanian System | REALM | RELATED FOREIGN SPECIES | GIRTY'S NAME (PROF. PAP. 58) |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| *Scacchinella gigantea* SCHELLWIEN | X | R | | | | | | | | C | | | | | | | | | | R | | | | | | | | | | | | | | | | | | E | *S variabilis* GEMMELLARO | |
| *Aulosteges beedei* KING | | | | | | | | | | | | | X | | | | | | | R | | | | | | | | | | | | | | | | | | B | | |
| *A guadalupensis* SHUMARD | | | | | | | | | | | | | | | | | | | | A | X | G | | | | | G | | | | | | | | | | E | *A delhousi* DAVIDSON | *Aulosteges guadalupensis* SHUMARD |
| *A magnicostatus* GIRTY | | | | | | | | | | X | | | C | X | | | | | | R | | | | | | | | | | | | | | | | | B | | *Aulosteges magnicostatus* n sp |
| *A medlicottianus* WAAGEN | | R | | | | | | | | A | | | A | | | | | | | X | | | | | | | | | | | | | | | | | E | | |
| *A subcostatus* KING | | | | | | | | | | | | | X | | | | | | | | | | | | | | | | | | | | | | | | B? | | |
| *A trigonalis* KING | | | | | | | | | | R | | | C | ? | | | X | | | X | | | | | | | | | | | | | | | | | B | | |
| *A tuberculatus* KING | | | | | | | | | | | | | | | | | | | | C | X | | | | | | | | | | | | | | | | B | | |
| *A wolfcampensis* KING | C | A | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | A? | | |
| *Strophalosia hystricula* GIRTY | | X | | | | | | | | | X | | X | X | | | | | | X | | | | | | | | | | | | | | | | | A? | | *Strophalosia hystricula* n sp |
| *Tegulifera bosei* KING | ? | C | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | E | *T deformis* SCHELLWIEN | |
| *Prorichthofenia lineata* KING | | | | | | | | | | C | C | | A | X | | | X | | X | | | | | | | | | | | | X | | | | | | B | | |
| *P permiana* (SHUMARD) | | | | | | | | | | | | | | | | | | | | A | | G | | | | | G | | ? | | | | | | | | B | | |
| *P tegulifera* KING | | | | | | | | | | C | C | C | | | | | | | | | | | | | | | | | | | | | | | | | B | | |
| *P wuddani* (BOSE) | | | | | | | | | | | | | C | X | | | | | X | X | | | | | | | | | | | | | | | | | B | | |
| *Parawaagenoconcha frederickensis* KING | | X | | | | | | | | ? | | | ? | | | | | | | | | | | | | | | | | | | | | | | | E | *P derbyensis* (TSCHERNYSCHEW) | |
| *Lyttonia nobilis americana* (GIRTY) | | | | | | | | | | C | C | | A | X | X | | | | | A | X | G | | X | | G | | X | | | | | | | | E | *L nobilis* WAAGEN | *Leptodus americanus* n sp *L guadalupensis* n sp |
| *L hartoni* KING | | | | | | | | | | R | | | R | | | | | | | R | | | | | | | | | | | | | | | | | B | | |
| *Liorhynchus biplicata* (SHUMARD) | | | | | | | | | | | | | R | | | | | | | | | G | | | | | G | | | A | | | | | | | D | | *Agnus biplicata* SHUMARD; var. *gaticia* & *semiplicata* |
| *L weeksi nobilis* (GIRTY) | | | | | | | | | | | | | | | | | | | | X | | | | | | | | | | | | | | | X | | D | | |
| *Agnoides bidentata* (WAAGEN) | | | | | | | | | | | | | X | | | | X | | | | | G | | | | | | | | | | | | | | | B | | *Agnus bidentata* n sp |
| *P elegans* (GIRTY) | | | | | | | | | | X | X | | X | | | | | | | X | | G | | | | | G | | | | | | | | | | B | | *Pugnax elegans* n sp |
| *P pingus* (GIRTY) | | | | | | | | | | X | | | X | | | | | | | R | | G | | | | | | | | | | | | | | | B | | *Pugnax pingus* n sp |
| *P shumardianus* (GIRTY) | | | | | | | | | | | | | R | | | | | | | | | | | | | | G | | | | | | | | | | B | | *Pugnax shumardiana* n sp |
| *P wallerianus* (SHUMARD) | | | | | | | | | | | | | | | | | | | | X | | G | | | | | G | | | | | | | | X | | D | | *Pugnax wallerianus* SHUMARD |
| *P cogenus* (SHUMARD) | | | | R | C | A | | | | X | | | R | ? | | G | | | | | | G? | | | | | | | | ? | | | | | | | | B | | *Agnus elegenus* SMITH; *Rhynchonella? reversa* SHUMARD |
| *P transversus* KING | | | | | | | | | | C | | | | | | | | | | | | | | | | | | | | | | | | | | | B | | |
| *Rhynchopora nevadensis* MEEK & WORTHEN | R | R | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | X | A,C | *R nikitini* TSCHERNYSCHEW | |
| *R taylori* GIRTY | | | | | | | | | | | | | | | | | | | | X | | | | | | | | | X | | | | | | X | | D | | |
| *Camarophoria dolo* KING | | | | R | | | | | | | | | | | | | | | | R | | | | | | | | | | | | | | | | | B | | |
| *C thevenini* ROTHPLETZ | | R | | R | | | | | | | | | R | | | | | | | | | | | | | | | | | | | | | | | | C | | |
| *C venusta* GIRTY | | R | | | | | | | | C | | | A | X | | X | X | | X | C | X | | | X | | G | | | | | | | | | | | E | *C pardoni* DAVIDSON | *Camarophoria venusta* n sp |
| *C? indentata* (SHUMARD) | | | | | | | | | | | | | | | | | | | | ? | | G | | ? | | G | | ? | | | | | | | | | B | | *Rhynchonella? indentata* SHUMARD |
| *C? longeva* (GIRTY) | | | | | | | | | | | | | | | | G? | | | | | | | | | | G | | | | | | | | | | | B | | *Rhynchonella? longeva* n sp |
| *Uncinulodes guadalupensis* (SHUMARD) | | R | | | | | | | | R | | | C | | | | | | | X | | | | | | G | | | | | | | | | | E | *U jabiensis* WAAGEN | *Rhynchonella? guadalupae* SHUMARD |

FIG. 4, TABLE OF FOSSILS



| SPECIES | WOLFCAMP: Hidalgo Tongue–Glass Mts. | WOLFCAMP: Madera Limestone–Gloss Mts. | WOLFCAMP: Upper Wolfcamp–Sierra Diablo | GYM: Hueco Mountains: Lower | GYM: Hueco Mountains: Middle | GYM: Hueco Mountains: Upper | GYM: Diablo Plateau | GYM: Wylie Mountains | GYM: Carrizo Mountains | HESS: Glass Mountains | HESS: Sierra Diablo | HESS: Finlay Mountains | LEONARD: Sierra Diablo: Glass Mountains | LEONARD: Sierra Diablo: Bone Canyon Member | LEONARD: Victoria Peak Member | LEONARD: Burnet Member–Guadalupe? | LEONARD: Baylor Mountains | LEONARD: Finlay Mountains | LEONARD: Skelter Area | WORD AND DELAWARE MTS.: Glass Mountains | WORD AND DELAWARE MTS.: Sierra Diablo | WORD AND DELAWARE MTS.: Delaware–Guadalupe Mts. | CAPITAN: Glass Mountains: Skelter Area | CAPITAN: Glass Mountains: Altuda Member | CAPITAN: Glass Mountains: Gilliam Member | CAPITAN: Glass Mountains: Tessey Member | CAPITAN: Glass Mountains: Guadalupe Mountains | LAS DELICIAS: Bissett Formation | LAS DELICIAS: Limestone of Cerro Caballo | LAS DELICIAS: Wadgemoderas Horizon | LAS DELICIAS: Limestone of Pichegua | NEW MEXICO: Abo Formation | NEW MEXICO: Yeso Formation | NEW MEXICO: San Andres Formation | Kolob (Arizona) | Precipice (Idaho, Utah, etc.) | Pennsylvanian System | REALM | RELATED FOREIGN SPECIES | GIRTY'S NAME (PROF PAPER) |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| *Spirifer condor* DOBRIGHY | X | X | | C | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | C | | |
| *S marcoui intraplica* KING | | | | | | | | | | | | | X | X | X | | | X | X | | | | | | | | | | | | | | | | | | | | | E | *S. marcoui* WAAGEN | |
| *S. (Neospirifer) bakeri* KING | | | | | | | | | | | | | R | | | | | | | X | | | | | | | | | | | | | | | | | | | E | *S. niger* WAAGEN | |
| *S. (Neospirifer) costella* KING | | | | | | | | | | | | | R | | | | | | | | | | | | | | | | | | | | | | | | | | E | *S. fasciger* KEYSERLING | |
| *S. (Neospirifer) hueccensis* KING | | | | X | | | | | | X | | | | | | | | | | | | | | | | | | | | | | | | | | | | B? | | |
| *S. (Neospirifer) mexicanus* SHUMARD | | | | | | | | | | | | | | | | | | | | | | G | | | | | | G | | | | | | | | | | E | *S. oldhamianus* WAAGEN | *Spirifer mexicanus* SHUMARD |
| *S. (Neospirifer) mexicanus compactus* GIRTY | | | | | | | | | | | | | | | | | | | | | | | | | | | | G | | | | | | | | | | E? | | *Spirifer mexicanus* var. *compactus* n. var. |
| *S. (Neospirifer) lævis* KING | | | | | | | | | | R | | | X | X | | | | | | | | | | | | | | | | | | | | | | | | | E | | |
| *S. (Neospirifer) pseudocameratus* GIRTY | | | | | | | | | | | | | A | X | X | | X | | X | A | X | G | | | | | G | | C | | | | | | X | | | D | | *Spirifer* sp. b |
| *S. (Neospirifer) texanus* MEEK | X | X | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | X | | | A | | |
| *S. (Neospirifer) triplicatus* HALL | C | C | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | X | | | A | | |
| *S. (Ethinia) sulcifer* SHUMARD | | | | | | | | | | | | | | | | | | | | A | | G | | | | | G | | | | | | | | | | | E | *S. rajah* SALTER | *Spirifer sulcifer* SHUM. *Spirifer* sp. B |
| *Squamularia guadalupensis* (SHUMARD) | | X | | | | | | | | C | | | C | X | X | | | | C | C | | G | X | A | | G | | | | | | ? | ? | | | | B,E | *S. indica* WAAGEN | *Squamularia guadalupensis* SHUMARD |
| *S. perplexa* (MC CHESNEY) | X | ? | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | X | | | A | | |
| *Ambocoelia planoconvexa* var. *guadalupensis* GIRTY | | | | | X | | | | | | | | X | | | X | | | X | X | | G | | | | | G | | | | | | | | X | | B,D | | *Ambocoelia planoconvexa* var. *guadalupensis* n. var. |
| *A planoconvexa* (SHUMARD) | C | ? | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | X | | | A | | |
| *Martinia rhomboidalis* GIRTY | | | | | | | | | | | | | X | | X | X | | | | X | | G | | A | | | G | | | | | | | | | | | E | *M. variabilis* GEMELLARO | *Martinia rhomboidalis* n. var. |
| *M. wellsi campensis* KING | A | C | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | E | *M. triquetra* TSCHERNYSCHEW (non GEMMELLARO) | |
| *Spiriferina angulata* KING | | | | | | | | | | X | X | | | | | | | | | | | | | | | | | | | | | | | | | | | B? | | |
| *S. hilli* GIRTY | | | | | | | | | | X | | | X | | | | | | X | | | | | | | | | | | | X | | | | | | | E | *S. multiplicata* (MOWERBY) | *Spiriferina hilli* n. sp. |
| *S. texa* GIRTY | | | | | | | | | | | | | | | | | | | | X | | G | | X | | | G | | | | X | | | | | | | E | *S. cristata* (SCHLOTHEIM) | *Spiriferina texa* n. sp. *S. walleri* n. sp. |
| *S. pyramidalis* GIRTY | | | | | | | | | | | | | | | | | | | | R | | G | | | | | G | | | | | | | | | | | B | | *Spiriferina pyramidalis* n. sp. |
| *Punctospirifer billingsi* (SHUMARD) | | | | | | | | | | | | | | | | | | | | X | | G | | X | | | G | | | | | | | | | | | E | *P. ornata* (WAAGEN) | *Spiriferina billingsi* SHUMARD |
| *Rhentuckyensis* (SHUMARD) | R | R | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | X | | | A | | |
| *Hustedia biparsita* GIRTY | | | | | | | | | | | | | R | | | | | | | | | G | | | | | | | | | | | | | | | | B | | *Hustedia biparsita* n. sp. |
| *H. hessensis* KING | | | | | | | | | | X | X | | | | | | | | | | | | | | | | | | | | | | | | | | | B | | |
| *H. hueccensis* KING | | | | | | X | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | B | | |
| *H. meekana* (SHUMARD) | | | | | | | | | | X | | | A | | X | X | X | | X | A | X | G | | X | | | G | | X | | X | | | | | | | E | *Hustedia indica* (WAAGEN) | *Hustedia meekana* SHUMARD |
| *H. mormoni* (MARCOU) | C | A | | C | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | X | | | A | | |
| *H. mormoni papillata* (SHUMARD) | | | | | | | | | | | | | X | | | G | | | | C | | G | | | | | G | | | | | | | | | | | B | | *Hustedia papillata* SHUMARD |
| *H. pusilla* (GIRTY) | | ? | | | | | | | | X | | | | | | G | | | | X | | | | | | | | | X | | X | | | | | | B? | | *Pugnax? pusilla* n. sp. |
| *Composita emarginata* GIRTY | | | | | | | | | | | | | | | | | | | | | | G | | | | | G | | | | | | | | | | | B | | *Composita emarginata* n. sp. |
| *C. emarginata effinis* GIRTY | | | | | | | | | | | | | | | | | | | | A | X | G | | X | | | G | | X | | | | | | | | | B,E | *C. extra* (DIENER) | *Composita emarginata* var. *effinis* n. var. |
| *C. mexicana* (HALL) | | | | X | C | A | X | | | X | X | | X | | X | G | | | X | | | | | | | | | | | | | | X | X | | X | | B | | *Composita mexicana* HALL. *Camara* var. *guadalupensis* Vish |
| *C. mira* (GIRTY) | | | ? | | | | | | | X | | | C | | | X | X | | X | C | X | | | | | | | X | | | | | | | X | | | B,D | | |
| *C. persimilata* (MEEK) | | | | | | | | | | | | | | | | | | | | R | | | | | | | | | | | | | | | X | | | D | | |
| *C. subtilita* (HALL) | C | X | | C | | | | X | | A | X | | A | X | X | | X | | X | | | | | | | | | | | | | | X | X | | X | | A,C,E | | |
| *C. subtilita angusta* KING | X | X | A | | | | | | | | | | ? | | | | | | | | | | | | | | | | | | | | | | | | | | B | | |
| *Dielasma bovidens* (MORTON) | X | X | | X | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | X | A,C | | |
| *D. problematicum novilunæ* KING | | | | | | | | | | | | | | | | | | | | R | | | | | | | | | | | | | | | | | | E | *D. problematicum* (DAVIDSON) | |
| *D. shafterense* KING | | | | | | | | | | | | | | | | | | | R | | | | | | | | | | | | | | | | | | | B | | |
| *D. spatiolatum* GIRTY | | | | | | | | | | | | | ? | | | | | | | X | | G | X | | | | G | | | | | | | | | | | E | *D. truncatum* (WAAGEN) | *Dielasma spatiolatum* n. sp. *D. curvatum* n. sp. |
| *D. prolongatum* GIRTY | | | | | | | | | | | | | | | | | | | | | | G | | | | | G | | | | | | | | | | | B | | *Dielasma prolongatum* n. sp. |
| *D. schucherti minor* KING | | | | | | | | | | | | | | | | | | | | X | | | | | | | | | | | | | | | | | | E | *D. bipilex* WAAGEN | |
| *Dielasmina guadalupensis* GIRTY | | | | | | | | | | R | | | | | | | | | | | | G | | | | | G | | | | | | | | | | | E | | *Dielasmina guadalupensis* n. sp. |
| *Notothyris schucherti* GIRTY | | | | | | | | | | | | | | | | | | | | | | G | | | | | G | | | | | | | | | | | E | | *Notothyris schucherti* Girty var. *data* Girty |
| *Heterelasma shumardianum* GIRTY | | | | | | | | | | | | | | | | | | | | | | G | | | | | G | | | | | | | | | | | B? | | *Heterelasma shumardianum* n. sp. |

**LEGEND**

REALM
- A-Descended from American Pennsylvanian species
- B-Descended from American Permian species
- C-Tethyan
- D-Andean
- E-Arctic

RELATIVE ABUNDANCE
- A-Abundant
- C-Common
- X-Sporadically common
- R-Rare
- ?-Questionable
- G-Listed by Girty where not found by King

# PLATES I TO XLIV

Except where otherwise indicated, all figures on these plates are natural size.

## PLATE I

Figure— PAGE

1. *Orbiculoidea missouriensis* (Shumard) ..... 42
   Brachial valve from the *Waagenoceras* zone (Word) at Las Delicias, Mexico. Y.P.M. 12698.

2–4. *Rhipidomella hessensis*, n. sp. ..... 43
   2a–d. Brachial, pedicle, lateral, and posterior views of a cotype from loc. 107. Y.P.M. 12642.
   3a–b. Pedicle and lateral views of a large specimen from loc. 222. Y.P.M. 12643.
   4. Brachial view of a cotype from loc. 107. Upper Hess formation, Glass Mountains. T. 10413.

5–7. *Rhipidomella leonardensis*, n. sp. ..... 43–44
   5a–b. Pedicle exterior and interior views of a cotype from loc. 123 (×3). Y.P.M. 12647.
   6. Pedicle view of a cotype from loc. 123. T. 10507.
   7. Brachial view of a cotype from loc. 123 (×3). Y.P.M. 12647. Leonard formation, Glass Mountains.


8–9. *Rhipidomella mesoplatys baylorensis*, n. var. ..... 44
   8. Brachial view of a paratype from loc. 485. Bone Canyon limestone, Sierra Diablo. T. 11051.
   9a–b. Posterior and brachial views of the holotype, from loc. 503. Y.P.M. 12648. Leonard formation, Baylor Mountains.

10–11. *Rhipidomella mesoplatys*, n. sp. ..... 44
   10a–c. Posterior, pedicle, and brachial (×2) views of the holotype, from loc. 123. Y.P.M. 12680.
   11a–b. Lateral and brachial (×2) views of a somewhat distorted specimen from loc. 256. T. 10551. Leonard and Word (?) formations, Glass Mountains.

12–13. *Rhipidomella transversa*, n. sp. ..... 44
   12a–c. Pedicle, lateral, and brachial (×3) views of the holotype, from loc. 123. Y.P.M. 12650.
   13a–c. Brachial, posterior, and pedicle (×3) views of a specimen from loc. T15a. Leonard formation, Glass Mountains.

14–15. *Orthotichia kozlowskii*, n. sp. ..... 45
   14a–b. Brachial and pedicle views of the holotype, from loc. 92. Y.P.M. 12651. Wolfcamp formation, Glass Mountains.
   15a–b. Brachial and pedicle views of a somewhat crushed specimen from loc. 456. Wolfcamp formation, Sierra Diablo.

16–20. *Parenteletes cooperi*, n. gen., n. sp. ..... 41
   16. Pedicle valve of a cotype from loc. 273, in which the posterior part has been ground away, showing the arrangement of the internal plates. Gaptank formation. Y.P.M. 12678.
   17a–d. Brachial, pedicle, lateral, and posterior views of a cotype from loc. 199. Y.P.M. 12679.
   18. Interior of the pedicle valve of a cotype from loc. 95S, showing arrangement of internal plates. Y.P.M. S1494.
   19a–b. Pedicle and posterior views of a broken cotype from loc. 199. Y.P.M. 12679.
   20. Posterior view of a cotype from loc. Tba. Wolfcamp formation, *Uddenites* zone, Glass Mountains.


1
2a
2b
2c
2d
3a
3b
4
6
5a
5b
7
8
9a
9b
10a
10b
11a
12a
13a
11b
12b
13b
10c
13c
12c
14a
14b
15a
15b
16
17a
17b
17c
17d
18
19a
19b
20

## PLATE II

Figure— PAGE

1–3. *Parenteletes cooperi*, n. gen., n. sp. 49

1a–c. Brachial, pedicle, and lateral views of a specimen from loc. T. 65.

2. Brachial view of a specimen from loc. 273. Y.P.M. 12678.

3a–d. Lateral, anterior, posterior, and antero-ventral views of a specimen from loc. T. 65. Gaptank formation, Marathon basin.

4–8. *Enteletes dumblei* Girty 45

4. Brachial interior view of a specimen from loc. Tx. T. 11047.

5. Pedicle interior view of a specimen from loc. Tx. T. 11047.

6a–c. Pedicle exterior views, and oblique lateral view of pedicle interior, of a specimen from loc. Tx. T. 11047. Word formation, Glass Mountains.

7a–c. Brachial, lateral, and posterior views of a specimen from loc. 222. Y.P.M. 12674.


8a–d. Brachial, ventral, lateral, and posterior views of a specimen from loc. 222. Y.P.M. 12674. Upper Hess formation, Glass Mountains.

9–11. *Enteletes liumbonus*, n. sp. 46

9. Pedicle view of a paratype from loc. 17. Y.P.M. 12668.

10a–c. Lateral, brachial, and posterior views of a paratype from loc. T. 15a. Hess formation, Glass Mountains.

11a–b. Pedicle and lateral views of a paratype from loc. 381. Y.P.M. 12669. Gym formation, Hueco Mountains.


1a
1b
1c
2
3a
3b
3c
3d
5
6a
6b
6c
4
7a
7b
7c
8a
8b
8c
8d
9
10a
10b
10c
11a
11b

## PLATE III

Figure— PAGE

1–2. *Enteletes liumbonus*, n. sp. 46

1a–d. Brachial, pedicle, lateral, and posterior views of a paratype from loc. 17. Y.P.M. 12668. Hess formation, Glass Mountains.

2a–c. Brachial, pedicle, and lateral views of the holotype from loc. 467. Y.P.M. 12670. Hess formation, Sierra Diablo.

3–8. *Enteletes plummeri*, n. sp. 47

3a–b. Brachial and lateral views of a specimen from loc. T. 15. Lower Leonard formation, Glass Mountains. T. 10410.

4a–c. Brachial, pedicle, and lateral views of a specimen from loc. T. 8. T. 10410. Upper Hess formation, Glass Mountains.

5a–b. Brachial and lateral views of a specimen from loc. T. 9.

6a–b. Brachial and pedicle views of a cotype from loc. 232. Y.P.M. 12666.

7. Posterior view of a cotype from loc. 104. Y.P.M. 12667.

8. Lateral view of a specimen from loc. T. 15. Lower Leonard formation, Glass Mountains.

9. *Enteletes wolfcampensis*, n. sp. 47

9a–c. Pedicle, brachial, and lateral views of the holotype, from loc. 196. Y.P.M. 12645. Wolfcamp formation, Glass Mountains.

10–12. *Enteletes leonardensis*, n. sp. 46

10a–c. Brachial, pedicle, and posterior views of a specimen from loc. 28. Y.P.M. 12671.

11a–c. Lateral, posterior, and brachial views of the holotype, from loc. 128. Y.P.M. 12673.


12a–b. Brachial and posterior views of a specimen from loc. 37. T. 11034. Leonard formation, Glass Mountains.

13–15. *Enteletes wordensis*, n. sp. 47–48

13a–d. Brachial, pedicle, lateral, and posterior views of the holotype, from loc. 264. Y.P.M. 12656.

14a–b. Brachial and lateral views of a brachial valve from loc. 255. Y.P.M. 12655.

15a–c. Brachial, posterior, and lateral views of a brachial valve from loc. 255. T. 10550. Word formation, Glass Mountains.


1a
1b
1c
1d
2a
2b
2c
3a
3b
4a
4b
4c
5a
5b
6a
6b
7
8
9a
9b
9c
10a
10b
10c
11a
11b
11c
12a
12b
13a
13b
13c
13d
14a
14b
15a
15b
15c

## PLATE IV

Figure— PAGE

1–3. *Streptorhynchus lamellatum*, n. sp. 49

1a–c. Brachial, lateral, and posterior views of the holotype, from loc. 245. In 1c the pedicle valve is up. Note dorsal cardinal area. Y.P.M. 11399. Leonard formation, Glass Mountains.

2. Brachial view of a specimen from loc. 105. Y.P.M. 11398. Hess formation, Glass Mountains.

3. Brachial view of a specimen from loc. T. 93. T. 10684. Wolfcamp formation, Glass Mountains.

4–6. *Streptorhynchus pygmæum* Girty 49–50

4. Brachial view of a specimen from loc. T. 240. T. 10573. Word formation, Glass Mountains.

5a–c. Pedicle, brachial, and posterior views of a specimen from loc. 26S. Y.P.M. 11387. Leonard formation, Glass Mountains.

6a–b. Pedicle and lateral views of a large specimen from loc. T. 148. T. 10578. Word formation, Glass Mountains.

7–10. *Streptorhynchus pyramidale*, n. sp. 50–51

7a–c. Exterior, interior, and lateral views of a brachial valve of a cotype from loc. 93. Y.P.M. 11392.

8a–c. Pedicle, cardinal, and lateral views of a pedicle valve of a cotype from loc. 93; the apical part has been ground off. Y.P.M. 11392.

9. Brachial view of a cotype from loc. 93. Y.P.M. 11392.

10. Brachial view of a cotype from loc. 93. T. 10581. Wolfcamp formation, Glass Mountains.

11–13. *Streptorhynchus? undulatum*, n. sp. 51

11a–d. Brachial, pedicle, posterior, and lateral views of the holotype, from loc. 211. In 11d the pedicle valve is up, the brachial down. Y.P.M. 11406.

12a–b. Pedicle and posterior views of a specimen from loc. T. 9. In 12b the pedicle valve is up, the brachial down. T. 10561.

13. Posterior view of a specimen from loc. T. 9, with the cardinal area exfoliated. T. 10661. Hess formation, Glass Mountains.

14–15. *Orthothetina* sp. (cf. *Orthothetes mutabilis* Girty) 52


14a–b. Brachial and lateral views of a specimen from loc. 91. Y.P.M. 11420. Wolfcamp formation, Glass Mountains.

15a–b. Brachial and posterior views of a specimen from loc. 354. Y.P.M. 11424. Limestone of Strawn age, Hueco Mountains.

16–17. *Meekella difficilis* Girty 53–54

16a–d. Pedicle, brachial, lateral, and posterior views of a specimen from loc. D. 1. In 16d the pedicle valve is up, the brachial down.

17. Pedicle valve from loc. 235. Y.P.M. 10897. Leonard formation, Glass Mountains.


1a
1b
1c
4
5a
5b
5c
6a
6b
2
3
7a
7b
7c
8a
8b
8c
9
10
11c
11a
11b
11d
12a
12b
13
14a
14b
15a
15b
16a
16b
16c
16d
17

## PLATE V

Figure— PAGE

1. *Meekella difficilis* Girty ..... 53

1a–b. Brachial and posterior views of a brachial valve from loc. 26S. Y.P.M. 10992.

2–7. *Meekella attenuata* Girty ..... 52–53

2a–b. Exterior and interior views of a pedicle valve from loc. Tx. T. 11028. Word formation, Glass Mountains.

3a–b. Lateral and cardinal views of a specimen from loc. 123. In 3b the pedicle valve is up, the brachial down. This specimen is exceptionally small and finely plicated, and the cardinal area is unusually high. Y.P.M. 10908. Leonard formation, Glass Mountains.

4a–d. Pedicle, brachial, lateral, and posterior views of a specimen from loc. 242. In 4d the pedicle valve is up, the brachial down. Y.P.M. 10921.

5a–d. Brachial, pedicle, lateral, and posterior views of a specimen from loc. T. 240a. In 5d the pedicle valve is up, the brachial down. T. 10700.

6a–c. Lateral, posterior, and anterior views of a large cardinal process from loc. Tx, of undetermined relationships, perhaps belonging to a *Meekella*.

7a–d. Pedicle, brachial, lateral, and posterior views of a specimen from loc. Tx. 148. In 7d the pedicle valve is up, the brachial down. T. 10709. Word formation, Glass Mountains.

8–9. *Meekella hessensis*, n. sp. ..... 56

8a–d. Pedicle, brachial, lateral, and posterior views of the holotype, from loc. 122. Y.P.M. 10950.


9a–b. Lateral and pedicle views of a specimen from loc. 222, more coarsely plicate than the holotype. Hess formation, Glass Mountains.

10–12. *Meekella globosa*, n. sp. ..... 54

10. Pedicle valve of a large specimen from loc. 128. T. 10609.

11a–c. Pedicle, lateral, and posterior views of a cardinal process from loc. 128. Y.P.M. 10945.

12a–c. Brachial, lateral, and posterior views of the holotype, from loc. 151. Y.P.M. 10944. Leonard formation, Glass Mountains.


1a
1b
2a
3a
3b
4a
5a
5b
5c
5d
6a
6b
4b
7a
7b
7c
7d
6c
4d
4c
8a
8b
8c
8d
9a
9b
10
11a
11b
11c
12a
12b
12c

## PLATE VI

Figure— PAGE

1–4. *Meekella irregularis texana*, n. var. ..... 55

1a–c. Pedicle, lateral, and posterior views of a pedicle valve from loc. 70.

2a–c. Pedicle, lateral, and posterior views of a pedicle valve from loc. 199, the holotype. Y.P.M. 10966.

3. Brachial valve of a specimen from loc. 95. Y.P.M. 10967.


4a–b. Pedicle and lateral views of a pedicle valve from loc. 88S. Y.P.M. 10968. Wolfcamp formation, Glass Mountains.

5–7. *Meekella grandis*, n. sp. ..... 54–55

5a–c. Lateral, brachial, and posterior views of the holotype, from loc. 174. In 5c the pedicle valve is up, the brachial down. Y.P.M. 10589. Leonard formation, Glass Mountains.

6a–d. Ventral, posterior, inverted, anterior, and lateral views of a cardinal process from loc. 105. Y.P.M. 10890. Hess formation, Glass Mountains.

7. Brachial valve of a specimen from loc. Tg. Leonard formation, Shafter.


1a
1b
1c
2a
2b
2c
3
4a
4b
5a
5b
5c
6a
6b
6c
6d
7

## PLATE VII

Figure— PAGE

1–5. *Meekella mexicana* Girty ..... 55

1a–d. Pedicle, brachial, lateral, and posterior views of a specimen from loc. 434. In 1d the pedicle valve is up, the brachial down. Y.P.M. 10943.

2a–c. Lateral, pedicle, and brachial views of a specimen from loc. 434. Y.P.M. 10943.

3a–b. Brachial and posterior views of a specimen from loc. 434.

4. Posterior view of a specimen from loc. 434.

5. Posterior view of a specimen from loc. 434. Y.P.M. 10943. Middle Gym formation, Glass Mountains.

6–8. *Meekella skenoides* Girty ..... 56

6a–d. Pedicle, brachial, lateral, and posterior views of a specimen from loc. 192. Y.P.M. 11000.

7a–d. Pedicle, lateral, brachial, and posterior views of a specimen from loc. 192. The apical part of the pedicle valve has been reconstructed with clay. In 7d the pedicle valve is up, the brachial down. T. 10748.

8. Fragmentary pedicle valve from loc. 44–46. Y.P.M. 11331. Word formation, Glass Mountains.

9. *Meekella striatocostata* (Cox), *Strophalosia hystricula* Girty, *and Streptorhynchus* sp. ..... 56–57; 96

9a–c. Pedicle, interior, and lateral views of a pedicle valve from loc. 93S. Y.P.M. 11911. Wolfcamp formation, Glass Mountains.


10–13. *Geyerella americana* Girty ..... 57

10a–b. Lateral and pedicle views of a pedicle valve from loc. 85, which appears to belong to this genus but is not typical of the species. Y.P.M. 10999.

11a–b. Lateral and posterior views of a specimen from loc. 35. In 11b the cardinal area is twisted to the left; it is not sharply defined, and therefore not easily distinguishable. Y.P.M. 10995.

12a–b. Lateral view of a large specimen from loc. T. 15, and longitudinal section of same, showing the median septum below, with one of the dental lamellae above joining it at an angle. Apices of both valves broken.

13. Pedicle view of a shell from loc. 17. Y.P.M. 10997. Hess (or possibly Wolfcamp) formation, Glass Mountains.


1a
1b
1c
1d
2a
3a
3b
2b
2c
4
5
6a
7a
7b
8
6d
6c
6b
7d
9a
9b
7c
9c
10a
10b
11a
11b
12a
12b
13

## PLATE VIII

Figure— PAGE

1–2. *Derbya nasuta* Girty ..... 59

1a–d. Pedicle, brachial, lateral, and posterior views of a distorted shell from loc. D. 3. Note the much greater elevation of the cardinal area on one side than on the other. In 1d the pedicle valve is directed upward. Leonard formation, Glass Mountains.

2. Pedicle view, showing septum, of a specimen from loc. 476. Y.P.M. 11430. Hess formation, Sierra Diablo.

3. *Derbya cymbula* Hall and Clarke? ..... 59

3a–b. Posterior and brachial views of a brachial valve from loc. 169a. Note the holes penetrating the shell; they probably were made by some boring organism. Y.P.M. 11409. Wolfcamp formation, Glass Mountains.

4–6. *Derbya buchi* (d'Orbigny) ..... 58


4a–d. Posterior, brachial, pedicle, and lateral views of a specimen from loc. 441. Y.P.M. 11414. Upper Gym formation, Hueco Mountains.

5. Brachial valve from loc. 93. Y.P.M. 11410.

6a–b. Posterior and pedicle views of a pedicle valve from loc. 93. T. 10600. Wolfcamp formation, Glass Mountains.

7. *Derbya elevata*, n. sp. ..... 59

7a–c. Cardinal, pedicle, and lateral views of the holotype from loc. 265. Y.P.M. 11417. Word formation, Glass Mountains.


1a
1b
1c
1d
2
3a
4a
5
3b
4b
6a
4d
4c
6b
7a
7b
7c

## PLATE IX

Figure— PAGE

1–4. *Chonetes permianus* Shumard ..... 62–63

1a–b. Exterior and interior brachial views of a specimen from loc. 6. Y.P.M. 10811.

2a–b. Exterior and interior pedicle views of a specimen from loc. 6. Y.P.M. 10811. Word formation (basal limestone), Glass Mountains.

3. Pedicle view of a specimen from loc. 484. Y.P.M. 10817. Bone Canyon limestone, Sierra Diablo.

4. Pedicle view of a specimen from loc. 128. T. 10615. Leonard formation, Glass Mountains.

5. *Chonetes verneuilianus* Norwood and Pratten ..... 64

5a–b. Pedicle and brachial views of a specimen from loc. 75. Y.P.M. 10792. Wolfcamp formation, Glass Mountains.

6. *Chonetes hillanus* Girty ..... 61

6a–b. Internal mould of a pedicle valve, from loc. 248, and posterior view of the same. Word formation (upper limestone), Glass Mountains. Y.P.M. 10818.

7–11. *Chonetes quadratus,* n. sp. ..... 63

7. Pedicle view of a specimen from loc. 246. Word formation, Glass Mountains. T. 10660.

8. Posterior view of a paratype from Bass Ranch, Grand Canyon. Y.P.M. 10830.

9. Pedicle view of the holotype from Bass Ranch, Kaibab formation, Grand Canyon. Y.P.M. 10830.

10. Pedicle view of a paratype from loc. 247. Y.P.M. 10833.

11. Pedicle view of a specimen from loc. T. 148. T. 10635. Word formation, Glass Mountains.

12–13. *Chonetes biplicatus,* n. sp. ..... 60

12a–c. Pedicle, lateral, and posterior views of the holotype from loc. 476. Y.P.M. 10824. Hess formation, Sierra Diablo.

13. Pedicle view of a specimen from loc. 193. T. 10686. Wolfcamp formation, Glass Mountains.

14. *Chonetes granulifer* Owen ..... 60

Pedicle view of a specimen from loc. 88. Y.P.M. 10796. Word formation, *Uddenites* zone, Glass Mountains.

15–16. *Chonetes spinoliratus,* n. sp. ..... 63

15a–c. Lateral, pedicle (×3), and posterior (×3) views of the holotype, from loc. 91a. Y.P.M. 10825.

16a–b. Pedicle and brachial views of a specimen from loc. 92. T. 10588. Wolfcamp formation, Glass Mountains.

17–20. *Chonetes deliciasensis,* n. sp. ..... 60

17a–d. Pedicle, brachial, lateral, and posterior views of a cotype from section 2, bed 12. Y.P.M. 10787.

18. Pedicle view of an internal mold from section 2, bed 12. Y.P.M. 10787.

19a–c. Posterior, pedicle, and brachial views of a cotype from section 2, bed 12. Y.P.M. 10787.

20. Exterior of pedicle valve of a cotype from section 1, bed 11. Y.P.M. 10789. Limestone of Cerro Caballo, near Las Delicias.

21–22. *Chonetes hessensis,* n. sp. ..... 61

21a–c. Pedicle, brachial, and posterior views of a cotype from loc. 107. T. 10589.

22a–c. Lateral, pedicle (×3), and brachial (×3) views of a cotype from loc. 107. Y.P.M. 10820. Hess formation, Glass Mountains.

23. *Chonetes spinoliratus diabloensis,* n. var. ..... 63

23a–b. Brachial and pedicle views of the holotype, from loc. 479. Y.P.M. 10829. Hess formation, Sierra Diablo.

24. *Chonetes ostiolatus* Girty ..... 62

Pedicle view of a specimen from loc. Tn. T. 10590. Cibolo limestone.

25. *Chonetes subliratus* Girty ..... 63–64

25a–b. Pedicle and posterior views of a pedicle valve of rather large size from loc. T. 239. T. 10730. Word formation, Glass Mountains.


26. *Chonetes molengraaffi* Broili? ..... 61

Pedicle fragment from loc. 512. Y.P.M. 10819. Leonard formation, Finlay Mountains.

27–28. *Orthotetella wolfcampensis,* n. gen. n. sp. ..... 51–52

27a–b. Pedicle and posterior views of a pedicle valve from loc. 93. T. 10446.

28a–c. Posterior, pedicle, and pedicle interior views of the holotype, from loc. 93. Y.P.M. 11366. Wolfcamp formation, Glass Mountains.


1a
1b
2a
2b
5a
5b
8
12a
12b
3
4
7
10
12c
14
6b
9
11
17a
17d
15a
16a
6b
13
17c
17b
20
19b
15b
15c
19a
18
19c
21a
21b
21c
22a
24
23a
26
22b
22c
23b
25a
25b
27b
27a
28a
28b
28c

## PLATE X

Figure— PAGE

1–7. *Chonetes subliratus* Girty ..... 63–64

1a–b. Pedicle and brachial views of a shell from loc. 123, showing the inner shell surface. T. 10733.

2. Pedicle view of a small specimen from loc. 123, inner surface shown. Y.P.M. 10839. Leonard formation, Glass Mountains.

3. Interior of a pedicle valve from loc. 46, showing only the outer smooth shell layer. Y.P.M. 10856.

4a–b. Posterior and pedicle views of a specimen from loc. 250, showing outer smooth shell layer. Y.P.M. 10868.

5. Pedicle valve from loc. 253, the surface of which is formed by the outer smooth layer, showing the penetrating holes. T. 10742. Word formation, Glass Mountains.

6. Pedicle valve from loc. 512. Y.P.M. 10847. Leonard formation, Finlay Mountains.

7a–c. Pedicle, brachial, and posterior views of a specimen from loc. 123. Y.P.M. 10839. Leonard formation, Glass Mountains.

8. *Isogramma millepunctata* (Meek and Worthen) ..... 65

Part of a pedicle valve (internal mold) from loc. 203. Y.P.M. 10837: Wolfcamp formation, Glass Mountains.

9–11. *Productus guadalupensis* Girty ..... 67

9a–c. Pedicle, anterior, and lateral views of a specimen from loc. 246. Y.P.M. 10743.

10a–c. Pedicle, lateral, and posterior views of a specimen from loc. 246. Y.P.M. 10743.

11. Lateral view of a specimen from loc. T. 148. Upper Word formation, Glass Mountains.

12. *Productus dartoni*, n. sp. ..... 67

12a–c. Pedicle, lateral, and posterior views of a cotype from loc. 385. Y.P.M. 10740. Lower Gym formation, Hueco Mountains.

13. *Productus dar*toni, n. sp., and *Composita mira* Girty ..... 67, 129

13a–c. Posterior, anterior, and lateral views of a cotype from loc. 460. Y.P.M. 10739. Upper Wolfcamp formation, Sierra Diablo.

14. *Productus dartoni sullivanensis*, n. var. ..... 67

14a–c. Pedicle, posterior, and lateral views of a specimen from loc. 46. Y.P.M. 10767. Lower Word formation, Glass Mountains.


15. *Productus capitanensis* Girty ..... 66–67

15a–b. Pedicle and lateral views of a specimen from loc. 179. Y.P.M. 10773. Gilliam member, Glass Mountains.

16–17. *Productus arcticus* Whitfield ..... 66

16a–c. Pedicle, lateral, and brachial views of a specimen from loc. 247. Y.P.M. 10769.

17a–c. Posterior, pedicle, and lateral views of a specimen from loc. 247. Y.P.M. 10769. Upper Word formation, Glass Mountains.


1a
1b
2
3
4a
5
6
7a
7b
7c
4b
8
9b
9a
9c
10a
10b
12a
12b
12c
10c
11
13a
13b
13c
14a
14b
14c
15a
15b
16a
16b
17a
17b
17c
16c

## PLATE XI


Figure— PAGE

1. *Productus comancheanus* Girty ..... 67
   Pedicle view of a specimen from loc. 144. Y.P.M. 10745. Middle Word formation, Glass Mountains.

2. *Productus inca* d'Orbigny ..... 68–69
   2a–c. Posterior, pedicle, and lateral views of a specimen from loc. 421. Y.P.M. 10644. Middle Gym formation, Hueco Mountains.

3–6. *Productus hessensis*, n. sp. ..... 68
   3a–c. Pedicle, lateral, and brachial views of a specimen from loc. 507. Y.P.M. 10654.
   4a–b. Lateral and pedicle views of a specimen from loc. Tzz,aa. T. 10716. Gym formation, Wylie Mountains.
   5. Pedicle view of a specimen from loc. 222, only the visceral part of which remains. Y.P.M. 10655.
   6a–b. Lateral and pedicle views of the holotype, from loc. 106. Y.P.M. 10651. Hess formation, Glass Mountains.

7–8. *Productus huecoensis*, n. sp. ..... 68
   7a–c. Posterior, lateral, and pedicle views of the holotype, from loc. 380. Y.P.M. 10554. Lower Gym formation, Hueco Mountains.
   8a–b. Posterior and lateral views of a specimen from loc. 193, which is considerably smaller than those from the Hueco Mountains. Upper Wolfcamp formation, Glass Mountains.


1
2a
2b
2c
3a
3b
4a
4b
3c
6a
6b
5
7a
8a
7b
7c
8b

## PLATE XII

Figure— PAGE

1–6. *Productus ivesi* Newberry ..... 69–70

1a–b. Posterior and pedicle views of a specimen from loc. 174. Y.P.M. 10608.


2. Pedicle view of a specimen from loc. 123.

3. Interior of brachial valve from loc. T. 128. T. 9980.

4. Pedicle view of a specimen from loc. 174. T. 10049.

5a–d. a, c, d, lateral, brachial, and pedicle views of a specimen from loc. 123; b, posterior view of another specimen from the same locality. Y.P.M. 10629.

6. Interior of a very thick old pedicle valve from loc. 15. Y.P.M. 10593. Leonard formation, Glass Mountains.


1a
1b
2
3
4
5a
5b
6
5c
5d

## PLATE XIII

Figure— PAGE

1–3. *Productus multistriatus* Meek ..... 71

1. Pedicle view of a specimen from loc. T. 240. The anterior trail of the shell is not preserved. T. 10036.

2a–b. Lateral and pedicle views of a specimen from loc. 144. The anterior trail of the shell is not preserved. Y.P.M. 10647. Middle Word formation, Glass Mountains.

3a–c. Pedicle, lateral, and anterior views of a specimen from section 1, bed 14, showing the smaller size of the Mexican specimens. Y.P.M. 10648. Limestone of Cerro Caballo, Las Delicias, Coahuila.

4–5. *Productus inca* d'Orbigny ..... 68–69

4a–b. Pedicle and lateral views of a specimen from loc. 414. T. 10432.

5a–c. Pedicle, lateral, and brachial views of a specimen from loc. 415. Y.P.M. 10645. Middle Gym formation, Hueco Mountains.


6–9. *Productus indicus* Waagen ..... 72–73

6a–c. Pedicle, posterior, and lateral views of a specimen from loc. 46. Y.P.M. 10559. Word formation, Glass Mountains.

7. Pedicle view of a specimen from loc. 517. Y.P.M. 10562. Leonard formation, Shafter.

8. Part of pedicle valve from loc. T. 251. T. 10254.

9a–c. Posterior, lateral, and pedicle views of a specimen from loc. 46. T. 10250. Word formation, Glass Mountains.


1
2a
2b
3a
3b
3c
4a
4b
5a
5b
5c
6a
6b
6c
7
8
9a
9b
9c

## PLATE XIV

Figure— PAGE

1–3. *Productus gratiosus occidentalis* Schellwien ..... 71

1a–d. Pedicle, brachial, lateral, and posterior views of a specimen from loc. 93. Y.P.M. 10768.

2. Pedicle view of a specimen from loc. 93. T. 10307. Wolfcamp formation, Glass Mountains.

3a–c. Pedicle, lateral, and posterior views of a specimen from loc. Sd. Y.P.M. 12664. Gaptank formation, Marathon Basin.

4–9. *Productus leonardensis*, n. sp. ..... 70–71

4a–b. Brachial and pedicle views of a shell from loc. 128. Y.P.M. 10712.

5. Lateral view of a cotype from loc. 120. Y.P.M. 10706.

6a–c. Lateral, anterior, and pedicle views of a pedicle valve from loc. Tb. T. 10337.

7. External mold of a dorsal valve from loc. 10. Y.P.M. 10715.

8. Produced anterior part of a shell from loc. 128. Y.P.M. 10712.

9. Part of a pedicle valve from loc. 128, showing spines along posterior margin, and upturned marginal part of shell. Y.P.M. 10712. Leonard formation, Glass Mountains.

10–14. *Productus occidentalis* Newberry ..... 72–73

10. Interior of brachial valve from loc. 123, showing brachial ridges. Y.P.M. 10566.

11. Pedicle view of a specimen from loc. 174.

12. Pedicle view of a specimen from loc. 174. Y.P.M. 10565.

13. Brachial view of a large, broad specimen from loc. 128, showing spine projecting from one of the ears. Y.P.M. 10563. Leonard formation, Glass Mountains.


14a–b. Pedicle and lateral views of a specimen from loc. 247. T. 10319. Upper Word formation, Glass Mountains.

15–17. *Productus semireticulatus hermosanus* Girty? ..... 73–74

15. Mold of brachial valve from loc. 88S. Y.P.M. 10698.

16. Part of a pedicle valve from loc. 87. T. 10288.

17. Part of a pedicle valve from loc. 195. Y.P.M. 10694. Wolfcamp formation, Glass Mountains.


1a
1b
1c
1d
2
3a
3b
3c
4a
4b
5
6a
7
8
6b
6c
9
10
11
12
13
14a
14b
15
16
17

## PLATE XV

Figure— PAGE

1–5. *Productus schucherti*, n. sp. ..... 73

1a–c. Posterior, lateral, and brachial views of a specimen from loc. 484. Y.P.M. 10765.

2a–c. Posterior, pedicle, and lateral views of a specimen from loc. 484. T. 10311. Bone Canyon limestone, Sierra Diablo.

3a–b. Brachial and pedicle views of a cotype from loc. 120. Y.P.M. 10763.

4a–c. Lateral and pedicle views of a cotype from loc. 120. Y.P.M. 10763.

5a–b. Posterior and pedicle views of a specimen from loc. D1. Leonard formation, Glass Mountains.

6–11. *Productus semistriatus* Meek ..... 74

6a–c. Anterior, posterior, and lateral views of a specimen from loc. 92. Y.P.M. 10667. Wolfcamp formation, Glass Mountains.


7a–b. Lateral and brachial views of a specimen from loc. 376. T. 10286. Lower Gym formation, Hueco Mountains.

8a–c. Lateral, posterior, and pedicle views of an exceptionally broad specimen from loc. 88S. Y.P.M. 10675. Wolfcamp formation, *Uddenites* zone, Glass Mountains.

9a–b. Posterior and pedicle views of a specimen from loc. 456. Y.P.M. 10666. Upper Wolfcamp formation, Sierra Diablo.

10a–b. Lateral and pedicle views of a specimen from loc. 382. T. 10276. Lower Gym formation, Hueco Mountains.

11. Pedicle view of a specimen from loc. 93S. Y.P.M. 10669. Wolfcamp formation, Glass Mountains.


1a
1b
1c
4a
2a
3a
4b
3b
4c
2b
2c
5a
5b
6b
6a
6c
7a
8a
9a
10a
7b
8b
9b
10b
11
8c

## PLATE XVI

Figure— PAGE

1–5. *Productus wolfcampensis,* n. sp. ..... 74–75

1a–b. Pedicle and lateral views of a specimen from loc. 460. Y.P.M. 10663. Upper Wolfcamp formation, Sierra Diablo.

2a–b. Posterior and lateral views of a specimen from loc. 198. T. 10343. Wolfcamp formation, Glass Mountains.

3a–c. Pedicle, lateral, and posterior views of the holotype, from loc. Tmm. T. 10339. Lower Gym formation, Hueco Mountains.

4a–b. Pedicle and lateral views of a specimen from loc. 456. Y.P.M. 10662.

5. Pedicle view of an exceptionally coarsely costate specimen from loc. 456. Y.P.M. 10662. Upper Wolfcamp formation, Sierra Diablo.

6–7. *Linoproductus cora* (d'Orbigny) ..... 75–76


6a–c. Brachial, pedicle, and lateral views of a specimen from loc. 434. Y.P.M. 11529.

7a–b. Pedicle and posterior views of a specimen from loc. 434. Y.P.M. 11529. Middle Gym formation, Hueco Mountains.

8–9. *Linoproductus cora angustus,* n. var. ..... 76

8a–b. Lateral and pedicle views of a specimen from loc. 15. T. 10057. Leonard formation, Glass Mountains.

9. Pedicle view of a cotype from loc. 105. Y.P.M. 11519. Hess formation, Glass, Mountains.


1a
1b
2a
2b
3a
3b
3c
4a
4b
5
6a
6b
6c
7a
7b
8a
8b
9

## PLATE XVII

Figure— PAGE

1–2. *Linoproductus (Cancrinella) villiersi* (d'Orbigny) ..... 78

1a–b. Pedicle and lateral views of a specimen from loc. 414. Y.P.M. 10780.

2a–b. Pedicle and lateral views of a specimen from loc. 414. Y.P.M. 10780. Middle Gym formation, Hueco Mountains.

3–5. *Linoproductus girtyi*, n. sp. ..... 76

3. Pedicle view of an incomplete specimen from loc. 23–24. Y.P.M. 11501. Leonard formation, Glass Mountains.

4. Pedicle view of an incomplete specimen from loc. T. 240. T. 10681. Word formation, Glass Mountains.

5a–b. Pedicle and posterior view of the holotype, from loc. Ax. Y.P.M. 11507. Leonard formation, Glass Mountains.

6–7. *Linoproductus (Cancrinella) phosphaticus* (Girty) ..... 77–78

6. Lateral view of a specimen from loc. 247. Y.P.M. 10775.

7. Pedicle view of a specimen from loc. 247. Upper Word formation, Glass Mountains. T. 10584.

8–9. *Linoproductus nasutus*, n. sp. ..... 76

8a–c. Pedicle, lateral, and posterior views of a paratype from loc. 247. T. 10096.

9a–b. Pedicle and lateral views of the holotype, from loc. 246. Y.P.M. 11523. Upper Word formation, Glass Mountains.

10–15. *Linoproductus waagenianus* (Girty) ..... 77

10a–c. Pedicle, brachial, and lateral views of a specimen from loc. T. 119. T. 10329. Leonard formation, Glass Mountains.

11a–c. Pedicle, posterior, and lateral views of a specimen from loc. 46, showing broad lateral extension of ears. T. 10330.

12. View of three specimens that are attached to one another, from loc. 46. Y.P.M. 11482.

13. Brachial view of a specimen from loc. 46, showing partial envelopment of brachial side by enrollment of pedicle valve. Y.P.M. 11482.

14a–c. Pedicle, posterior, and brachial views of a specimen from loc. 46. Y.P.M. 11482.

15a–b. Pedicle and posterior views of a specimen from loc. 46. T. 10330. Lower Word limestone, Glass Mountains.

16–19. *Striatifera pinniformis* (Girty) ..... 78

16a–b. Pedicle and lateral views of a specimen from loc. 120. Y.P.M. 11634.

17a–b. Posterior and pedicle views of a specimen from loc. 21.


18. Lateral view of a specimen from loc. 37. Y.P.M. 11631. Leonard formation, Glass Mountains.

19a–c. Posterior, pedicle, and brachial views of a specimen from section 1, bed 14. Limestone of Cerro Caballo, near Las Delicias, Coahuila. Y.P.M. 11630.

20–21. *Buxtonia occidentalis*, n. sp. ..... 79–80

20a–b. Pedicle and brachial views of a cotype from loc. 93. Y.P.M. 11528.

21a–b. Pedicle and lateral views of a cotype from loc. 91a. Y.P.M. 11552. Wolfcamp formation, Glass Mountains.


1a
1b
2a
2b
3
4
5a
5b
6
7
8a
8b
8c
9a
9b
10a
10b
10c
11c
11a
11b
12
13
14a
14b
14c
15a
15b
16a
16b
17a
18
17b
19a
19b
19c
20a
20b
21a
21b

## PLATE XVIII


Figure— PAGE

1–3. *Buxtonia peruviana* (d'Orbigny) ..... 80

1a–d. Pedicle, brachial, lateral, and posterior views of a specimen from loc. Tmm. Lower Gym formation, Hueco Mountains.

2a–d. Posterior, lateral, brachial, and pedicle views of a specimen from loc. 472. Y.P.M. 11620. Hess formation, Sierra Diablo.

3a–b. Lateral and pedicle views of a specimen from loc. 377b. Y.P.M. 11619. Lower Gym formation, Hueco Mountains.


1a
1b
1c
1d
2a
2b
2c
2d
3a
3b

## PLATE XIX

Figure— PAGE

1. *Buxtonia victorioensis*, n. sp. ..... 80

1a–c. Posterior, pedicle, and lateral views of the holotype, from loc. 498. Y.P.M. 11615. Victorio Peak limestone, Sierra Diablo.

2–4. *Waagenoconcha leonardensis*, n. sp. ..... 80–81

2a–b. Pedicle and lateral views of a specimen from loc. Tb. T. 10345.


3a–b. Lateral and posterior views of the holotype, from loc. 120; only one-half of the shell is preserved. Y.P.M. 11614.

4a–c. Pedicle, lateral, and posterior views of a specimen from loc. 15. Y.P.M. 11613. Leonard formation, Glass Mountains.

5–6. *Waagenoconcha montpelierensis* (Girty) ..... 81

5. Posterior view of a specimen from loc. T. 240. T. 10265.

6a–d. Posterior, pedicle, lateral, and brachial views of a specimen from loc. 146. Y.P.M. 11674. Word formation, Glass Mountains.


1a
1b
1c
2a
2b
3a
4a
4b
3b
4c
5
6a
6b
6c
6d

## PLATE XX

Figure— PAGE

1–3. *Overtonia cristato-tuberculata* (Kozlowski) ..... 82

1. Pedicle view of a specimen from loc. Tba. T. 9994.

2a–b. Pedicle and lateral views of a specimen from loc. 94. Y.P.M. 11499.

3a. Pedicle view of a specimen from loc. 94. Y.P.M. 11499. Wolfcamp formation, *Uddenites* zone, Glass Mountains.

4–5. *Avonia meekana* (Girty) ..... 83

4a–b. Pedicle and posterior views of a specimen from loc. 193. T. 10253.

5a–b. Pedicle and lateral views of a specimen from loc. 144. Y.P.M. 11530. Middle Word formation, Glass Mountains.

6. *Avonia ? incurvata*, n. sp. ..... 83

6a–c. Pedicle, lateral, and posterior views of the holotype, from loc. 46. Y.P.M. 11583. Lower Word formation, Glass Mountains.

7–9. *Avonia boulei* (Kozlowski) ..... 82–83

7a–c. Pedicle, lateral, and posterior views of a specimen from loc. 193. T. 10375.

8a–c. Pedicle, lateral, and posterior views of a specimen from loc. 193. Y.P.M. 11585.

9a–c. Pedicle, posterior, and brachial views of a specimen from loc. 87. Y.P.M. 11589. Wolfcamp formation, Glass Mountains.

10–11. *Avonia subhorrida* (Meek) ..... 84

10a–b. Pedicle and lateral views of a specimen from loc. Tx. 148. T. 10283.

11. Pedicle view of a specimen from loc. T. 239. T. 10255. Middle and upper Word formation, Glass Mountains.

12–15. *Avonia subhorrida rugatula* (Girty) ..... 84–85

12a–b. Pedicle and lateral views of a specimen from loc. 440a. Y.P.M. 11608. Upper Gym formation, Hueco Mountains.

13. Lateral view of a specimen from loc. 137. T. 10256. Middle Word formation, Glass Mountains.

14. Pedicle view of a specimen from loc. 123. Y.P.M. 11607. Leonard formation, Glass Mountains.

15. Pedicle view of a specimen from loc. 137. Word formation, Glass Mountains.

16–24. *Avonia signata* (Girty) ..... 83

16a–b. Pedicle and lateral views of a specimen from loc. 144, showing the large hollow spines on the ears. Y.P.M. 11531.

17a–b. Pedicle and lateral views of a specimen from loc. 144. T. 10235.

18. Pedicle view of a specimen from loc. 148.

19. Lateral view of a specimen from loc. 253. Y.P.M. 11545.


20. Interior of brachial valve from loc. Tx. T. 11070.

21a–d. Brachial, lateral, posterior, and pedicle views of a specimen from loc. 247. Y.P.M. 11543.

22a–b. Pedicle and posterior views of a specimen from loc. 247. Y.P.M. 11543.

23a–b. Posterior and pedicle views of a specimen from loc. T. 148. T. 10232.

24a–b. Pedicle and lateral views of a specimen from loc. 246. Y.P.M. 11544. Middle and upper Word formation, Glass Mountains.


1
2a
2b
4a
4b
5a
5b
6a
6b
6c
7a
7b
7c
8a
8b
8c
9a
9b
9c
10a
10b
11
12a
13
14
12b
17b
15
16a
16b
17a
21a
21b
20
18
19
21c
21d
23a
22a
22b
23b
24a
24b

## PLATE XXI

Figure— PAGE

1–5. *Avonia walcottiana* (Girty) ..... 85

1. Pedicle view of a specimen from loc. x. T. 11039.

2a–b. Pedicle and posterior views of a specimen from loc. 248. Y.P.M. 11571.

3a–b. Pedicle and lateral views of a specimen from loc. 144. T. 11313.

4. Pedicle view of a specimen from loc. 240. Y.P.M. 11576.

5a–c. Pedicle, lateral, and posterior views of a specimen from loc. 144. Y.P.M. 11575. Word formation, Glass Mountains.

6–7. *Marginifera dugoutensis*, n. sp. ..... 87

6. Pedicle view of a cotype from loc. 193. Y.P.M. 11773.

7a–b. Posterior and pedicle views of a cotype from loc. 193. Y.P.M. 11773. Wolfcamp formation, Glass Mountains.

8–13. *Avonia walcottiana costata*, n. var. ..... 85

8. Pedicle view of a cotype from loc. T. 240. T. 10282.

9a–b. Pedicle and lateral views of a cotype from loc. T. 240. T. 10282.

10a–b. Pedicle and posterior views of a cotype from loc. T. 240. T. 10282.

11. Pedicle view of a cotype from loc. T. 240. T. 10282.

12a–b. Pedicle and lateral views of a cotype from loc. T. 240. T. 10282.

13. Lateral view of a specimen from loc. 241. Y.P.M. 11568. Word formation, Glass Mountains.

14–18. *Marginifera capaci* (d'Orbigny) ..... 86

14a–c. Pedicle, brachial, and lateral views of a specimen from loc. 439. Y.P.M. 11772. Upper Gym formation, Hueco Mountains.

15a–c. Pedicle, posterior, and lateral views of a specimen from loc. 376. Y.P.M. 11770.

16a–b. Pedicle and posterior views of a specimen from loc. 384. Lower Gym formation, Hueco Mountains.

17. Pedicle view of a specimen from loc. 196. Y.P.M. 11763.

18a–b. Pedicle and posterior views of a specimen from loc. 88. Wolfcamp formation, Glass Mountains.

19–21. *Marginifera cristobalensis* Girty ..... 86–87

19a–b. Lateral and posterior views of a specimen from loc. 120. T. 10188.

20a–c. Pedicle, lateral, and posterior views of a specimen from loc. 120. Y.P.M. 11641.

21. Pedicle view of a specimen from loc. 125. Y.P.M. 11662. Leonard formation, Glass Mountains.

22–24. *Marginifera manzanica* Girty ..... 87


22a–c. Pedicle, brachial, and posterior views of a specimen from loc. 4. Y.P.M. 11687.

23. Posterior view of a specimen from loc. 4. Y.P.M. 11687.

24. Pedicle view of a specimen from loc. 4. T. 10076. Leonard formation, Glass Mountains.

25. *Horridonia texana*, n. sp., with *Agathiceras girtyi* Böse ..... 85–86

25a–c. Pedicle, posterior, and lateral views of the holotype, which is broken in the central part, showing a specimen of *Agathiceras girtyi*. Loc. 255. Y.P.M. 11508. Middle Word formation, Glass Mountains.


1
2a
2b
3a
3b
7a
4
5a
5b
5c
6
7b
9a
9b
10a
10b
11
12a
12b
14a
14b
14c
15a
15b
15c
13
16a
16b
17
18a
18b
19a
19b
20a
20b
20c
21
22a
22b
22c
23
24
25a
25b
25c

## PLATE XXII

Figure— PAGE

1–2. *Marginifera popei minor,* n. var. 88–89

1a–c. Pedicle, posterior, and lateral views of a cotype from section 1, bed 14. Y.P.M. 11635.

2. Pedicle view of a cotype from section 1, bed 14. Y.P.M. 11635. Limestone of Cerro Caballo, near Las Delicias, Coahuila.

3–10. *Marginifera reticulata,* n. sp. 89

3. Brachial view of a specimen from loc. 4. Y.P.M. 11724.

4. Pedicle view of a specimen from loc. 226. Y.P.M. 11727.

5a–b. Lateral and pedicle views of a specimen from loc. 228. Y.P.M. 11729.

6a–c. Pedicle, posterior, and lateral views of a cotype from loc. Tb. T. 10056.

7a–c. Lateral, pedicle, and posterior views of a specimen from loc. Tb. T. 10056.

8. Pedicle view of a specimen from loc. D. 4.

9a–b. Pedicle and lateral views of a specimen from loc. Tb. T. 10056.

10a–b. Pedicle and lateral views of a specimen from loc. Tb. T. 10056. Leonard formation, Glass Mountains.

11–15. *Marginifera reticulata angusta,* n. var. 89

11. Pedicle view of a specimen from loc. 4. Y.P.M. 11807.

12. Pedicle view of a cotype from loc. 26S. Y.P.M. 11806.

13a–b. Pedicle and lateral views of a cotype from loc. T. 15. T. 10290.

14a–b. Pedicle and posterior views of a cotype from loc. 26S. Y.P.M. 11806.

15. Pedicle view of a specimen from loc. 15. T. 10290. Leonard formation, Glass Mountains.

16–19. *Marginifera popei* (Girty) 88

16. Pedicle view of a specimen from loc. 144. T. 10354.

17. Pedicle view of a specimen from loc. 247.

18. Pedicle view of a specimen from loc. 246. Y.P.M. 11753.

19. Pedicle view of a specimen from loc. 246. Y.P.M. 11753. Word formation, Glass Mountains.

20–25. *Marginifera opima* (Girty) 87–88

20. Interior view of a brachial valve from loc. Tx, showing incomplete marginal ridge. T. 11072.


21. Lateral view of a specimen from loc. 247. Y.P.M. 11707.

22a–c. Brachial, pedicle, and lateral views of a specimen from loc. 253. Y.P.M. 11718.

23a–c. Pedicle, lateral, and posterior views of a specimen from loc. 247.

24a–b. Pedicle and posterior views of a specimen from loc. 144. Y.P.M. 11706.

25a–b. Lateral and pedicle views of a specimen from loc. 247. Y.P.M. 11707. Word formation, Glass Mountains.


1a
1b
1c
2
3
4
5
6a
6b
6c
7a
8
5b
10a
10b
7b
7c
9a
11
12
13a
13b
14a
9b
16
17
18
14b
15
19
20
21
22a
23a
23b
23c
22b
22c
24a
24b
25a
25b

## PLATE XXIII

Figure— PAGE

1–3. *Marginifera? whitei,* n. sp. 90

1a–c. Pedicle, brachial, and lateral views of the holotype, from loc. 107. Y.P.M. 11685. Hess formation, Glass Mountains.

2a–c. Pedicle, lateral, and posterior views of a specimen from loc. 479. Y.P.M. 11686.

3a–b. Pedicle and posterior views of a specimen from loc. 479. Y.P.M. 11686. Hess formation, Sierra Diablo.

4. *Marginifera wabashensis* (Norwood and Pratten) 90

4a–c. Pedicle, lateral, and posterior views of a specimen from loc. 201. Y.P.M. 11778. Wolfcamp formation, *Uddenites* zone, Glass Mountains.

5–7. *Marginifera? texana* (Girty) 89–90

5. Pedicle view of a specimen from loc. 247. Y.P.M. 11733.

6. Pedicle view of a specimen from loc. 247. Y.P.M. 11733.

7. Pedicle view of a specimen from loc. 247. Y.P.M. 11733. Upper Word formation, Glass Mountains.

8–11. *Marginifera? wordensis,* n. sp. 90–91

8. Pedicle view of a specimen from loc. Tx. T. 9991.

9. Pedicle view of a specimen from loc. Tx. T. 9991.

10. Pedidle view of a cotype from loc. T. 240. T. 9998.

11. Pedicle view of a cotype from loc. T. 240. T. 9998. Word formation, Glass Mountains.

12. *Marginifera?* sp.

12a–c. Pedicle, brachial, and lateral views of a specimen of doubtful relationships from loc. Tb. T. 10987. Leonard formation, Glass Mountains.

13–19. *Marginifera sublaevis,* n. sp. 89

13a–c. Pedicle, posterior, and lateral views of a specimen from loc. 512. Y.P.M. 11802. Leonard formation, Finlay Mountains.

14a–c. Pedicle, posterior, and lateral views of a specimen from loc. 485. Y.P.M. 11796. Bone Canyon limestone, Sierra Diablo.

15a–c. Pedicle, lateral, and posterior views of a cotype from loc. 120. Y.P.M. 11792.

16a–b. Pedicle and lateral views of a specimen from loc. T. 119. T. 10051.

17. Interior of a brachial valve from loc. 241, supposed to belong to this species. Y.P.M. 11797.

18a–b. Brachial and lateral views of a specimen from loc. T. 9. T. 9954.

19. Pedicle view of a cotype from loc. 120. Y.P.M. 11792. Leonard formation, Glass Mountains.

20–25. *Scacchinella gigantea* Schellwien 91–92

20a-b. Pedicle and brachial views of a specimen from loc. 208. The apex of the pedicle valve has been broken along the lines of weakness formed by the transverse partitions. T. 10239.


21a–c. Lateral, pedicle, and brachial views of a specimen from loc. 208, the apex of which is broken, but not along transverse partitions. Y.P.M. 12826.

22a–b. Posterior and brachial views of a small specimen from loc. 208. Y.P.M. 12826.

23. Posterior view of a specimen from loc. 208. Y.P.M. 12826. Hess formation, Glass Mountains.

24a–c. Posterior, lateral, and pedicle views of a pedicle valve from loc. 208. The apex is broken along a transverse partition. T. 10239.

25a–b. Posterior and brachial view of a specimen from loc. 208. The apex of the pedicle valve is broken off. Y.P.M. 12826. Hess formation, Glass Mountains.


1a
1b
1c
2a
2b
2c
3a
3b
4a
4b
4c
5
6
7
8
9
10
11
12a
12b
12c
13a
13b
13c
14a
14b
14c
15a
15b
15c
16a
16b
17
18a
18b
19
20a
20b
21a
21b
21c
22a
22b
23
24a
24b
24c
25a
25b

## PLATE XXIV

Figure— PAGE

1–5. *Scacchinella gigantea* Schellwien ..... 91–92


1a-c. Brachial, pedicle, and posterior views of a part of a long pedicle valve from loc. 35. It can be seen that the fragment is broken at both ends along the transverse partitions. Y.P.M. 12828. Hess formation, Glass Mountains.

2a–b. Posterior and brachial views of a large specimen from loc. 71, the apical part of which is not preserved. The ventral median septum may well be seen. Y.P.M. 12827. Wolfcamp formation, Glass Mountains.

3a–b. Lateral and posterior views of an elongate pedicle valve from loc. 35, showing a transverse angulation in the cardinal area. Y.P.M. 12828.

4a–b. Posterior and lateral views of a large specimen from loc. 208, showing the entire apex of the pedicle valve. Y.P.M. 12826.

5. Posterior view of a specimen from loc. 208. The apex of the pedicle valve is broken along the transverse partitions. A median depression may be seen in the cardinal area, which is due to the exfoliation of the surface of the area and excavation along the inside of the median septum. T. 10239. Hess formation, Glass Mountains.


1a
1b
1c
2a
3a
3b
2b
4a
4b
5

## PLATE XXV

Figure— PAGE

1–4. *Aulosteges magnicostatus* Girty ..... 93

1a–b. Pedicle and lateral views of a specimen from loc. 123. T. 10085.

2a–b. Pedicle and lateral views of a specimen from loc. 26S. Y.P.M. 11890.

3. Lateral view of a specimen from loc. 123. Y.P.M. 11902.

4a–b. Pedicle and lateral views of a specimen from loc. 228. T. 10096. Leonard formation, Glass Mountains.

5–7. *Aulosteges subcostatus*, n. sp. ..... 94

5a–b. Exterior and interior of a pedicle valve from loc. T. 119. T. 10155.

6a–d. Pedicle, brachial, posterior, and lateral views of a specimen from loc. T. 119. T. 10155.

7a–b. Brachial and lateral views of the holotype, from loc. 123. Y.P.M. 11837. Leonard formation, Glass Mountains.

8–11. *Aulosteges guadalupensis* Shumard ..... 93

8a–b. Pedicle and lateral views of a specimen from loc. 144. Y.P.M. 11824.

9. Pedicle valve from loc. 144. P.Y.M. 11824.

10. Pedicle valve from loc. Tx.

11. Lateral view of a pedicle valve from loc. 148. T. 10259. Word formation, Glass Mountains.

12–15. *Aulosteges beedei*, n. sp. ..... 92–93

12a–b. Brachial and lateral views of a cotype from loc. 123. Y.P.M. 11876.

13a–c. Pedicle, brachial, and lateral views of a cotype from loc. 123. Y.P.M. 11876.

14a–c. Pedicle, brachial, and lateral views of a cotype from loc. 123. T. 10690. Leonard formation, Glass Mountains.


15a–b. Exterior and interior of a brachial valve from loc. Tx, showing the remarkable cardinal process. Word formation, Glass Mountains. T. 11045.

16–18. *Scacchinella gigantea* Schellwien ..... 91–92

16. Part of a pedicle valve from loc. 208, which is broken along transverse partitions. Y.P.M. 12826.

17. Fragment of a brachial valve from loc. 35, showing spinose surface. T. 10241.

18a–b. Posterior and lateral views of a specimen from loc. 208, broken at the apex along transverse partitions, which show especially well in 18b. Y.P.M. 12826. Hess formation, Glass Mountains.


1a
1b
2a
2b
3
4a
4b
5a
5b
6a
6b
7a
7b
6c
6d
8a
8b
9
12a
13a
13b
10
11
13c
15a
12b
14a
14b
14c
15b
16
17
18a
18b

## PLATE XXVI


Figure— PAGE

1–6. *Aulosteges wolfcampensis*, n. sp. ..... 95–96

1. Lateral view of a cotype from loc. 193. Y.P.M. 11849.
2. Mold of exterior of a brachial valve of a cotype from loc. 93. Y.P.M. 11847.
3. Interior of a brachial valve from loc. 93. Y.P.M. 11847.
4. Cardinal process of a cotype from loc. 93. T. 10158.
5a–b. Pedicle and lateral views of a cotype from loc. 88S. Y.P.M. 11839.
6a–b. Lateral and pedicle views of a cotype from loc. 93. T. 10158. Wolfcamp formation, Glass Mountains.

7–11. *Aulosteges medlicottianus* Waagen ..... 93–94

7a–c. Pedicle, brachial, and lateral views of a specimen from loc. D. 3. Leonard formation, Glass Mountains.
8a–c. Lateral, pedicle, and brachial views of a specimen from loc. 108. Y.P.M. 11868. Hess formation, Glass Mountains.
9a–b. Pedicle and brachial views of a specimen from loc. 44. T. 10253. Word formation, Glass Mountains.
10. Brachial view of a specimen from loc. 121. Y.P.M. 11870.
11a–c. Pedicle, brachial, and lateral views of a specimen from loc. 124. Leonard formation, Glass Mountains. T. 11079.


1
2
3
4
5a
5b
6a
6b
7a
7b
7c
8a
9a
10
8b
8c
9b
11a
11b
11c

## PLATE XXVII

Figure— PAGE

1–3. *Aulosteges trigonalis*, n. sp. ..... 94–95

1a–b. Pedicle and lateral views of the holotype, from loc. 231. Y.P.M. 11883. Leonard formation, Glass Mountains.

2a–b. Pedicle and lateral views of a specimen from loc. D. 239.

3. Pedicle view of a specimen from loc. Tx. T. 11014. Word formation, Glass Mountains.

4–7. *Aulosteges tuberculatus*, n. sp. ..... 95

4. Part of a pedicle valve of a cotype from loc. 247. Y.P.M. 11815.

5. Brachial valve of a cotype from loc. T. 239. T. 10608.

6a–b. Lateral and pedicle views of a cotype from loc. 247. Y.P.M. 11815.

7. Pedicle valve of a cotype from loc. T. 239. T. 10608. Word formation, Glass Mountains.

8–9. *Aulosteges medlicottianus* Waagen ..... 93–94

8. Brachial view of a specimen from loc. 124. Y.P.M. 11855. Leonard formation, Glass Mountains.

9. Lateral view of a specimen from loc. 222. Y.P.M. 11874. Hess formation, Glass Mountains.

10–13. *Strophalosia hystricula* Girty ..... 96

10. Posterior view of a pedicle valve from loc. 128. Y.P.M. 11908.

11. Pedicle valve from loc. 128. Y.P.M. 11908.

12. Part of a pedicle valve from loc. 128. Y.P.M. 11908.


13. Part of a pedicle valve from loc. 128. Y.P.M. 11908. Leonard formation, Glass Mountains.

14–16. *Teguliferina bösei*, n. sp. ..... 96–97

14. Large piece from loc. 93, showing many intergrown specimens attached to one another by spines. Y.P.M. 12661.

15a–c. Anterior, apertural, and brachial views of the holotype, from loc. 93. Y.P.M. 12829. In 15c the articulation of the brachial valve may be seen, with part of the pedicle valve growing over it.

16. Two intergrown specimens, showing the position of the brachial valve. Loc. 93. Y.P.M. 12661. Wolfcamp formation, Glass Mountains.


1a
1b
2a
2b
3
4
5
8
9
6a
6b
7
10
11
12
13
14
15a
15b
16
15c


## PLATE XXVIII

Figure— PAGE

1. *Prorichtofenia likharewi* King, n. sp. 98–99

1a. A large colony from the upper side. In some specimens the characteristic feature of the species, the sharp median ridge opposite the inside of the cardinal area, is visible. Y.P.M. 12009.

1b. The same from the lower side, showing their radial arrangement. Leonard formation, Glass Mountains, loc. 128.



1a

1b

## PLATE XXIX

Figure— PAGE

1–5. *Prorichthofenia likharewi*, n. sp. ..... 98–99

1a–b. Lateral and apertural views of a cotype from loc. 120. Y.P.M. 12008.

2a–b. Lateral and apertural views of a cotype from loc. 120. T. 10596.

3a–b. Lateral and apertural views of a cotype fro mloc. 120 Y.P.M. 12008. Leonard formation, Glass Mountains.

4a–b. Posterior and apical views of an internal mold from the limestone of Pichagüia, near Las Delicias, Coahuila. There appears to have been a high median ridge opposite the cardinal area, which proves the specimen to belong to this species. Y.P.M. 12658.

5a–b. Apertural and lateral views of a specimen from loc. 479. Y.P.M. 12010. Hess formation, Sierra Diablo.


6–9. *Prorichthofenia permiana* (Shumard) ..... 99

6a–c. Apertural and two lateral views of a specimen from loc. 144. The brachial valve is broken, but the remaining part is in place, and partly open. Y.P.M. 11969.

7a–b. Apertural and lateral views of three intergrown specimens from loc. 144. Y.P.M. 11969.

8. A small cluster from loc. 144. Y.P.M. 11687.

9. An intergrown mass from loc. 144. T. 10629. Word formation, Glass Mountains.


1a
1b
2a
2b
3a
3b
4a
4b
5a
5b
6a
6b
6c
7a
7b
8
9

## PLATE XXX

Figure— PAGE

1. *Prorichthofenia likharewi*, n. sp. ..... 98– 99

1a–b. Lateral and apertural views of a specimen from loc. 479. Hess formation, Sierra Diablo.

2–7. *Prorichthofenia teguliferoides*, n. sp. ..... 99–100

2a–b. Pedicle and brachial views of a cotype from loc. 479. T. 11064.

3a–b. Pedicle and brachial views of a ctoype from loc. 479. Y.P.M. 11957.

4a–b. Pedicle and brachial views of a specimen from loc. 479. Y.P.M. 11957. Hess formation, Sierra Diablo.

5a–c. Pedicle, brachial, and lateral views of a specimen from loc. 208. The shell is preserved in limestone, and the details shown in Figs. 2–3 are not preserved. Y.P.M. 11952. Hess formation, Glass Mountains.

6a–b. Apertural and lateral views of a specimen from loc. 511. T. 10157.

7. Apertural view of a specimen from loc. 511. Y.P.M. 11958. Hess formation, Finlay Mountains.

8–13. *Prorichthofenia permiana* (Shumard) ..... 99

8a–b. Apertural and lateral views of a specimen from loc. 55. Brachial valve in place. T. 10558.


9. Apertural view of a specimen from loc. Tc. Brachial valve not present. T. 10613.

10a–b. Interior and aperture of a brachial valve from loc. 144, which shows marginal spines, directed internally. Y.P.M. 12659.

11a–b. Lateral and apertural views of a specimen from loc. 247. 11b shows part of a brachial valve in place. Y.P.M. 11920.

12. A slab showing, above, *Lyttonia nobilis americanus* (Girty) n. var., with the brachial valve in place over the inside of the pedicle one; below, part of a *Prorichthofenia permiana*, showing the inside of the cardinal area and the prostrate spines lying inside the ventral cup. From loc. Tx. 148. T. 10677.

13a–b. Colony from loc. 144. Y.P.M. 11969. Word formation, Glass Mountains.


1a
1b
2a
2b
3a
3b
5a
5b
5c
6a
6b
4a
4b
7
8a
8b
9
10a
10b
11a
11b
12
13a
13b

## PLATE XXXI

Figure— PAGE

1–6. *Prorichthofenia uddeni* (Böse) ..... 100

1. Apertural view of a specimen from loc. 128, showing the median ridge on the cardinal side of the cone. T. 10114.

2. Apertural view of a specimen from loc. 128. Y.P.M. 11948.

3a–b. Lateral and apertural views of a specimen from loc. 128. Y.P.M. 11948.

4a–b. Lateral and apertural views of a specimen from loc. 128. T. 10114.

5. Apertural view of a specimen from loc. 120. Y.P.M. 11946.

6a–b. Lateral and apertural views of a specimen from loc. 120, showing the median ridge on the cardinal side of the cone. Y.P.M. 11946. Leonard formation, Glass Mountains.

7–9. *Parakeyserlingina fredericksi*, n. sp. ..... 102

7a–b. Exterior and interior of the holotype, from loc. T. 93. The transition from forward-directed septal loops to loops perpendicular to the median line can be clearly seen. T. 10671.

8. Exterior of a specimen from loc. T. 93. The outer part of the shell has been removed in places, revealing the loops within. T. 10671.

9. Interior of a part of a specimen, from loc. 87 low, showing the divided character of the loops. Y.P.M. 12039. Wolfcamp formation, Glass Mountains.


10–13. *Lyttonia nobilis americanus* (Girty) ..... 103–104

10a–b. Pedicle and lateral views of a specimen from loc. 128. In 10a the shell has been removed, showing the septa within. T. 10572. Leonard formation, Glass Mountains.

11. Mold of the interior of a pedicle valve from loc. 251. T. 10650.

12. Interior of part of a brachial valve from loc. 247. Y.P.M. 12041.

13. Interior of part of a pedicle valve from loc. 247. Y.P.M. 12041. Word formation, Glass Mountains.


1
2
3a
3b
4a
4b
5
6a
6b
7a
7b
10a
10b
8
9
11
12
13

## PLATE XXXII

Figure— PAGE

1–9. *Lyttonia nobilis americanus* (Girty) ..... 103–104; 54

1a–b. Exterior and interior of a pedicle valve from loc. 476. Within the shell may be seen a second, smaller pedicle valve (upper left of Fig. 1b). Y.P.M. 12820. Hess formation, Sierra Diablo.

2. Exterior of part of a brachial valve from loc. 250, showing granular nature of its surface. T. 10557.

3a–b. Interior and exterior of a pedicle valve from loc. T. 240a, showing the tangled posterior outgrowth of the shell and the double character of the first-formed septa, resembling *Parakeyserlingina*. T. 10074.

4. Interior of a pedicle valve from loc. 51, showing the posterior expansion of the shell. T. 11020. Word formation, Glass Mountains.


5. Axial part of the brachial valve of a *Lyttonia*, from loc. 128. Y.P.M. 12042.

6. Exterior of a pedicle valve from loc. 128, through a cavity in which may be seen the interior of the brachial valve. Y.P.M. 12042. Leonard formation, Glass Mountains.

7. Exterior of a pedicle valve, which is largely exfoliated so that the mold of the interior of the shell is visible; from loc. 152. Y.P.M. 12017. Word formation, Glass Mountains.

8. Piece of rock from loc. 151, showing the outside of a *Lyttonia*, traces of the septa showing through the shell. To the left are a *Linoproductus waagenianus* and a *Meekella globosa*. Y.P.M. 12660. Leonard formation, Glass Mountains.

9. Exterior of part of a brachial valve from loc. Tx. 148. T. 10528. Word formation, Glass Mountains.


1a
1b
2
4
3a
3b
7
5
6
8
9

## PLATE XXXIII

Figure— PAGE

1–8. *Leiorhynchus bisulcatum* (Shumard) ..... 105

1a–d. Pedicle, brachial, lateral, and anterior views of a specimen from section 1, bed 19. Y.P.M. 12558.

2a–c. Pedicle, brachial, and lateral views of a specimen from section 2, bed 6. Y.P.M. 12556.

3. Anterior view of a specimen from section 2, bed 6. Y.P.M. 12556.

4a–c. Brachial, lateral, and anterior views of a specimen from section 2, bed 6. Y.P.M. 12556.

5a–c. Pedicle, brachial, and anterior views of a specimen from section 2, bed 6. Y.P.M. 12556.

6. Anterior view of a specimen from section 2, bed 6. T. 11089.

7a–c. Pedicle, brachial, and anterior views of a specimen from section 2, bed 6. Y.P.M. 12556.

8. Pedicle view of a specimen from section 2, bed 6. *Waagenoceras* horizon near Las Delicias, Coahuila.

9–11. *Leiorhynchus weeksi nobilis* (Girty) ..... 105–106

9a–d. Pedicle, brachial, lateral, and anterior views of a specimen from loc. 265. Y.P.M. 12553.

10a–c. Brachial, lateral, and anterior views of a specimen from loc. 144. Y.P.M. 12552.

11. Anterior view of a specimen from loc. 144. T. 10598. Word formation, Glass Mountains.

12–13. *Pugnoides elegans* (Girty) ..... 106

12a–d. Pedicle, brachial, lateral, and anterior views of a specimen from loc. 106. Y.P.M. 12527.

13a–c. Brachial, pedicle, and anterior views of a specimen from loc. 108. Y.P.M. 12697. Hess formation, Glass Mountains.

14–17. *Pugnoides transversus*, n. sp. ..... 108–109

14a–d. Brachial, pedicle, lateral, and posterior views of the holotype, from loc. 222. Y.P.M. 12518.

15a–c. Pedicle, posterior, and lateral views of a specimen from loc. T. 107. T. 10674.


16. Brachial view of a specimen from loc. 212. Y.P.M. 12519.

17a–c. Brachial, lateral, and posterior views of a specimen from loc. 107. Upper Hess formation, Glass Mountains. T. 10670.

18–19. *Lyttonia hortoni*, n. sp. ..... 104

18. Interior of pedicle valve of the holotype, from loc. 7; only the right half is shown. Y.P.M. 12062. Leonard formation, Glass Mountains.

19. Interior of part of a pedicle valve from loc. T. 140, showing the interrupted nature of the septa. T. 10586. Word formation, Glass Mountains.


1a
1b
1c
1d
2a
2b
2c
3
4a
4b
4c
5a
5b
5c
6
7a
7b
7c
8
9a
9b
9c
7d
10a
10b
10c
12a
12b
12c
12d
11
13a
13b
13c
14d
15a
15b
16
14a
14b
14c
17c
15c
17a
17b
19
18

## PLATE XXXIV

Figure— PAGE

1. *Pugnoides bidentatus* (Girty) ..... 107–108

1a–c. Brachial, pedicle, and lateral views of a specimen from loc. 491. Y.P.M. 12562. Bone Canyon limestone, Delaware Mountains.

2–4. *Pugnoides elegans* (Girty) ..... 106

2a–d. Pedicle, brachial, lateral, and anterior views of a specimen from loc. 108. Y.P.M. 12549. Upper Hess formation, Glass Mountains.

3a–b. Pedicle and brachial views of a specimen from loc. 106. T. 10582. Hess formation, Glass Mountains.

4a–b. Brachial and lateral views of a specimen from loc. 104. T. 10645. Leonard formation, Glass Mountains.

5–9. *Pugnoides texanus* (Shumard) ..... 108

5a–d. Brachial, pedicle, lateral, and anterior views of a specimen from loc. 223. Y.P.M. 12530.

6a–d. Brachial, pedicle, lateral, and anterior views of a specimen from loc. 223. Y.P.M. 12530. Upper Hess formation, Glass Mountains.

7a–c. Brachial, pedicle, and anterior views of a specimen from loc. 440a. Y.P.M. 12527. Upper Gym formation, Hueco Mountains.

8a–b. Brachial, pedicle, and anterior views of a specimen from loc. 223. T. 10666.

9. Anterior view of a specimen from loc. 223. T. 10666. Upper Hess formation, Glass Mountains.

10–12. *Pugnoides swallovianus* (Shumard) ..... 107

10a–b. Exterior and interior of a brachial valve from loc. Tx. T. 11037.

11a–c. Brachial, pedicle, and anterior views of a specimen from loc. T. 240. T. 10470.

12a–c. Brachial, lateral, and anterior views of a specimen from loc. T. 240. T. 16470. Word formation, Glass Mountains.

13–14. *Pugnoides shumardianus* (Girty) ..... 107

13a–d. Brachial, pedicle, lateral, and anterior views of a specimen from loc. T. 15a.

14a–c. Brachial, lateral, and anterior views of a small specimen from loc. T. 15a. Leonard formation, Glass Mountains.

15–17. *Pugnoides pinguis* (Girty) ..... 106

15a–d. Pedicle, brachial, lateral, and anterior views of a specimen from loc. 124. Leonard formation, Glass Mountains. T. 10017.

16a–b. Pedicle and lateral views of a specimen from loc. 17. Y.P.M. 12521. Hess formation, Glass Mountains.

17. Anterior view of a specimen from loc. D. 3. Leonard formation, Glass Mountains.

18. *Rhynchopora illinoisensis* (Worthen) ..... 109

18a–d. Pedicle, brachial, lateral, and anterior views of a specimen from loc. T. 88. In 18d the pedicle valve is up, the brachial down. Wolfcamp formation, *Uddenites* zone, Glass Mountains.

19–23. *Rhynchopora taylori* Girty ..... 109–110

19. Anterior view of a specimen from loc. 144. Y.P.M. 12641.

20a–d. Brachial, pedicle, lateral, and anterior views of a specimen from loc. T. 240. Word formation, Glass Mountains. T. 10998a.

21a–b. Brachial and anterior views of a specimen from loc. 436. Y.P.M. 12636. Upper Gym formation, Hueco Mountains.

22a–c. Brachial, lateral, and anterior views of a brachial valve from loc. T. 240. Word formation, Glass Mountains. T. 10998.

23a–b. Brachial and anterior views of a specimen from loc. 436. This and the specimen illustrated in Fig. 21 are not typical because of their small size, but agree with the larger specimens in number of plications. Y.P.M. 12636. Upper Gym formation, Hueco Mountains.

24–27. *Camarophoria deloi*, n. sp. ..... 110

24a–c. Brachial, lateral, and posterior views of the holotype, from loc. 376. In 24c the shell surface is broken, so that the spondylium may be seen.

25a–c. Brachial, lateral, and anterior views of a specimen from loc. 241. Y.P.M. 12693. Word formation, Glass Mountains.

26. Brachial view of a specimen from Sawmill Canyon, Arizona. Y.P.M. 12694.

27a–b. Posterior and brachial views of a specimen from Sawmill Canyon, Arizona. Lower Permian, Santa Rita Mountains.

28. *Camarophoria thevenini* Kozlowski ..... 110

28a–c. Pedicle, brachial, and lateral views of a specimen from loc. 123. Y.P.M. 12695. Leonard formation, Glass Mountains.

29–31. *Camarophoria venusta* Girty ..... 110–111


29a–b. Posterior and brachial views of a specimen from loc. 107. Hess formation, Glass Mountains.

30a–b. Anterior and brachial views of a specimen from loc. T. 15a.

31a–b. Lateral and brachial views of a specimen from loc. D. 1. Leonard formation, Glass Mountains.

32. *Camarophoria? indentata* (Shumard) ..... 111–112

32a–c. Brachial, pedicle, and lateral views of an internal mold from section 1, bed 14. Y.P.M. 12691. Limestone of Cerro Caballo, near Las Delicias, Coahuila.


1a
1b
1c
2a
2b
2c
2d
3a
3b
5a
5b
5c
5d
6a
6b
6c
6d
4a
4b
7a
7b
7c
8a
8b
9
10a
10b
11a
11b
13a
13b
13c
13d
14a
12a
12b
12c
11c
14b
14c
15a
15b
15c
15d
16a
16b
17
18a
18b
19
18c
18d
20a
20b
20c
20d
21a
21b
22a
22b
22c
23a
23b
24a
24b
24c
26
27b
28a
28b
28c
25a
25b
25c
27a
29a
30a
32a
32b
32c
29b
30b
31a
31b

## PLATE XXXV

Figure— PAGE

1–5. *Camarophoria venusta* Girty ..... 110–111

1a–c. Pedicle, lateral, and posterior views of a relatively broad specimen from loc. 484. Y.P.M. 12587. Bone Canyon limestone, Sierra Diablo.

2a–b. Pedicle and lateral views of a specimen from loc. 227. Y.P.M. 12614.

3a–d. Brachial, pedicle, lateral, and anterior views of a specimen from loc. 174. Y.P.M. 12620. Leonard formation, Glass Mountains.

4. Pedicle view of an exceptionally broad specimen from loc. 484. In breadth it resembles *Pugnoides transversus*, but it contains a spondylium. Y.P.M. 12587. Bone Canyon limestone, Sierra Diablo.

5. Interior, viewed from the front. Note cruralium above and spondylium below. The hair-like extension from the right-hand side of the cruralium is one of the crura. Loc. 107. Y.P.M. 12607. Upper Hess formation, Glass Mountains.

6–7. *Uncinuloides guadalupensis* (Shumard) ..... 112–113

6a–c. Brachial, pedicle, and half lateral views of a specimen from loc. T. 128. In 6c may be seen the cruralium projecting spoon-like from the apex of the brachial valve, with one of the hair-like crura just below it on the far side. T. 10167.


7a–c. Pedicle, lateral, and anterior views of a pedicle valve from loc. 120, showing *Uncinulus*-like anterior part. Y.P.M. 12582. Leonard formation, Glass Mountains.

8–10. *Spirifer condor* (d'Orbigny) ..... 113–114

8a–c. Posterior, brachial, and lateral views of a specimen from loc. Tmm. T. 10341.

9a–b. Pedicle and brachial views of a specimen from loc. Tmm, the posterior part of which has been so eroded that the median part of the pedicle valve shows the inner cavity of the shell, filled with limestone, surrounded by the apical callosity. T. 10341.

10. Interior of a pedicle valve from loc. 376. Y.P.M. 12493. Lower Gym formation, Hueco Mountains.


1a
1b
1c
2a
2b
3a
3b
3c
3d
4
5
6a
6b
6c
7a
7b
7c
8a
8b
2286
8c
9a
2286
10
9b

## PLATE XXXVI

Figure— PAGE

1. *Spirifer condor* (d'Orbigny) ..... 113–114


1a–b. Pedicle and brachial views of a specimen from loc. Tmm. The apical part of the pedicle valve has had the shell worn away, so that only the internal mold is visible. T. 10341. Lower Gym formation, Hueco Mountains.

2–4. *Spirifer marcoui infraplica*, n. var. ..... 114–115

2a–c. Posterior, pedicle, and lateral views of an incomplete pedicle valve of a cotype from loc. 499. Y.P.M. 12402. Victorio Peak limestone, Sierra Diablo.

3a–c. Lateral, posterior, and pedicle views of a pedicle valve of a cotype from loc. Tg. T. 10043.

4a–c. Cardinal, brachial, and posterior views of a brachial valve from loc. Tg. T. 10043. Leonard formation, transition beds, Shafter.


1a
1b
2a
2b
2c
3a
2288
3b
2288
4a
3c
4b
4c

## PLATE XXXVII

Figure— PAGE

1–3. *Spirifer (Neospirifer) costella,* n. sp. ..... 115

1. Part of a pedicle valve from loc. 123. Y.P.M. 12439.

2. Pedicle view of a specimen from loc. 126.

3a–d. Posterior, pedicle, brachial, and lateral views of the holotype, from loc. 174. Y.P.M. 12437. Leonard formation, Glass Mountains.

4–6. *Spirifer (Neospirifer) bakeri,* n. sp. ..... 115


4a–b. Pedicle and posterior views of a pedicle valve from loc. 46. Y.P.M. 12500.

5a–b. Posterior and pedicle views of a pedicle valve from loc. 58.

6a–c. Pedicle, posterior, and lateral views of the holotype, from loc. 46. Y.P.M. 12500. Word formation, Glass Mountains.

7. *Spirifer (Neospirifer) mexicanus latus,* n. var. ..... 116

7a–d. Brachial, pedicle, lateral, and posterior views of the holotype, from loc. 123. Y.P.M. 12504. Leonard formation, Glass Mountains.


1
2
3a
3b
3c
3d
4a
4b
5a
5b
6a
6b
6c
7a
7b
7c
7d

## PLATE XXXVIII

Figure— PAGE

1. *Spirifer (Neospirifer) mexicanus latus*, n. var. ..... 116

1a–c. Pedicle, brachial, and posterior views of a specimen from loc. 126. Y.P.M. 12505. Leonard formation, Glass Mountains.

2–3. *Spirifer (Neospirifer) huecoensis*, n. sp. ..... 116–117

2a–c. Pedicle, posterior, and lateral views of the holotype, from loc. 389. In 2b the pedicle valve is up, the brachial down. Y.P.M. 12441. Lower Gym formation, Hueco Mountains.


3a–b. Exterior and interior of a pedicle valve from loc. 107. Y.P.M. 12442. Hess formation, Glass Mountains.

4–6. *Spirifer (Neospirifer) pseudocameratus* Girty ..... 116

4. Lateral view of a specimen from loc. 493. T. 10129. Victorio Peak limestone, Sierra Diablo.

5a–b. Brachial and posterior views of a specimen from loc. 44–46. Y.P.M. 12446. Word formation, Glass Mountains.

6. Brachial view of a specimen from loc. 493. Y.P.M. 12449. Victorio Peak limestone, Sierra Diablo.


1a
1b
1c
2a
2b
2c
3a
3b
4
5a
5b
6

## PLATE XXXIX

Figure— PAGE

1–3. *Spirifer (Neospirifer) pseudocameratus* Girty ..... 116

1. Pedicle view of an internal mold from loc. 6. Y.P.M. 12480. Word formation, Glass Mountains.

2. Lateral view of a specimen from loc. 174. T. 10228.

3a–c. Posterior, pedicle, and brachial views of a specimen from loc. 121. Y.P.M. 12458. Leonard formation, Glass Mountains.

4. *Spirifer (Neospirifer) triplicatus* Hall ..... 117


4a–c. Brachial, lateral, and posterior views of a specimen from loc. 201. Y.P.M. 12424. Wolfcamp formation, *Uddenites* zone, Glass Mountains.

5. *Spirifer (Neospirifer) texanus* Meek ..... 117

5a–b. Lateral and pedicle views of a part of a pedicle valve from loc. 88. Y.P.M. 12399. Wolfcamp formation, *Uddenites* zone, Glass Mountains.

6. *Spirifer (Elivina) sulcifer* Shumard ..... 118

6a–c. Pedicle, brachial, and lateral views of a large specimen from loc. 247. The brachial valve is somewhat crushed inward. Y.P.M. 12414. Upper Word formation, Glass Mountains.


1
2
3a
3b
3c
4a
4b
4c
5a
5b
6a
6b
6c

## PLATE XL

Figure— PAGE

1–5. *Spirifer (Elivina) sulcifer* Shumard ..... 118

1a–c. Lateral, brachial, and posterior views of the posterior part of a specimen from loc. 144. Y.P.M. 12410.

2a–b. Brachial and posterior views of a specimen from loc. 247. Y.P.M. 12414. Word formation, Glass Mountains.

3. Interior of a pedicle valve from loc. 158. Y.P.M. 12418. Upper Word formation, Glass Mountains.

4a–d. Pedicle, brachial, lateral, and posterior views of a specimen from loc. 146. T. 10179.

5a–b. Pedicle and brachial views of a specimen from loc. 247, the posterior part of which is very much eroded. Word formation, Glass Mountains. T. 10174.

6–8. *Squamularia guadalupensis* (Shumard) ..... 118–119

6. Pedicle valve from loc. D. 1. Leonard formation, Glass Mountains.

7a–c. Pedicle, lateral, and posterior views of a specimen from loc. 1. Y.P.M. 12370. Hess formation, Glass Mountains.

8a–b. Brachial and lateral views of an internal mold from loc. 246. T. 10599. Word formation, Glass Mountains.

9–13. *Martinia wolfcampensis*, n. sp. ..... 121


9a–d. Brachial, pedicle, lateral, and anterior views of the holotype, from loc. 88S. Y.P.M. 12295.

10. Interior of pedicle valve from loc. 93. Y.P.M. 12296.

11a–c. Pedicle, brachial, and lateral views of a specimen from loc. 94. The shell is dorsally crushed, so that the true thickness can not be seen. Y.P.M. 12287.

12. Posterior view of a specimen from loc. 88. T. 10010.

13. Pedicle view of a specimen from loc. 88. The apex of the pedicle valve is crushed. T. 10010. Wolfcamp formation, Glass Mountains.


1a
1b
1c
2a
2b
3
4a
4b
4c
4d
5a
5b
6
7a
7b
7c
8a
8b
9a
9b
9c
9d
10
11a
11b
11c
12
13

## PLATE XLI

Figure— PAGE

1–5. *Ambocoelia guadalupensis* Girty ..... 119–120

1a–d. Pedicle, brachial, posterior, and lateral views of a specimen from loc. T. 120. T. 10061. Leonard formation, Glass Mountains.

2. Brachial view of an exceptionally small specimen rather doubtfully referred to this species. Loc. 414. Y.P.M. 12347. Middle Gym formation, Hueco Mountains.

3a–d. Pedicle, brachial, lateral, and posterior views of a specimen from loc. 491. Y.P.M. 12348. Bone Canyon limestone, Delaware Mountains.

4a–b. Pedicle and posterior views of a specimen from loc. 265. Y.P.M. 12340.

5a–b. Pedicle and lateral views of a specimen from loc. Tc. T. 10065. Word formation, Glass Mountains.

6–10. *Martinia rhomboidalis* Girty ..... 120


6a–b. Brachial and posterior views of a specimen from loc. 77. The locality is supposed to be in the Wolfcamp, but probably this and other fossils were derived from float originating above. T. 10171.

7a–b. Posterior and brachial views of a specimen from loc. 120. T. 10153. Leonard formation, Glass Mountains.

8a–c. Pedicle, lateral, and posterior views of a pedicle valve from loc. 57. Y.P.M. 12319. Capitan formation, Altuda member, Glass Mountains.

9a–c. Posterior, pedicle, and lateral views of a specimen from loc. 123. Y.P.M. 12325. Leonard formation, Glass Mountains.

10. Large piece from loc. 57 containing several specimens. Y.P.M. 12319. Capitan formation, Altuda member, Glass Mountains.


1a
1b
1c
3a
3b
3c
1d
2
4a
4b
5a
5b
3d
6a
7a
7b
6b
8a
8b
8c
9a
10
9b
9c

## PLATE XLII

Figure— PAGE

1–6. *Spiriferina hilli* Girty ..... 122
1. Brachial view of a specimen from loc. D. 9. Leonard formation, Glass Mountains.
2. Pedicle view of a broad, non-typical specimen from loc. 208. Y.P.M. 12270. Hess formation, Glass Mountains.
3. Pedicle view of a specimen from loc. D. 20.
4. Posterior view of a specimen from loc. D. 20. Leonard formation, Glass Mountains.
5. Lateral view of a specimen from Cerro Pichagüia. Y.P.M. 12271.
6a–b. Posterior and pedicle views of a specimen from Cerro Pichagüia. Y.P.M. 12271. Limestone of Pichagüia, near Las Delicias, Coahuila.

7–11. *Spiriferina laxa* Girty ..... 122–123
7. Interior of the apical part of a brachial valve from Cerro Pichagüia, showing the cardinal process, and hinge-sockets. Y.P.M. 12318.
8a–c. Exterior, interior, and lateral views of a pedicle valve from Cerro Pichagüia, showing median septum and hinge-teeth. Limestone of Pichagüia, near Las Delicias, Coahuila. Y.P.M. 12318.
9a–c. Pedicle, lateral, and posterior views of a pedicle valve from loc. T. 148. T. 9988.
10. Posterior view of a specimen from loc. 192. T. 9987.
11. Pedicle view of an exceptionally broad specimen from loc. 46. Y.P.M. 12311. Word formation, Glass Mountains.

12–13. *Spiriferina angulata*, n. sp. ..... 122
12a–c. Pedicle, anterior, and lateral views of the holotype, from loc. 478. Y.P.M. 12301. Hess formation, Sierra Diablo.
13. Brachial view of a fragmentary specimen from loc. 106. T. 10675. Hess formation, Glass Mountains.

14. *Spiriferina pyramidalis* Girty ..... 123–124
14a–c. Posterior, brachial, and lateral views of a specimen from loc. 264. Y.P.M. 12308. Word formation, Glass Mountains.

15–18. *Punctospirifer billingsii* (Shumard) ..... 124–125
15a–c. Exterior, interior, and lateral views of a pedicle valve from loc. Tx. T. 10991.
16. Part of a brachial valve from loc. T. 148, showing nature of surface. T. 9958.
17a–b. Pedicle and lateral views of a pedicle valve from loc. 241. Y.P.M. 12285.
18a–c. Brachial, lateral, and posterior views of a specimen from loc. Tc. T. 9967. Word formation, Glass Mountains.

19. *Hustedia bipartita* Girty ..... 125
19a–c. Pedicle, lateral, and posterior views of a specimen from loc. D. 2. Leonard formation, Glass Mountains.

20–26. *Hustedia pusilla* (Girty) ..... 127–128
20a–c. Pedicle, brachial, and lateral views of a specimen from an unknown locality and horizon. T. 11032.
21a–b. Pedicle and anterior views of a specimen from an unknown locality and horizon. T. 11030.
22. Pedicle view of a specimen from an unknown locality and horizon. Y.P.M. 12686.
23. Anterior view of a specimen from an unknown locality and horizon.
24. Pedicle view of a specimen from loc. 195a, doubtfully conspecific with *H. pusilla*. Y.P.M. 12690. Wolfcamp formation, Glass Mountains.
25. Pedicle view of a specimen from an unknown locality and horizon.
26a–b. Front and posterior views of the jugum of an etched specimen from loc. 46. Y.P.M. 12688. Word formation, Glass Mountains.


27–32. *Hustedia mormoni papillata* (Shumard) ..... 127
27. Lateral view of a specimen from an unknown locality and horizon.
28. Brachial view of a specimen from an unknown locality and horizon. T. 11024.
29. Exterior of an etched pedicle valve from an unknown locality. T. 11018. Word formation.
30a–c. Brachial, lateral, and posterior views of a specimen from an unknown locality and horizon. Y.P.M. 12683.
31a–d. Pedicle, brachial, lateral, and posterior views of a specimen from loc. 243. Y.P.M. 12187.
32a–b. Brachial and lateral views of a specimen from loc. 239. Word formation, Glass Mountains.

33. *Hustedia mormoni* (Marcou) ..... 127
33a–c. Brachial, lateral, and posterior views of a specimen from loc. 93. Y.P.M. 12174. Wolfcamp formation, Glass Mountains.

34–39. *Hustedia meekana* (Shumard) ..... 126–127
34a–d. Pedicle, brachial, lateral, and posterior views of a specimen from loc. 123. T. 10023. Leonard formation, Glass Mountains.
35. Brachial view of a specimen from loc. 491. Y.P.M. 12075. Bone Canyon limestone, Delaware Mountains.
36. Brachial view of a specimen from loc. T. 107. T. 9971. Hess formation, Glass Mountains.
37a–d. Pedicle, brachial, lateral, and posterior views of a specimen from loc. 123. T. 10023. Leonard formation, Glass Mountains.
38a–c. Brachial, pedicle, and lateral views of a specimen from loc. T. 243. T. 9973. Word formation, Glass Mountains.
39. Brachial view of a specimen from loc. 25. Y.P.M. 12094. Leonard formation, Glass Mountains.

40–42. *Hustedia huecoensis*, n. sp. ..... 125–126
40a–c. Brachial, pedicle, and lateral views of a cotype from loc. 440a. Y.P.M. 12123.
41a–d. Brachial, pedicle, lateral, and posterior views of a cotype from loc. 440a. T. 11004.
42. Pedicle valve from loc. 440a. Y.P.M. 12123. Upper Gym formation, Hueco Mountains.

43–46. *Hustedia hessensis*, n. sp. ..... 125
43a–c. Pedicle, lateral, and posterior views of a specimen from loc. 106. T. 10064. Hess formation, Glass Mountains.
44a–c. Brachial, lateral, and posterior views of a cotype from loc. 478. Y.P.M. 12127.
45. Brachial view of a cotype from loc. 478. Y.P.M. 12127.
46. Brachial view of a specimen from loc. 474. T. 10146. Hess formation, Sierra Diablo.


1
2
3
4
5
6a
6b
7
8a
9a
9b
9c
10
11
8b
12a
12b
12c
13
15a
15b
15c
14a
14b
14c
8c
16
17a
17b
18a
18b
18c
19a
19b
19c
20a
20b
20c
21a
21b
22
23
24
25
26a
26b
27
28
29
30a
30b
30c
31a
31b
31c
31d
32a
32b
33a
33b
33c
34a
34b
34c
34d
35
36
37a
37b
37c
37d
38a
38b
38c
39
40a
40b
40c
41a
41b
41c
41d
42
43a
43b
43c
44a
44b
44c
45
46

## PLATE XLIII

Figure— PAGE

1–11. *Composita mexicana* (Hall) ..... 128–129

1a–e. Pedicle, brachial, lateral, anterior, and posterior views of a specimen from loc. 223. T. 9993.

2a–e. Pedicle, brachial, lateral, anterior, and posterior views of a specimen from loc. 223. T. 9993.

3a–e. Pedicle, brachial, lateral, anterior, and posterior views of a specimen from loc. 223. T. 9993. Hess formation, Glass Mountains.

4a–d. Brachial, posterior, pedicle, and lateral views of a specimen from loc. 397. Y.P.M. 12220. Lower Gym formation, Hueco Mountains.

5a–e. Pedicle, brachial, lateral, anterior, and posterior views of a specimen from loc. 223. T. 9993. Hess formation, Glass Mountains.

6. Pedicle view of a specimen from loc. 436b. Y.P.M. 12217. Upper Gym formation, Hueco Mountains.

7a–b. Brachial and lateral views of a specimen from an unknown locality and horizon. Y.P.M. 12681.

8a–c. Pedicle, brachial, and posterior views of a specimen from loc. 223. T. 9993.

9a–b. Brachial and lateral views of a specimen from loc. 223. T. 9993. Hess formation, Glass Mountains.

10a–d. Pedicle, brachial, lateral, and anterior views of a specimen from loc. 436b. Y.P.M. 12217. Upper Gym formation, Hueco Mountains.

11a–b. Pedicle and lateral views of a specimen from loc. 451. Gym formation, Diablo Plateau.

12–17. *Composita emarginata affinis* Girty ..... 128

12a–d. Brachial, lateral, posterior, and anterior views of a specimen from loc. 247. T. 10118.

13. Brachial view of an internal mold from loc. 247. Y.P.M. 12249.

14a–b. Posterior and brachial views of an internal mold from loc. 247. T. 10118.

15a–d. Pedicle, brachial, lateral, and anterior views of a specimen from loc. 247. T. 10118. Middle and upper Word formation, Glass Mountains.

16a–b. Brachial and lateral views of a specimen from bed 12, section 12. Y.P.M. 12251. Cerro Caballo limestone, Las Delicias, Coahuila.

17. Pedicle view of a specimen from loc. 145. Y.P.M. 12258. Middle and upper Word formation, Glass Mountains.


18–19. *Composita persinuata* (Meek) ..... 130

18a–c. Pedicle, posterior, and anterior views of a specimen from loc. 55. T. 10591.

19a–e. Pedicle, brachial, lateral, anterior, and posterior views of a specimen from loc. 241. Y.P.M. 12277. Word formation, Glass Mountains.

20. *Composita subtilita* (Hall) ..... 130

20a–b. Lateral and brachial views of a specimen from loc. 93. Y.P.M. 12699. Wolfcamp formation, Glass Mountains,


1a
1b
1c
1d
1e
2a
2b
2c
2d
2e
3a
3b
3c
3d
3e
4a
4b
4c
5a
5b
5c
5d
5e
6
4d
7a
7b
8a
8b
8c
9a
9b
10a
10c
10d
10b
11a
11b
12a
12c
12d
13
14a
14b
15a
15b
15c
15d
16a
16b
17
18a
18b
20a
19a
19b
19c
19d
19e
20b

## PLATE XLIV

Figure— PAGE

1–2. *Composita subtilita* (Hall) ..... 130

1a–b. Pedicle and anterior views of a specimen from loc. T.107. T. 10137. Hess formation, Glass Mountains.

2a–b. Lateral and pedicle views of a specimen from loc. D. 3. Word formation?, Glass Mountains.

3–8. *Composita mira* (Girty) ..... 129

3a–c. Brachial, lateral, and posterior views of a specimen from loc. Te. T. 10207. Gym formation, Three-mile Mountain, near Van Horn.

4a–d. Brachial, pedicle, lateral, and posterior views of a specimen from loc. 174. Y.P.M. 12203. Leonard formation, Glass Mountains.

5a–c. Brachial, lateral, and pedicle views of a specimen from loc. 484. Y.P.M. 12205. Bone Canyon member, Leonard formation, Sierra Diablo.

6a–c. Brachial, pedicle, and lateral views of a specimen from loc. 128. T. 10210.

7. Pedicle view of a specimen from loc. 174. Y.P.M. 12203. Leonard formation, Glass Mountains.

8a–b. Pedicle and lateral views of a specimen from loc. 46. Y.P.M. 12196. Word formation, Glass Mountains.

9–10. *Composita subtilita angusta,* n. var. ..... 130–131

9a–d. Brachial, pedicle, lateral, and anterior views of a cotype from loc. 456. Y.P.M. 12191. Upper Wolfcamp formation, Sierra Diablo.

10a–d. Pedicle, brachial, lateral, and posterior views of a cotype from loc. 199. T. 10143. *Uddenites* member, Wolfcamp formation, Glass Mountains.

11. *Dielasma bovidens* (Morton) ..... 132

11a–c. Brachial, lateral, and posterior views of a specimen from loc. 199. Y.P.M. 12140. *Uddenites* member, Wolfcamp formation, Glass Mountains.

12–14. *Dielasma schucherti minor,* n. var. ..... 133

12a–d. Brachial, pedicle, lateral, and anterior views of a specimen from loc. 239. Y.P.M. 12156.

13a–b. Brachial and anterior views of a specimen from loc. 239. T. 9982.

14a–b. Brachial and anterior views of a specimen from loc. 239. Y.P.M. 12156. Word formation, Glass Mountains.

15. *Dielasmina guadalupensis* Girty ..... 133

15a–d. Lateral, brachial, and anterior views of a specimen from loc. 223. Y.P.M. 12159. Hess formation, Glass Mountains.

16. *Dielasma problematicum wordense,* n. var. ..... 131–132

16a–e. Brachial, pedicle, lateral, posterior, and anterior views of the holotype from loc. T.144. T. 10312. Word formation, Glass Mountains.

17–20. *Dielasma spatulatum* Girty ..... 132

17. Etched interior of a brachial valve from loc. Tx. T. 11062.


18a–c. Brachial, pedicle, and lateral views of a specimen from loc. T. 240. T. 10151.

19a–c. Pedicle, lateral, and anterior views of a specimen from loc. 144. Y.P.M. 12144.

20a–d. Pedicle, brachial, posterior, and anterior views of a specimen from loc. T. 240. T. 10151. Word formation, Glass Mountains.

21. *Dielasma shafterense,* n. sp. ..... 132

21a–d. Pedicle, brachial, lateral, and posterior views of the holotype from loc. Tp. T. 10025. Cibolo limestone, Shafter district.


1a
1b
2a
2b
3a
3b
3c
4a
4b
4c
4d
5a
5b
6a
6b
6c
7
8a
8b
5c
9a
9b
9c
9d
10a
10b
10c
10d
12a
12b
12c
12d
13a
13b
14a
14b
11a
11b
11c
16a
16b
16c
16d
16e
15a
15b
15c
17
18a
18b
18c
19a
19b
19c
19d
20a
20b
20c
20d
21a
21b
21c

# INDEX


PAGE

Abo sandstone ..... 25, 26, 28, 29, 30
Acambona ..... 41
Admiral formation ..... 19
Adrianites marathonensis ..... 17
Agathiceras girtyi ..... 17
  texanum ..... 11, 33
Akasaka limestone ..... 39
Alaska ..... 31, 35
Alethopteris ..... 29
Allorisma capax ..... 22
Alta formation ..... 17
Altuda ..... 10, 13
Ambocoelia ..... 119
  guadalupensis ..... 31, 32, 119
  planoconvexa ..... 41, 120
Amdrup's Land ..... 34
Americus limestone ..... 22, 23
Andean "Upper Carboniferous" ..... 7, 35, 36, 37, 38, 41
Annularia ..... 20
Anomia craniolaris ..... 42
Anomites glabra ..... 120
  scabriculus ..... 79
  semireticulatus ..... 66
Apache formation ..... 13
Apache Mountains ..... 13
Araucarites ..... 23
Arbuckle Mountains ..... 21, 23
Archer County ..... 19
Ardmore Basin ..... 23
Arizona ..... 5, 9, 13, 24, 25, 28, 29, 32, 36
Arizona Plateau ..... 29, 30
Arroyo formation ..... 20
Artinsk stage ..... 31, 32, 33, 35
Aspicosaurus ..... 26
Athyridae ..... 41, 42, 128
Atremata ..... 37
Atripidae ..... 41
Aulosteges Beedei ..... 92
  guadalupensis ..... 11, 15, 39, 93
  helmerson ..... 92
  hispidus ..... 31, 32
  magnicostatus ..... 11, 12, 15, 93
  medlicottianus ..... 39, 93
  subcostatus King ..... 94
  triagonalis ..... 94
  tuberculatus ..... 95
  wolfcampensis ..... 6
Aurin, F. L. ..... 21
Aviculopecten coloradoensis ..... 25
Avonia ..... 39
  boulei ..... 6, 7, 16, 16, 25, 36, 82
  incurvata ..... 83
  latidorsata ..... 83
  meekana ..... 83
  signata ..... 83
  subhorrida ..... 31, 32, 84
  subhorrida rugatula ..... 25, 84
  walcottiana costata ..... 85
  walcottiana ..... 85
Baird shale ..... 33
Baker, A. A. ..... 30
Baker, C. L. ..... 17, 26, 28
Bakewellia parva ..... 33
Banff, Alberta ..... 33
Banks Island ..... 33, 34
Baylor County ..... 19
Baylor Mountains ..... 14
Bear Island ..... 35
Beede, J. W. ..... 5, 17, 21, 22, 35
Belle Plains formation ..... 19
Bellerophon ..... 29, 30, 31, 32
  carbonarius ..... 25
  crassus ..... 25
  majusculus ..... 30
  percarinatus ..... 25
Big Blue group ..... 23
Big Wichita River ..... 19
Bisbee district ..... 24
Bissett formation ..... 10
Blaine formation ..... 20, 21, 22, 23, 28
Blanchard, W. G. ..... 12, 13, 27, 28
Bolivia ..... 35
Bone Canyon ..... 11, 12, 17, 28
Bone Canyon black limestone ..... 14
Bone Springs ..... 28
Böse, Emil ..... 5, 6, 8, 19, 20, 26
Brackett Draw ..... 15
Brachythyris ..... 41
Brazos River ..... 18, 19
Bryan, Kirk ..... 30
Buck Creek formation ..... 23
Bullwagon dolomite ..... 20
Buxtonia longa ..... 39
  occidentalis ..... 39, 79
  maxima ..... 19

PAGE

Buxtonia ..... 79
  peruviana ..... 16, 25, 26, 36, 39, 80
  victorioensis ..... 15, 80
Callipteris ..... 19, 23, 26
  conferta ..... 29
California ..... 31, 33
Camerophorella ..... 41
Camarophoria ..... 110
  deloi ..... 16, 25, 40, 110
  cf. purdoni ..... 13, 14
  indentata ..... 111
  thevenenì ..... 36, 40, 110
  venusta ..... 33, 40, 110
Canadian Cordillera ..... 33
Cape Sheridan ..... 34
Capitan limestone ..... 8, 10, 12, 13, 27, 28, 29, 38, 40
Captorhinus ..... 20
Carrizo Mountains ..... 17
Carlsbad ..... 13, 27
Castile gypsum ..... 12, 13
Cathedral Mountain ..... 10
Cedar Hills ..... 23
Cerro Caballo ..... 18
Cerro Pichagüia ..... 18
Chaetetes milleporaceus ..... 24
Chase formation ..... 23
Chickasha formation ..... 21, 23
Chinati Mountains ..... 17
Chinati Series ..... 17
Chiricahua Mountains ..... 25
Chonetella nasuta ..... 34
Chonetes ..... 38, 60
  biplicatus ..... 60
  deliciaensis ..... 60
  geinitzianus ..... 36
  glaber ..... 36
  granulifer ..... 60
  hessensis ..... 61
  hillanus ..... 12, 61
  molengraafi ..... 17, 61
  ostiolatus ..... 31, 32
  permianus ..... 62
  quadratus ..... 29, 63
  spinoliratus diabloensis ..... 63
  spinoliratus ..... 63
  subliratus ..... 17, 63
  verneuilianus ..... 64
Chonetidae ..... 60
Choza formation ..... 20
Chupadera ..... 13, 15, 16, 17, 26, 27, 28
Cibolo Basin ..... 17
Cibolo Creek ..... 17
Cibolo formation ..... 17
Cieneguita formation ..... 17
Cimarron ..... 23
Cisco group ..... 6, 18, 20, 23, 40
Clay slide ..... 9
Clear Fork ..... 19, 20, 21, 22, 23, 28
Clepsydrops ..... 26
Cliothyridina ..... 41
Clisophyllum gabbi ..... 33
Cloud Chief gypsum ..... 21
Cloudcroft, New Mexico ..... 28
Clyde formation ..... 19
Coahuila, State of ..... 18
Coconino ..... 25, 29, 30, 31
Colorado, State of ..... 30
Colorado River ..... 20
Coleman Junction limestone ..... 18, 19, 21
Comphostrobus ..... 19
Composita ..... 24, 28, 41
  emarginata affinis ..... 10, 12, 15, 41, 128
  mexicana ..... 5, 9, 11, 16, 24, 25, 26, 27, 28, 128
  mira ..... 31, 32, 129
  peperinuata ..... 31, 32, 130
  subtilita ..... 24, 26, 27, 41, 130
  subtilita angusta ..... 130
  xetra ..... 41
Concho River ..... 20
Cooks Range ..... 24
Cora ..... 35
Cottonwood limestone ..... 18, 21, 22
Council Grove formation ..... 23
Crandall, K. H. ..... 13
Crania? sp. ..... 42
  permiana ..... 97
  retzius ..... 42
Craniidae ..... 42
Cricotus ..... 20
Cross, W. ..... 30
Crow Flat ..... 15
Crum, H. E. ..... 30
Culm sandstone ..... 34
Cutler formation ..... 30

PAGE

Cyathophyllum ... 35
Cyrtia ... 41
Cyrtina ... 41
Dalberg, Texas ... 17
Darton, N. H. ... 11, 13, 25, 26, 27, 28, 29
Davis, M. J. ... 12, 13, 27, 28
Day Creek dolomite ... 21
Deer Mountain ... 16
Defiance uplift ... 29
Delaware Mountains ... 5, 11,12, 13, 14, 17, 27
Delaware Mountain formation ... 10, 12, 13, 14, 15, 27, 28
Delo, D. M. ... 5
Delthyris ... 41
Deltopecten occidentalis latiformis ... 33
Deming, New Mexico ... 24
Derby, O. A. ... 26
Derbya buchi ... 16, 36, 38, 58
  crassa ... 38
  crenulata ... 38, 58
  cymbula ... 38, 59
  elevata ... 12
  nasuta ... 38
  waagen ... 58
Diablo Mountains ... 26
Diablo Plateau ... 15, 16, 17
Diadectes ... 26
Diasparactus ... 26
Dicranophyllum ... 19
Dielasma ... 41, 133
  bovidens ... 41, 132
  problematicum wordenses ... 131
  schucherti ... 21, 22
  schucherti minor ... 22, 133
  shafterense ... 132
  wordensis ... 131
Dielasmina ... 42
  guadalupensis ... 133
Discinidae ... 42
Djulfa ... 38
Dog Creek shale ... 21
D'Orbigny ... 35, 37
Dos Alamos ... 15
Dobbin, C. E. ... 30
Double Mountain ... 20, 21, 22, 23, 28
Douglas, I. A. ... 35
Douglas, Arizona ... 25
Dugout Mountain ... 7, 8, 19
Dunbar, C. O. ... 5, 18, 23
Duncan formation ... 21
Dundee, Archer Co., Texas ... 19
Echinocrinus ... 24, 25
Elcabrosaurus ... 26
Ellesmere Island ... 32, 34, 35
Elmdale ... 22, 23
El Paso limestone ... 24
Elvina ... 117
Emory's United States and Mexican Boundary Survey ... 5
Enteletes ... 11, 37, 45
  audii ... 36, 37
  dumblei ... 8, 16, 36, 45
  leonardensis ... 11, 46
  liumbonus ... 11, 46
  plummeri ... 8, 37, 47
  wolfcampensis ... 47
  wordensis ... 11, 47
Eolyttonia ... 100
Eskridge shale ... 22
Euconispira taggarti ... 25
Euphemus carbonarius arenarius ... 33
Eumetria ... 41
Ferghana ... 39
Fielden Peninsula ... 34
Finlay Done ... 17
Finlay Mountains ... 17
Flower Pot ... 23
Florida Mountains ... 16, 17, 24
Franklin Mountains ... 16, 17
Frijole limestone ... 12, 13
Fulda, Texas ... 19
Fusulina ... 14, 15, 22, 23
  californica ... 33
  cylindrica ... 24
  elongata ... 10, 33
  emaciata ... 22
  gracilis ... 33
  longissimoidea ... 22
Fusulinella ... 24
Galiuro Mountains ... 25
Gaptank formation ... 6, 7, 8, 38
Garber formation ... 21
Gastrioceras ... 12, 31
  n. sp. ... 20
  roadense ... 10, 32
  serratum ... 12
  simulator ... 32
Gemmellaroia ... 40
Geyerella ... 38, 57
  americana ... 8, 57
Gigantoperis ... 19, 20, 21, 29
Gilliam sandstone ... 10

PAGE

Gilliland Canyon ... 10
Gilluly, James ... 30
Girty, G. H. ... 5, 6, 11, 12, 13, 14, 24, 25, 26, 27, 28, 29, 30, 34, 37, 38, 41
Glass Mountains ... 6, 7, 8, 9, 10, 11, 13, 14, 17, 18, 19, 20, 21, 22, 25, 26, 27
Glenopteris ... 23
Glen Rose limestone ... 17
Glorieta ... 25, 26, 30
Gompnostrobus ... 26
Goodridge formation ... 30
Gould, C. N. ... 21, 28
Grand Canyon ... 25, 28, 29, 30
Guadalupe ... 11, 27, 28
Guadalupe Mountains ... 5, 10, 11, 12, 13, 15, 25, 27, 28, 37
Guadalupe Point ... 11, 12, 13
Guadalupian fauna ... 6
Gym ... 8, 9, 15, 16, 17, 24, 25, 26, 27, 28
Haack, W. ... 18
Hale, Allen D. ... 30
Hall, J. ... 5
Harper red sandstone ... 23
Harrisburg gypsiferous member ... 29
Hart limestone ... 23
Heiberg Island ... 32, 34, 35
Hemiptichina ... 41
Hennessey formation ... 21
Hermit age ... 30
Hermit shale ... 29
Hermosa limestone ... 30
Hess Canyon ... 10
Hess formation 7, 8, 9, 10, 11, 14, 15, 16, 17, 20, 25, 26, 28, 38, 40
Hess limestone ... 14
Heterelasma ... 41
Hill, R. T. ... 5
Holbrook Dome ... 29
Holm's Land ... 37
Horridonia ... 35, 39, 85
  texana ... 85
  timanica ... 33, 34
Hueco formation ... 14
Hueco Mountains ... 5, 8, 15, 16, 24, 25, 26
Hustedia compressa ... 33
  huecoensis ... 16
  indica ... 41
  meekana ... 25
  mormoni ... 25
  pusilla ... 127
Idaho ... 30, 31
Inyo Mountains ... 31
Iron Mountain ... 9
Isogramma ... 38, 64
  millepunctata ... 65
Kaibab ... 9, 25, 27, 29, 30, 31, 32, 37
Kaibab Plateau ... 29
Kansas ... 18, 21, 22, 23, 30
Karnic Alps ... 7, 8
Kensuella ... 39
Kent County ... 20
Keyserlingina ... 40
  darvasica ... 101
Kiangsiella ... 38
Kindle, E. M. ... 33
King, Philip B. ... 5, 13
King Survey ... 31
Kirk, Edwin ... 13, 31
Kniker, Hedwig ... 22, 35
Kuiu Island ... 33
Kozlowski, Roman ... 35, 36, 37
Labidosaurus ... 20
Lake Titicaca ... 35, 36
Lakewood, New Mexico ... 22
Las Delicias, hacienda of ... 18
Las Puerticitas ... 18
Last Chance Canyon ... 28
Lee, W. T. ... 26
Leiorhynchus ... 40, 104
  bisulcatum ... 105
  rockymontanus ... 40
  weeksi ... 32, 33
  weeksi nobilis ... 105
Leonard ... 5, 8, 9, 10, 11, 12, 14, 15, 17, 18, 20, 21, 22, 25, 26, 27, 28, 29, 30, 40
Leonard-Double Mountain types ... 20
Leonard Mountain ... 9, 10
Lewis, F. E. ... 21
Limnocolis ... 26
Linoproductus ... 38, 75
  aagardi ... 38
  cancrini ... 38
  cancriniformis ... 36, 38
  cora ... 16, 24
  cora angustus ... 76
  girtyi ... 76
  nasutus ... 76
  phosphaticus ... 31, 32, 38, 77
  villiersi ... 7, 16, 36, 38, 78
  waagenianus ... 15, 31, 32, 38, 77
Little Dragoon Mountains ... 25

PAGE
Lloyd, E. Russell ..... 12, 13, 27
Longwell, C. R. ..... 30
Lo Ping ..... 38
Lueders formation ..... 20
Luna County ..... 24, 26
Lyttonia ..... 7, 8, 9, 10, 12, 14, 33, 40
hortoni ..... 104
nobilis americanus ..... 103
waagen ..... 102
Lyttoniidae ..... 100
Magdalena group ..... 26
Malone Mountains ..... 17
Mansfield, G. R. ..... 31
Manzano ..... 25, 27, 28, 29
Marathon, Texas ..... 6, 23, 26
Marcou, J. ..... 5
Marginifera ..... 25, 27, 39
capaci ..... 7, 16, 36, 39, 86
cristobalensis ..... 9, 11, 15, 27, 39, 86
duboutensis ..... 39, 87
lasallensis ..... 39, 87
manzanica ..... 9, 15, 27, 39, 87
opimus ..... 12, 15, 87
popei ..... 12, 88
reticulata angusta ..... 89
reticulata ..... 39, 89
sublaevis ..... 15, 17, 89
texana ..... 89
waagen ..... 86
wabashensis ..... 90
whitei ..... 39, 90
wordensis ..... 39, 90
Marion formation ..... 23
Martinia ..... 120
rhomboidalis ..... 41, 120
subquadrata ..... 10
triqueta ..... 41
wolfcampensis ..... 6, 121
Martiniopsis ..... 41
Maybelle, Baylor County, Texas ..... 19
McKittrick Canyon ..... 12
McKnight, E. T. ..... 30
Medlicottia ..... 17, 20
copei ..... 20
burckhardti ..... 20
Meek, F. B. ..... 31, 37
Meekella ..... 52
Meekella attenuata ..... 11, 52
difficilis ..... 53
globosa ..... 54
grandis ..... 54
hessensis ..... 56
irregularis texana ..... 6, 7, 38, 55
mexicana ..... 16, 22, 24, 26, 55
mutilirata ..... 53
pyramidalis ..... 25, 29
skenoides ..... 15, 56
striatocostata ..... 56
Melville Island ..... 33
Meristellidae ..... 41
Mexico ..... 18
Meyer ..... 35
Michelina ..... 24
Military Crossing of the Big Wichita ..... 19
Miller, A. K. ..... 6, 8
Miser, H. D. ..... 30
Mississippian ..... 28, 38
Moab region ..... 30
Mogollon Plateau ..... 29
Montana ..... 30, 31
Montoya limestone ..... 24
Moore, R. C. ..... 18, 20, 23, 30, 37
Morgan, George D. ..... 23
Mowry Mine ..... 25
Murchisonia terebra ..... 25
Myalina copei ..... 23
wyomingensis ..... 33
Nacimento range ..... 26
Naco limestone ..... 24, 26
Nation River ..... 33
Neospirifer ..... 115
Neotremata ..... 37, 42
Neva limestone ..... 22, 23
Nevada ..... 31
Newberry, I. S. ..... 37
New Mexico ..... 9, 13, 15, 16, 21, 24, 25, 26, 27, 29, 30, 36
Nosoni sandstone ..... 33
Notothyris ..... 41
Novaya Zemlya ..... 35
Odontopteris ..... 22, 23
fischeri ..... 19
neuropteroides ..... 19
Officer, H. G. ..... 21
Ojo Bonito ..... 17
Oklahoma ..... 21, 22, 23
Oldhamina ..... 40
Omphalotrochus ..... 22, 25, 31
obtuspira ..... 25
whitneyi ..... 25, 33
Onyx Mine ..... 25
PAGE
Ophiacodon ..... 26
Orbicula morrisi Davidson ..... 42
Orbiculoidea arenaria ..... 33
convexa ..... 29
d'Orbigny ..... 42
missouriensis ..... 32, 42
Ordovician ..... 14
Orthacea ..... 43
Orthis buchi ..... 58
Orthis? morgiana ..... 44
Orthis umbonata ..... 119
Orthis striatella ..... 60
Orthotetella ..... 38
wolfcampensis ..... 51
Orthothetina cf. Orthothetes mutabilis ..... 52
Orthothetina ..... 52
Orthothetinae ..... 38
Orthotichia ..... 37, 44
Orthotichia kozlokskii ..... 7, 36, 45
Orthotichia morgani kozlowski ..... 36
Overtonia cristato-tuberculata ..... 82
Overtonia ..... 39, 82
Owenyo limestone ..... 31
Paige, Sidney ..... 30
Paleozoic ..... 26
Paraceltites elegans ..... 11, 12
Paradise, Arizona ..... 25
Parakeyserlingina ..... 40, 100, 102
Paraenteletes cooperi ..... 7, 49
Parenteletes ..... 37, 48
Paraphorhynchus obscurum ..... 33
Parry Islands ..... 33, 35
Pastrioceras? serratum ..... 12
Peabody Museum ..... 5
Pecopteris ..... 26, 29
Pecos City, Texas ..... 26
Pecos River ..... 27
Pedernal Mountain ..... 26
Penasco River ..... 28
Pennsylvanian ..... 5, 6, 7, 8, 10, 14, 18, 22, 23, 24, 25, 26, 28, 29, 30, 31, 37
Peritrochia erebus ..... 11
Permian ..... 5, 6, 8, 9, 10, 11, 12, 13, 14, 16, 17, 18, 19, 20, 21 22, 23, 24, 25, 26, 27, 29, 30, 31, 37
near Las Delicias, Coahuila, Mexico ..... 18
Arctic America ..... 33
Canadian Rockies ..... 33
Carrizo and Wylie Mountains ..... 17
Diablo Plateau ..... 15
Kansas ..... 22
Ellesmere and Heiberg Islands ..... 34
Finlay Mountains ..... 17
Hueco and Franklin Mountains ..... 16
Mid-Continent Region ..... 18
North Central Texas ..... 18
Northern California ..... 33
Oklahoma and Texas Panhandle ..... 21
Shafter District ..... 17
southeastern and central Utah and southwestern Colorado 30
Perrinites ..... 9, 11, 20
compressus ..... 7, 8, 20
cumminsi ..... 7, 8, 19, 20
hilli ..... 20
vidriensis ..... 7, 8, 9, 17, 18, 20
Perry, Oklahoma ..... 21
Peru ..... 35, 36
Phosphoria ..... 10, 27, 30, 31, 32, 33, 35, 37
Pinto Canyon ..... 17
Plagioglypta canna ..... 33
Pleurophorus ..... 10
Plummer, F. B. ..... 6, 18, 19, 20, 23, 37
Poikilosakos ..... 40
Pope, Captain ..... 5
Pre-Cambrian ..... 14, 26, 29
Pre-Comanchean ..... 15
Pre-Triassic ..... 29
Pre-Wichita ..... 26
Productidae ..... 65
Productus ..... 25, 38, 75
andii ..... 58
aff. P. semireticulatus ..... 24
articus ..... 34, 35, 66
borealis ..... 33
capitanensis ..... 66
cf. abichi ..... 33
cf. spiralis ..... 33
comancheanus ..... 67
cora ..... 75
dartoni ..... 67
geniculatus ..... 31, 32
gratiosus occidentalis ..... 71
guadalupensis ..... 39, 67
guadalupensis var. comancheanus ..... 11
hessensis ..... 16, 17, 68
horridus ..... 39, 85
huecoensis ..... 16, 68
humboldti ..... 39
inca ..... 15, 16, 24, 25, 27, 36, 38, 68
indicus ..... 72
inflatus ..... 34

| | Page |
|---|---|
| invesi | 9, 15, 17, 25, 26, 27, 28, 29, 30, 33, 69 |
| leei | 25 |
| leonardensis | 9, 11, 70 |
| mexicanus | 27 |
| multistriatus | 31, 32, 71 |
| nebraskensis | 26 |
| occidentalis | 15, 25, 27, 30, 72 |
| pustulosa | 79 |
| schucherti | 73 |
| semireticulatus | 24, 33 |
| semireticulatus hermosanus | 73 |
| semistriatus | 16, 74 |
| simbriatus | 82 |
| sulcatus var. borealis | 33 |
| timanicus | 33, 39 |
| wolfcampensis | 16, 74 |
| youngianus | 82 |
| Prommel, H. W. C. | 30 |
| Prorichthofenia | 7, 8, 9, 10, 40 |
| likharewi | 98 |
| novum | 97 |
| permiana | 99 |
| teguliferoides | 8, 14, 17, 39, 99 |
| uddeni | 100 |
| Prothalassoceras | 8 |
| welleri | 7, 8 |
| Prouddenites fauna | 6 |
| Pseudosyrinx | 41 |
| Ptychospira | 41 |
| Pugnoides | 40 |
| bidentatus | 11, 107 |
| elegans | 106 |
| osagensis | 24, 40 |
| osagensis pusilla | 26 |
| pingus | 106 |
| shumardianus | 107 |
| swallovianus | 31, 32, 107 |
| texanus | 9, 11, 16, 24, 26, 108 |
| transversus | 108 |
| utah | 36 |
| Punctospirifer *billingsii* | 124 |
| kentuckyensis | 125 |
| Pustula | 39, 79 |
| nevadaensis | 30, 31, 32 |
| aff. P. porrecta | 25 |
| semipunctata | 39, 79 |
| subhorrida var. rugatula? | 11 |
| Putnam formation | 18, 23 |
| Quadrant | 31 |
| Quanah, Texas | 20 |
| Quartermaster formation | 21 |
| *Queen member* | 13, 27 |
| Red River | 19, 20 |
| Redwall | 28, 29 |
| Reeside, J. B. | 11, 13, 27, 28, 29, 30 |
| Rex chert member | 31 |
| Rhipidomella | 37 |
| cargonaria | 43 |
| hessensis | 37, 43 |
| leonardensis | 43 |
| mesoplatys | 44 |
| Oehlert | 43 |
| subquadrata | 37 |
| transversa | 44 |
| Rhynchonella guadalupae | 112 |
| Rhynchonellidae | 40 |
| Rhyncopora | 40 |
| Rhyncopora illinoisensis | 40, 109 |
| illinoisensis R. nikitini | 40 |
| taylori | 40, 109 |
| Richardson, G. B. | 5, 14, 28 |
| Richthofenidae | 39, 96 |
| Rico formation | 30 |
| Road Canyon | 10 |
| Rocky Arroyo | 28 |
| Rocky Mountains | 30 |
| Romer, A. S. | 19, 20, 23, 26 |
| Rowleyella | 41 |
| Ruedemann, R. | 13 |
| Runnels County | 20 |
| Russel Gap | 28 |
| Russia | 31, 35 |
| Rustler limestone | 13, 22 |
| Sacramento Mountains | 27, 28 |
| Salt Basin | 12, 14, 15 |
| Salt Crotan Creek | 20 |
| Salt Flat | 15 |
| Salt Plain | 23 |
| Salt Range | 36, 38, 41 |
| San Angelo formation | 20, 21, 23 |
| San Andres | 9, 13, 21, 25, 26, 27, 28, 30 |
| San Rafael Swell | 30, 32 |
| San Rita Mountains | 25 |
| Sawmill Canyon | 25 |
| Scacchinella | 39 |
| gigantea | 6, 7, 8, 91 |
| Schizophoria | 37 |
| Schwagerina | 7, 16, 17, 24, 31, 35 |
| kansasensis | 22 |
| princeps | 35 |
| robusta | 33 |
| Schellwien, E. | 55 |
| Schuchert, Charles | 5, 6, 22, 25, 35 |
| Scott, Gayle | 19 |
| Sellards, E. H. | 5 |
| Seven Heart Gap | 11, 12, 13 |
| Seven Rivers member | 13, 27 |
| Seymouria | 17, 20 |
| Shafter | 17, 18 |
| Shafter Mine | 17 |
| Shumard, B. F. | 5, 37 |
| Shumard, George C. | 5 |
| Sierra del Sobaco | 18 |
| Sierra *Diablo* | 8, 11, 12, 14, 15, 17, 18 |
| Smith, I. P. | 6 |
| Snyder's Hill | 25 |
| Sonora | 25 |
| Sosio limestone | 8, 38, 39 |
| Spehenacodon | 26 |
| Sphenophyllum | 23 |
| Sphenopteris | 26 |
| Spirifer limestone | 35 |
| Spirifer ambiguus | 128 |
| bakeri | 115 |
| condor | 6, 7, 16, 36, 41, 113 |
| costella | 41, 115 |
| fasciger | 41 |
| huecoensis | 16 |
| *lyra* | 41 |
| marcoui | 41 |
| marcoui infraplica | 15, 41, 114 |
| mexicanus | 41 |
| mexicanus latus | 116 |
| oldhaminus | 41 |
| pseudocameratus | 15, 31, 32, 41 |
| pseudocameratus | 116 |
| rajah | 41 |
| rockymontanus | 36 |
| Sowerby | 113 |
| sulcifer | 12, 41, 118 |
| supramosquensis | 34, 35 |
| texanus | 117 |
| triplicatus | 117 |
| Spiriferella | 41 |
| Spiriferidae | 113 |
| Spiriferina angulata | 122 |
| campestris | 36 |
| d'Orbigny | 121 |
| hilli | 122 |
| laxa | 12, 122 |
| pulchra | 31, 32 |
| pyramidalis | 123 |
| Spirigerella | 41 |
| Spitzbergen | 32, 34, 35 |
| Squamularia *guadalupensis* | 10, 25, 27, 41, 118 |
| gemmellaro | 118 |
| indica | 41 |
| perplexa | 27, 41, 119 |
| Stacheoceras | 20, 33 |
| bowmani | 10 |
| gilli amense | 17 |
| Steinmann, ——— | 35 |
| Stenopoceras dumblei | 23 |
| Stillwater formation | 21, 23 |
| Store Bjornekap | 34 |
| Stoyanow, A. A. | 25 |
| Straparollus subquadratus | 25 |
| pernodosus | 25 |
| Stratford formation | 23 |
| Streptorhynchus | 38 |
| lamellatus | 38, 49 |
| pygmaeum | 49 |
| pyramidalis | 38, 50 |
| undulatus | 8, 51 |
| Striatifera | 39, 78 |
| pinniformis | 78 |
| Strophalosia | 39 |
| hustricula | 96 |
| Supai formation | 28, 29, 30 |
| Svarte Veg | 34 |
| *Syringothyris* | 41 |
| Taeniopteris | 23, 26 |
| Tank Canyon | 28 |
| Tectarea | 40 |
| Teguliferina | 8, 40 |
| bosei | 7, 40, 96 |
| deformis | 40 |
| schellwien | 96 |
| Tensleep sandstone | 31 |
| Terebratula geinitziana | 109 |
| michelini | 43 |
| schlotheimi | 110 |
| Terebratulidae | 42, 131 |
| Terebratuloida | 40 |
| Tessey | 10 |
| Tethyan seaways | 37 |
| Texas | 23, 24, 25, 26, 27, 28, 29 |
| Texas Bureau of Economic Geology | 5 |

PAGE

Texas Company ..... 5
Timan ranges ..... 35
Timor, Island of ..... 36
Tombstone, Arizona ..... 25
Tom Green County ..... 20
Tomlinson, C. W. ..... 23
Trans-Pecos Texas ..... 5, 6, 10, 11, 13, 16, 17, 18, 21, 22, 27, 29, 30, 32, 34, 35, 36, 37
Triassic ..... 21, 32, 33
Trinity sand ..... 21
Triticites ..... 22
  cullomensis ..... 6
  secalicus ..... 6
  ventricosus ..... 22
Trogkofel ..... 7, 8, 38, 39, 40
Tschernyschew, T. ..... 38
Tschernyschewia ..... 39
Tucson, Arizona ..... 25
Tularosa ..... 26
Udden, J. A. ..... 17
Uddenites zone ..... 6, 7, 17, 22, 23, 36
Uinta Mountains ..... 31
Illmania ..... 23
Uncinella ..... 41
Uncinuloides ..... 40
  guadalupensis ..... 112
  novum ..... 112
Ural Mountains ..... 35, 36
United States Geological Survey ..... 29
Ussuriland ..... 32
Utah ..... 27, 30, 31
Vale formation ..... 20
Van Horn, Texas ..... 17
Vanoos formation ..... 23
Victoria Island ..... 33
Victorio Peak ..... 11, 14, 15, 28
Vidrio dolomite ..... 10
Voltzia ..... 23

PAGE

Waagenoceras ..... 12, 17, 18, 20
  dieneri ..... 10, 18
Waagenoconcha ..... 36, 39, 80
  irginae ..... 36
  leonardensis ..... 80
  montpelierensis ..... 6, 7, 15, 30, 31, 32, 33, 36, 81
Wabaunsee ..... 22
Walchia ..... 19, 23, 26, 29
Walchia cf. W. gracilia ..... 29
Walchia pinniformis ..... 23
Waterman Mountains ..... 25
Weber quartzite ..... 31
Weller, S. ..... 37
Wellington formation ..... 21, 23
Wells formation ..... 31
White, C. A. ..... 19
White, David ..... 11, 19, 22, 23, 26
Whitehorse ..... 21, 22
Wichita-Clear Fork boundary ..... 20
Wichita ..... 18, 19, 21, 22, 23, 28
Wichita Mountains ..... 21
Wichita River Valley ..... 19
Wildwood limestone ..... 33
Willis, Robin ..... 21, 27, 28
Winkelman, Texas ..... 25
Wolfcamp formation ..... 6, 7, 8, 10, 14, 16, 17, 19, 20, 22, 26, 28, 29, 38, 40
Word formation ..... 5, 9, 10, 11, 12, 13, 17, 18, 20, 22, 27, 29, 40
Word Ranch ..... 9, 17
Wreford ..... 23
Wylie Mountains ..... 17
Wyoming ..... 30, 31
Yakowlew, N. ..... 55
Yale University ..... 5
Yeso formation ..... 25, 26, 28, 30
Yukon ..... 33
Zechstein fauna ..... 41
Zuni Mountains ..... 26
Zuni uplift ..... 29